

PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁷ : C12N 15/16, C07K 14/72	A2	(11) International Publication Number: WO 00/22131
		(43) International Publication Date: 20 April 2000 (20.04.00)

(21) International Application Number: PCT/US99/24065

(22) International Filing Date: 13 October 1999 (13.10.99)

(30) Priority Data:

09/170,496	13 October 1998 (13.10.98)	US
60/108,029	12 November 1998 (12.11.98)	US
60/109,213	20 November 1998 (20.11.98)	US
60/110,060	27 November 1998 (27.11.98)	US
60/120,416	16 February 1999 (16.02.99)	US
60/121,852	26 February 1999 (26.02.99)	US
60/123,944	12 March 1999 (12.03.99)	US
60/123,945	12 March 1999 (12.03.99)	US
60/123,948	12 March 1999 (12.03.99)	US
60/123,946	12 March 1999 (12.03.99)	US
60/123,949	12 March 1999 (12.03.99)	US
60/123,951	12 March 1999 (12.03.99)	US
60/136,436	28 May 1999 (28.05.99)	US
60/136,437	28 May 1999 (28.05.99)	US
60/136,439	28 May 1999 (28.05.99)	US
60/137,567	28 May 1999 (28.05.99)	US
60/137,127	28 May 1999 (28.05.99)	US
60/137,131	28 May 1999 (28.05.99)	US
60/141,448	30 June 1999 (30.06.99)	US
60/151,114	27 August 1999 (27.08.99)	US
60/152,524	3 September 1999 (03.09.99)	US
Not furnished	9 September 1999 (09.09.99)	US
60/156,633	29 September 1999 (29.09.99)	US
60/156,555	29 September 1999 (29.09.99)	US
60/156,634	29 September 1999 (29.09.99)	US
Not furnished	1 October 1999 (01.10.99)	US
Not furnished	1 October 1999 (01.10.99)	US
Not furnished	1 October 1999 (01.10.99)	US
Not furnished	1 October 1999 (01.10.99)	US
Not furnished	12 October 1999 (12.10.99)	US
Not furnished	12 October 1999 (12.10.99)	US

(63) Related by Continuation (CON) or Continuation-in-Part (CIP) to Earlier Application
US 09/170,496 (CIP)
Filed on 13 October 1998 (13.10.98)

(72) Inventors; and

(75) Inventors/Applicants (for US only): BEHAN, Dominic, P. [GB/US]; 11472 Roxboro Court, San Diego, CA 92131 (US). LEHMANN-BRUIINSMA, Karin [DE/US]; 12565 Pathos Lane, San Diego, CA 92129 (US). CHALMERS, Derek, T. [GB/US]; 347 Longden Lane, Solana Beach, CA 92150 (US). CHEN, Ruoping [CN/US]; 5296 Timber Branch Way, San Diego, CA 92130 (US). DANG, Huong, T. [US/US]; 5352 Oak Park Drive, San Diego, CA 92105 (US). GORE, Martin [GB/US]; 6868 Estrella Avenue, San Diego, CA 92120 (US). LIAW, Chen, W. [US/US]; 7668 Salix Place, San Diego, CA 92129 (US). LIN, I-Lin [-/US]; 8291-7 Gold Coast Drive, San Diego, CA 92126 (US). LOWITZ, Kevin [US/US]; Apartment C, 8031 Caminito de Pizza, San Diego, CA 92108 (US). WHITE, Carol [US/US]; 4260 Cleveland Avenue, San Diego, CA 92103 (US).

(74) Agents: MILLER, Suzanne, E. et al.; Woodcock Washburn Kurtz Mackiewicz & Norris LLP, 46th floor, One Liberty Place, Philadelphia, PA 19103 (US).

(81) Designated States: AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DE, DK, DM, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

Published

Without international search report and to be republished upon receipt of that report.

(54) Title: NON-ENDOGENOUS, CONSTITUTIVELY ACTIVATED HUMAN G PROTEIN-COUPLED RECEPTORS

(57) Abstract

The invention disclosed in this patent document relates to transmembrane receptors, more particularly to a human G protein-coupled receptor for which the endogenous ligand is unknown ("orphan GPCR receptors"), and most particularly to mutated (non-endogenous) versions of the human GPCRs for evidence of constitutive activity.

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav	TM	Turkmenistan
BF	Burkina Faso	GR	Greece		Republic of Macedonia	TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's	NZ	New Zealand		
CM	Cameroon		Republic of Korea	PL	Poland		
CN	China	KR	Republic of Korea	PT	Portugal		
CU	Cuba	KZ	Kazakhstan	RO	Romania		
CZ	Czech Republic	LC	Saint Lucia	RU	Russian Federation		
DE	Germany	LI	Liechtenstein	SD	Sudan		
DK	Denmark	LK	Sri Lanka	SE	Sweden		
EE	Estonia	LR	Liberia	SG	Singapore		

**NON-ENDOGENOUS, CONSTITUTIVELY ACTIVATED
HUMAN G PROTEIN-COUPLED RECEPTORS**

This patent application is a continuation-in-part of, and claims priority from, U.S. Serial Number 09/170,496, filed with the United States Patent and Trademark Office on

5 October 13, 1998. This application also claims the benefit of priority from the following provisional applications, all filed via U.S. Express Mail with the United States Patent and Trademark Office on the indicated dates: U.S. Provisional Number 60/110,060, filed November 27, 1998; U.S. Provisional Number 60/120,416, filed February 16, 1999; U.S. Provisional Number 60/121,852, filed February 26, 1999 claiming benefit of U.S.

10 Provisional Number 60/109,213, filed November 20, 1998; U.S. Provisional Number 60/123,944, filed March 12, 1999; U.S. Provisional Number 60/123,945, filed March 12, 1999; U.S. Provisional Number 60/123,948, filed March 12, 1999; U.S. Provisional Number 60/123,951, filed March 12, 1999; U.S. Provisional Number 60/123,946, filed March 12, 1999; U.S. Provisional Number 60/123,949, filed March 12, 1999; U.S.

15 Provisional Number 60/152,524, filed September 3, 1999, claiming benefit of U.S. Provisional Number 60/151,114, filed August 27, 1999 and U.S. Provisional Number 60/108,029, filed November 12, 1998; U.S. Provisional Number 60/136,436, filed May 28, 1999; U.S. Provisional Number 60/136,439, filed May 28, 1999; U.S. Provisional Number 60/136,567, filed May 28, 1999; U.S. Provisional Number 60/137,127, filed May 28,

20 1999; U.S. Provisional Number 60/137,131, filed May 28, 1999; U.S. Provisional Number

- 2 -

60/141,448, filed June 29, 1999 claiming benefit of U.S. Provisional Number 60/136,437, filed May 28, 1999; U.S. Provisional Number 60/156,633, filed September 29, 1999; U.S. Provisional Number 60/156,555, filed September 29, 1999; U.S. Provisional Number 60/156,634, filed September 29, 1999; U.S. Provisional Number ____ (Arena Pharmaceuticals, Inc. docket number: CHN10-1), filed September 29, 1999; U.S. Provisional Number ____ (Arena Pharmaceuticals, Inc. docket number: RUP6-1), filed October 1, 1999; U.S. Provisional Number ____ (Arena Pharmaceuticals, Inc. docket number: RUP7-1), filed October 1, 1999; U.S. Provisional Number ____ (Arena Pharmaceuticals, Inc. docket number: CHN6-1), filed October 1, 1999; U.S. Provisional Number ____ (Arena Pharmaceuticals, Inc. docket number: RUP5-1), filed October 1, 1999; and U.S. Provisional Number ____ (Arena Pharmaceuticals, Inc. docket number: CHN9-1), filed October 1, 1999. This application is also related to co-pending U.S. Serial Number ____ (Woodcock, Washburn, Kurtz, Makiewicz & Norris, LLP docket number AREN-0050), filed on October 12, 1999 (via U.S. Express Mail) and U.S. Serial Number 09/364,425, filed on July 30, 1999, both incorporated herein by reference. This application also claims priority to U.S. Serial Number ____ (Woodcock, Washburn, Kurtz, Makiewicz & Norris, LLP docket number AREN-0054), filed on October 12, 1999 (via U.S. Express Mail), incorporated by reference herein in its entirety. Each of the foregoing applications are incorporated by reference herein in their entirety.

20

FIELD OF THE INVENTION

The invention disclosed in this patent document relates to transmembrane receptors, and more particularly to human G protein-coupled receptors, and specifically to

- 3 -

GPCRs that have been altered to establish or enhance constitutive activity of the receptor. Preferably, the altered GPCRs are used for the direct identification of candidate compounds as receptor agonists, inverse agonists or partial agonists having potential applicability as therapeutic agents.

BACKGROUND OF THE INVENTION

Although a number of receptor classes exist in humans, by far the most abundant and therapeutically relevant is represented by the G protein-coupled receptor (GPCR or GPCRs) class. It is estimated that there are some 100,000 genes within the human genome, and of these, approximately 2%, or 2,000 genes, are estimated to code for GPCRs. Receptors, including GPCRs, for which the endogenous ligand has been identified are referred to as "known" receptors, while receptors for which the endogenous ligand has not been identified are referred to as "orphan" receptors. GPCRs represent an important area for the development of pharmaceutical products: from approximately 20 of the 100 known GPCRs, 60% of all prescription pharmaceuticals have been developed.

GPCRs share a common structural motif. All these receptors have seven sequences of between 22 to 24 hydrophobic amino acids that form seven alpha helices, each of which spans the membrane (each span is identified by number, *i.e.*, transmembrane-1 (TM-1), transmembrane-2 (TM-2), etc.). The transmembrane helices are joined by strands of amino acids between transmembrane-2 and transmembrane-3, transmembrane-4 and transmembrane-5, and transmembrane-6 and transmembrane-7 on the exterior, or "extracellular" side, of the cell membrane (these are referred to as "extracellular" regions 1, 2 and 3 (EC-1, EC-2 and EC-3), respectively). The transmembrane helices are also joined by strands of amino acids between transmembrane-1 and transmembrane-2, transmembrane-3 and transmembrane-4, and

- 4 -

transmembrane-5 and transmembrane-6 on the interior, or "intracellular" side, of the cell membrane (these are referred to as "intracellular" regions 1, 2 and 3 (IC-1, IC-2 and IC-3), respectively). The "carboxy" ("C") terminus of the receptor lies in the intracellular space within the cell, and the "amino" ("N") terminus of the receptor lies in the extracellular space
5 outside of the cell.

Generally, when an endogenous ligand binds with the receptor (often referred to as "activation" of the receptor), there is a change in the conformation of the intracellular region that allows for coupling between the intracellular region and an intracellular "G-protein." It has been reported that GPCRs are "promiscuous" with respect to G proteins, *i.e.*,
10 that a GPCR can interact with more than one G protein. *See*, Kenakin, T., 43 *Life Sciences* 1095 (1988). Although other G proteins exist, currently, Gq, Gs, Gi, Gz and Go are G proteins that have been identified. Endogenous ligand-activated GPCR coupling with the G-protein begins a signaling cascade process (referred to as "signal transduction"). Under normal conditions, signal transduction ultimately results in cellular activation or cellular inhibition.
15 It is thought that the IC-3 loop as well as the carboxy terminus of the receptor interact with the G protein.

Under physiological conditions, GPCRs exist in the cell membrane in equilibrium between two different conformations: an "inactive" state and an "active" state. A receptor in an inactive state is unable to link to the intracellular signaling transduction
20 pathway to produce a biological response. Changing the receptor conformation to the active state allows linkage to the transduction pathway (via the G-protein) and produces a biological response.

A receptor may be stabilized in an active state by an endogenous ligand or a

- 5 -

compound such as a drug. Recent discoveries, including but not exclusively limited to modifications to the amino acid sequence of the receptor, provide means other than endogenous ligands or drugs to promote and stabilize the receptor in the active state conformation. These means effectively stabilize the receptor in an active state by
5 simulating the effect of an endogenous ligand binding to the receptor. Stabilization by such ligand-independent means is termed "constitutive receptor activation."

SUMMARY OF THE INVENTION

Disclosed herein are non-endogenous versions of endogenous, human GPCRs and uses thereof.

10

BRIEF DESCRIPTION OF THE DRAWINGS

Figure 1 is a representation of 8XCRE-Luc reporter plasmid (*see*, Example 4(c)3.)

Figures 2A and 2B are graphic representations of the results of ATP and ADP binding to endogenous TDAG8 (2A) and comparisons in serum and serum free media (2B).

15

Figure 3 is a graphic representation of the comparative signaling results of CMV versus the GPCR Fusion Protein H9(F236K):Gsa.

DETAILED DESCRIPTION

The scientific literature that has evolved around receptors has adopted a number of terms to refer to ligands having various effects on receptors. For clarity and
20 consistency, the following definitions will be used throughout this patent document. To the extent that these definitions conflict with other definitions for these terms, the following definitions shall control:

AGONISTS shall mean materials (*e.g.*, ligands, candidate compounds) that

activate the intracellular response when they bind to the receptor, or enhance GTP binding to membranes.

AMINO ACID ABBREVIATIONS used herein are set out in Table A:

TABLE A		
5	ALANINE	ALA
	ARGININE	ARG
	ASPARAGINE	ASN
	ASPARTIC ACID	ASP
	CYSTEINE	CYS
10	GLUTAMIC ACID	GLU
	GLUTAMINE	GLN
	GLYCINE	GLY
	HISTIDINE	HIS
	ISOLEUCINE	ILE
15	LEUCINE	LEU
	LYSINE	LYS
	METHIONINE	MET
	PHENYLALANINE	PHE
	PROLINE	PRO
20	SERINE	SER
	THREONINE	THR
	TRYPTOPHAN	TRP
	TYROSINE	TYR
	VALINE	VAL

25 **PARTIAL AGONISTS** shall mean materials (*e.g.*, ligands, candidate compounds) that activate the intracellular response when they bind to the receptor to a lesser degree/extent than do agonists, or enhance GTP binding to membranes to a lesser degree/extent than do agonists.

ANTAGONIST shall mean materials (*e.g.*, ligands, candidate compounds) that
 30 competitively bind to the receptor at the same site as the agonists but which do not activate the intracellular response initiated by the active form of the receptor, and can thereby inhibit the intracellular responses by agonists or partial agonists. **ANTAGONISTS** do not diminish the baseline intracellular response in the absence of an agonist or partial agonist.

CANDIDATE COMPOUND shall mean a molecule (for example, and not limitation,

- 7 -

a chemical compound) that is amenable to a screening technique. Preferably, the phrase "candidate compound" does not include compounds which were publicly known to be compounds selected from the group consisting of inverse agonist, agonist or antagonist to a receptor, as previously determined by an indirect identification process ("indirectly identified compound"); more preferably, not including an indirectly identified compound which has previously been determined to have therapeutic efficacy in at least one mammal; and, most preferably, not including an indirectly identified compound which has previously been determined to have therapeutic utility in humans.

COMPOSITION means a material comprising at least one component; a "pharmaceutical composition" is an example of a composition.

COMPOUND EFFICACY shall mean a measurement of the ability of a compound to inhibit or stimulate receptor functionality, as opposed to receptor binding affinity. Exemplary means of detecting compound efficacy are disclosed in the Example section of this patent document.

CODON shall mean a grouping of three nucleotides (or equivalents to nucleotides) which generally comprise a nucleoside (adenosine (A), guanosine (G), cytidine (C), uridine (U) and thymidine (T)) coupled to a phosphate group and which, when translated, encodes an amino acid.

CONSTITUTIVELY ACTIVATED RECEPTOR shall mean a receptor subject to constitutive receptor activation. A constitutively activated receptor can be endogenous or non-endogenous.

CONSTITUTIVE RECEPTOR ACTIVATION shall mean stabilization of a receptor in the active state by means other than binding of the receptor with its endogenous

- 8 -

ligand or a chemical equivalent thereof.

CONTACT or **CONTACTING** shall mean bringing at least two moieties together, whether in an in vitro system or an in vivo system.

DIRECTLY IDENTIFYING or **DIRECTLY IDENTIFIED**, in relationship to the phrase "candidate compound", shall mean the screening of a candidate compound against a constitutively activated receptor, preferably a constitutively activated orphan receptor, and most preferably against a constitutively activated G-protein-coupled cell surface orphan receptor, and assessing the compound efficacy of such compound. This phrase is, under no circumstances, to be interpreted or understood to be encompassed by or to encompass the phrase "indirectly identifying" or "indirectly identified."

ENDOGENOUS shall mean a material that a mammal naturally produces. **ENDOGENOUS** in reference to, for example and not limitation, the term "receptor," shall mean that which is naturally produced by a mammal (for example, and not limitation, a human) or a virus. By contrast, the term **NON-ENDOGENOUS** in this context shall mean that which is not naturally produced by a mammal (for example, and not limitation, a human) or a virus. For example, and not limitation, a receptor which is not constitutively active in its endogenous form, but when manipulated becomes constitutively active, is most preferably referred to herein as a "non-endogenous, constitutively activated receptor." Both terms can be utilized to describe both "in vivo" and "in vitro" systems. For example, and not limitation, in a screening approach, the endogenous or non-endogenous receptor may be in reference to an in vitro screening system. As a further example and not limitation, where the genome of a mammal has been manipulated to include a non-endogenous constitutively activated receptor, screening of a candidate compound by means of an in vivo system is viable.

- 9 -

G PROTEIN COUPLED RECEPTOR FUSION PROTEIN and GPCR FUSION

PROTEIN, in the context of the invention disclosed herein, each mean a non-endogenous protein comprising an endogenous, constitutively activate GPCR or a non-endogenous, constitutively activated GPCR fused to at least one G protein, most preferably the alpha (α) subunit of such G protein (this being the subunit that binds GTP), with the G protein preferably being of the same type as the G protein that naturally couples with endogenous orphan GPCR. For example, and not limitation, in an endogenous state, if the G protein "G α " is the predominate G protein that couples with the GPCR, a GPCR Fusion Protein based upon the specific GPCR would be a non-endogenous protein comprising the GPCR fused to G α ; in some circumstances, as will be set forth below, a non-predominant G protein can be fused to the GPCR. The G protein can be fused directly to the c-terminus of the constitutively active GPCR or there may be spacers between the two.

HOST CELL shall mean a cell capable of having a Plasmid and/or Vector incorporated therein. In the case of a prokaryotic Host Cell, a Plasmid is typically replicated as a autonomous molecule as the Host Cell replicates (generally, the Plasmid is thereafter isolated for introduction into a eukaryotic Host Cell); in the case of a eukaryotic Host Cell, a Plasmid is integrated into the cellular DNA of the Host Cell such that when the eukaryotic Host Cell replicates, the Plasmid replicates. Preferably, for the purposes of the invention disclosed herein, the Host Cell is eukaryotic, more preferably, mammalian, and most preferably selected from the group consisting of 293, 293T and COS-7 cells.

INDIRECTLY IDENTIFYING or **INDIRECTLY IDENTIFIED** means the traditional approach to the drug discovery process involving identification of an endogenous ligand specific for an endogenous receptor, screening of candidate compounds against the

- 10 -

receptor for determination of those which interfere and/or compete with the ligand-receptor interaction, and assessing the efficacy of the compound for affecting at least one second messenger pathway associated with the activated receptor.

INHIBIT or **INHIBITING**, in relationship to the term "response" shall mean that a response is decreased or prevented in the presence of a compound as opposed to in the absence of the compound.

INVERSE AGONISTS shall mean materials (*e.g.*, ligand, candidate compound) which bind to either the endogenous form of the receptor or to the constitutively activated form of the receptor, and which inhibit the baseline intracellular response initiated by the active form of the receptor below the normal base level of activity which is observed in the absence of agonists or partial agonists, or decrease GTP binding to membranes. Preferably, the baseline intracellular response is inhibited in the presence of the inverse agonist by at least 30%, more preferably by at least 50%, and most preferably by at least 75%, as compared with the baseline response in the absence of the inverse agonist.

KNOWN RECEPTOR shall mean an endogenous receptor for which the endogenous ligand specific for that receptor has been identified.

LIGAND shall mean an endogenous, naturally occurring molecule specific for an endogenous, naturally occurring receptor.

MUTANT or **MUTATION** in reference to an endogenous receptor's nucleic acid and/or amino acid sequence shall mean a specified change or changes to such endogenous sequences such that a mutated form of an endogenous, non-constitutively activated receptor evidences constitutive activation of the receptor. In terms of equivalents to specific sequences, a subsequent mutated form of a human receptor is considered to be equivalent to

- 11 -

a first mutation of the human receptor if (a) the level of constitutive activation of the subsequent mutated form of a human receptor is substantially the same as that evidenced by the first mutation of the receptor; and (b) the percent sequence (amino acid and/or nucleic acid) homology between the subsequent mutated form of the receptor and the first mutation
5 of the receptor is at least about 80%, more preferably at least about 90% and most preferably at least 95%. Ideally, and owing to the fact that the most preferred cassettes disclosed herein for achieving constitutive activation includes a single amino acid and/or codon change between the endogenous and the non-endogenous forms of the GPCR, the percent sequence homology should be at least 98%.

10 **NON-ORPHAN RECEPTOR** shall mean an endogenous naturally occurring molecule specific for an endogenous naturally occurring ligand wherein the binding of a ligand to a receptor activates an intracellular signaling pathway.

ORPHAN RECEPTOR shall mean an endogenous receptor for which the endogenous ligand specific for that receptor has not been identified or is not known.

15 **PHARMACEUTICAL COMPOSITION** shall mean a composition comprising at least one active ingredient, whereby the composition is amenable to investigation for a specified, efficacious outcome in a mammal (for example, and not limitation, a human). Those of ordinary skill in the art will understand and appreciate the techniques appropriate for determining whether an active ingredient has a desired efficacious outcome based upon the
20 needs of the artisan.

PLASMID shall mean the combination of a Vector and cDNA. Generally, a Plasmid is introduced into a Host Cell for the purposes of replication and/or expression of the cDNA as a protein.

- 12 -

STIMULATE or **STIMULATING**, in relationship to the term "response" shall mean that a response is increased in the presence of a compound as opposed to in the absence of the compound.

VECTOR in reference to cDNA shall mean a circular DNA capable of incorporating
5 at least one cDNA and capable of incorporation into a Host Cell.

The order of the following sections is set forth for presentational efficiency and is not intended, nor should be construed, as a limitation on the disclosure or the claims to follow.

A. Introduction

The traditional study of receptors has always proceeded from the a priori assumption
10 (historically based) that the endogenous ligand must first be identified before discovery could proceed to find antagonists and other molecules that could affect the receptor. Even in cases where an antagonist might have been known first, the search immediately extended to looking for the endogenous ligand. This mode of thinking has persisted in receptor research even after the discovery of constitutively activated receptors. What has not been heretofore recognized
15 is that it is the active state of the receptor that is most useful for discovering agonists, partial agonists, and inverse agonists of the receptor. For those diseases which result from an overly active receptor or an under-active receptor, what is desired in a therapeutic drug is a compound which acts to diminish the active state of a receptor or enhance the activity of the receptor, respectively, not necessarily a drug which is an antagonist to the endogenous ligand.
20 This is because a compound that reduces or enhances the activity of the active receptor state need not bind at the same site as the endogenous ligand. Thus, as taught by a method of this invention, any search for therapeutic compounds should start by screening compounds against the ligand-independent active state.

- 13 -

B. Identification of Human GPCRs

The efforts of the Human Genome project has led to the identification of a plethora of information regarding nucleic acid sequences located within the human genome; it has been the case in this endeavor that genetic sequence information has been made available without an understanding or recognition as to whether or not any particular genomic sequence does or may contain open-reading frame information that translate human proteins. Several methods of identifying nucleic acid sequences within the human genome are within the purview of those having ordinary skill in the art. For example, and not limitation, a variety of human GPCRs, disclosed herein, were discovered by reviewing the GenBank™ database, while other GPCRs were discovered by utilizing a nucleic acid sequence of a GPCR, previously sequenced, to conduct a BLAST™ search of the EST database. Table B, below, lists several endogenous GPCRs that we have discovered, along with a GPCR's respective homologous receptor.

TABLE B

	Disclosed Human Orphan GPCRs	Accession Number Identified	Open Reading Frame (Base Pairs)	Per Cent Homology To Designated GPCR	Reference To Homologous GPCR (Accession No.)
15					
	hARE-3	AL033379	1,260 bp	52.3% LPA-R	U92642
20	hARE-4	AC006087	1,119 bp	36% P2Y5	AF000546
	hARE-5	AC006255	1,104 bp	32% <i>Oryzias latipes</i>	D43633
	hGPR27	AA775870	1,128 bp		
	hARE-1	AI090920	999 bp	43% KIAA0001	D13626
	hARE-2	AA359504	1,122 bp	53% GPR27	
25	hPPR1	H67224	1,053 bp	39% EBI1	L31581
	hG2A	AA754702	1,113 bp	31% GPR4	L36148

- 14 -

	hRUP3	AL035423	1,005 bp	30% <i>Drosophila melanogaster</i>	2133653
	hRUP4	AI307658	1,296 bp	32% pNPGPR 28% and 29 % <i>Zebra fish</i> Ya and Yb, respectively	NP_004876 AAC41276 and AAB94616
	hRUP5	AC005849	1,413 bp	25% DEZ 23% FMLPR	Q99788 P21462
5	hRUP6	AC005871	1,245 bp	48% GPR66	NP_006047
	hRUP7	AC007922	1,173 bp	43% H3R	AF140538
	hCHN3	EST 36581	1,113 bp	53% GPR27	
	hCHN4	AA804531	1,077 bp	32% thrombin	4503637
	hCHN6	EST 2134670	1,503 bp	36% edg-1	NP_001391
	hCHN8	EST 764455	1,029 bp	47% KIAA0001	D13626
10	hCHN9	EST 1541536	1,077 bp	41% LTB4R	NM_000752
	hCHN10	EST 1365839	1,055 bp	35% P2Y	NM_002563

Receptor homology is useful in terms of gaining an appreciation of a role of the receptors within the human body. As the patent document progresses, we will disclose techniques for mutating these receptors to establish non-endogenous, constitutively activated versions of these receptors.

The techniques disclosed herein have also been applied to other human, orphan GPCRs known to the art, as will be apparent as the patent document progresses.

C. Receptor Screening

Screening candidate compounds against a non-endogenous, constitutively activated version of the human GPCRs disclosed herein allows for the direct identification of candidate compounds which act at this cell surface receptor, without requiring use of the receptor's endogenous ligand. By determining areas within the body where the endogenous version of human GPCRs disclosed herein is expressed and/or over-expressed, it is possible to determine related disease/disorder states which are associated with the expression and/or over-expression

- 15 -

of the receptor; such an approach is disclosed in this patent document.

With respect to creation of a mutation that may evidence constitutive activation of the human GPCR disclosed herein is based upon the distance from the proline residue at which is presumed to be located within TM6 of the GPCR; this algorithmic technique is disclosed
5 in co-pending and commonly assigned patent document U.S. Serial Number 09/170,496, incorporated herein by reference. The algorithmic technique is not predicated upon traditional sequence "alignment" but rather a specified distance from the aforementioned TM6 proline residue. By mutating the amino acid residue located 16 amino acid residues from this residue (presumably located in the IC3 region of the receptor) to, most preferably, a lysine residue,
10 such activation may be obtained. Other amino acid residues may be useful in the mutation at this position to achieve this objective.

D. Disease/Disorder Identification and/or Selection

As will be set forth in greater detail below, most preferably inverse agonists to the non-endogenous, constitutively activated GPCR can be identified by the methodologies of this
15 invention. Such inverse agonists are ideal candidates as lead compounds in drug discovery programs for treating diseases related to this receptor. Because of the ability to directly identify inverse agonists to the GPCR, thereby allowing for the development of pharmaceutical compositions, a search for diseases and disorders associated with the GPCR is relevant. For example, scanning both diseased and normal tissue samples for the presence
20 of the GPCR now becomes more than an academic exercise or one which might be pursued along the path of identifying an endogenous ligand to the specific GPCR. Tissue scans can be conducted across a broad range of healthy and diseased tissues. Such tissue scans provide a preferred first step in associating a specific receptor with a disease and/or disorder. *See, for*

- 16 -

example, co-pending application (docket number ARE-0050) for exemplary dot-blot and RT-PCR results of several of the GPCRs disclosed herein.

Preferably, the DNA sequence of the human GPCR is used to make a probe for (a) dot-blot analysis against tissue-mRNA, and/or (b) RT-PCR identification of the expression
5 of the receptor in tissue samples. The presence of a receptor in a tissue source, or a diseased tissue, or the presence of the receptor at elevated concentrations in diseased tissue compared to a normal tissue, can be preferably utilized to identify a correlation with a treatment regimen, including but not limited to, a disease associated with that disease. Receptors can equally well be localized to regions of organs by this technique. Based on
10 the known functions of the specific tissues to which the receptor is localized, the putative functional role of the receptor can be deduced.

E. Screening of Candidate Compounds

1. Generic GPCR screening assay techniques

When a G protein receptor becomes constitutively active, it binds to a G protein (*e.g.*,
15 Gq, Gs, Gi, Gz, Go) and stimulates the binding of GTP to the G protein. The G protein then acts as a GTPase and slowly hydrolyzes the GTP to GDP, whereby the receptor, under normal conditions, becomes deactivated. However, constitutively activated receptors continue to exchange GDP to GTP. A non-hydrolyzable analog of GTP, [³⁵S]GTPγS, can be used to monitor enhanced binding to membranes which express constitutively activated receptors.
20 It is reported that [³⁵S]GTPγS can be used to monitor G protein coupling to membranes in the absence and presence of ligand. An example of this monitoring, among other examples well-known and available to those in the art, was reported by Traynor and Nahorski in 1995. The preferred use of this assay system is for initial screening of candidate compounds because the

- 17 -

system is generically applicable to all G protein-coupled receptors regardless of the particular G protein that interacts with the intracellular domain of the receptor.

2. Specific GPCR screening assay techniques

Once candidate compounds are identified using the "generic" G protein-coupled receptor assay (*i.e.*, an assay to select compounds that are agonists, partial agonists, or inverse agonists), further screening to confirm that the compounds have interacted at the receptor site is preferred. For example, a compound identified by the "generic" assay may not bind to the receptor, but may instead merely "uncouple" the G protein from the intracellular domain.

a. *Gs, Gz and Gi.*

10 Gs stimulates the enzyme adenylyl cyclase. Gi (and Gz and Go), on the other hand, inhibit this enzyme. Adenylyl cyclase catalyzes the conversion of ATP to cAMP; thus, constitutively activated GPCRs that couple the Gs protein are associated with increased cellular levels of cAMP. On the other hand, constitutively activated GPCRs that couple Gi (or Gz, Go) protein are associated with decreased cellular levels of cAMP. *See, generally,*

15 "Indirect Mechanisms of Synaptic Transmission," Chpt. 8, From Neuron To Brain (3rd Ed.) Nichols, J.G. et al eds. Sinauer Associates, Inc. (1992). Thus, assays that detect cAMP can be utilized to determine if a candidate compound is, *e.g.*, an inverse agonist to the receptor (*i.e.*, such a compound would decrease the levels of cAMP). A variety of approaches known

20 in the art for measuring cAMP can be utilized; a most preferred approach relies upon the use of anti-cAMP antibodies in an ELISA-based format. Another type of assay that can be utilized is a whole cell second messenger reporter system assay. Promoters on genes drive the expression of the proteins that a particular gene encodes. Cyclic AMP drives gene expression by promoting the binding of a cAMP-responsive DNA binding protein or

- 18 -

transcription factor (CREB) that then binds to the promoter at specific sites called cAMP response elements and drives the expression of the gene. Reporter systems can be constructed which have a promoter containing multiple cAMP response elements before the reporter gene, *e.g.*, β -galactosidase or luciferase. Thus, a constitutively activated Gs-linked receptor causes
5 the accumulation of cAMP that then activates the gene and expression of the reporter protein. The reporter protein such as β -galactosidase or luciferase can then be detected using standard biochemical assays (Chen et al. 1995).

b. Go and Gq.

10 Gq and Go are associated with activation of the enzyme phospholipase C, which in turn hydrolyzes the phospholipid PIP_2 , releasing two intracellular messengers: diacylglycerol (DAG) and inistol 1,4,5-triphoisphate (IP_3). Increased accumulation of IP_3 is associated with activation of Gq- and Go-associated receptors. *See, generally, "Indirect Mechanisms of Synaptic Transmission," Chpt. 8, From Neuron To Brain (3rd Ed.) Nichols,*
15 *J.G. et al eds. Sinauer Associates, Inc. (1992).* Assays that detect IP_3 accumulation can be utilized to determine if a candidate compound is, *e.g.*, an inverse agonist to a Gq- or Go-associated receptor (*i.e.*, such a compound would decrease the levels of IP_3). Gq-associated receptors can also been examined using an AP1 reporter assay in that Gq-dependent phospholipase C causes activation of genes containing AP1 elements; thus, activated Gq-
20 associated receptors will evidence an increase in the expression of such genes, whereby inverse agonists thereto will evidence a decrease in such expression, and agonists will evidence an increase in such expression. Commercially available assays for such detection are available.

- 19 -

3. GPCR Fusion Protein

The use of an endogenous, constitutively activate orphan GPCR or a non-endogenous, constitutively activated orphan GPCR, for use in screening of candidate compounds for the direct identification of inverse agonists, agonists and partial agonists provide an interesting screening challenge in that, by definition, the receptor is active even in the absence of an endogenous ligand bound thereto. Thus, in order to differentiate between, *e.g.*, the non-endogenous receptor in the presence of a candidate compound and the non-endogenous receptor in the absence of that compound, with an aim of such a differentiation to allow for an understanding as to whether such compound may be an inverse agonist, agonist, partial agonist or have no affect on such a receptor, it is preferred that an approach be utilized that can enhance such differentiation. A preferred approach is the use of a GPCR Fusion Protein.

Generally, once it is determined that a non-endogenous orphan GPCR has been constitutively activated using the assay techniques set forth above (as well as others), it is possible to determine the predominant G protein that couples with the endogenous GPCR. Coupling of the G protein to the GPCR provides a signaling pathway that can be assessed. Because it is most preferred that screening take place by use of a mammalian expression system, such a system will be expected to have endogenous G protein therein. Thus, by definition, in such a system, the non-endogenous, constitutively activated orphan GPCR will continuously signal. In this regard, it is preferred that this signal be enhanced such that in the presence of, *e.g.*, an inverse agonist to the receptor, it is more likely that it will be able to more readily differentiate, particularly in the context of screening, between the receptor when it is contacted with the inverse agonist.

The GPCR Fusion Protein is intended to enhance the efficacy of G protein coupling

with the non-endogenous GPCR. The GPCR Fusion Protein is preferred for screening with a non-endogenous, constitutively activated GPCR because such an approach increases the signal that is most preferably utilized in such screening techniques. This is important in facilitating a significant "signal to noise" ratio; such a significant ratio is import preferred for
5 the screening of candidate compounds as disclosed herein.

The construction of a construct useful for expression of a GPCR Fusion Protein is within the purview of those having ordinary skill in the art. Commercially available expression vectors and systems offer a variety of approaches that can fit the particular needs of an investigator. The criteria of importance for such a GPCR Fusion Protein construct is
10 that the endogenous GPCR sequence and the G protein sequence both be in-frame (preferably, the sequence for the endogenous GPCR is upstream of the G protein sequence) and that the "stop" codon of the GPCR must be deleted or replaced such that upon expression of the GPCR, the G protein can also be expressed. The GPCR can be linked directly to the G protein, or there can be spacer residues between the two (preferably, no more than about 12,
15 although this number can be readily ascertained by one of ordinary skill in the art). We have a preference (based upon convenience) of use of a spacer in that some restriction sites that are not used will, effectively, upon expression, become a spacer. Most preferably, the G-protein that couples to the non-endogenous GPCR will have been identified prior to the creation of the GPCR Fusion Protein construct. Because there are only a few G proteins that have been
20 identified, it is preferred that a construct comprising the sequence of the G protein (*i.e.*, a universal G protein construct) be available for insertion of an endogenous GPCR sequence therein; this provides for efficiency in the context of large-scale screening of a variety of different endogenous GPCRs having different sequences.

As noted above, constitutively activated GPCRs that couple to Gi, Gz and Go are expected to inhibit the formation of cAMP making assays based upon these types of GPCRs challenging (*i.e.*, the cAMP signal decreases upon activation thus making the direct identification of, *e.g.*, inverse agonists (which would further decrease this signal), interesting).

5 As will be disclosed herein, we have ascertained that for these types of receptors, it is possible to create a GPCR Fusion Protein that is not based upon the endogenous GPCR's endogenous G protein, in an effort to establish a viable cyclase-based assay. Thus, for example, a Gz coupled receptor such as H9, a GPCR Fusion Protein can be established that utilizes a Gs fusion protein - we believe that such a fusion construct, upon expression, "drives" or "forces"
10 the non-endogenous GPCR to couple with, *e.g.*, Gs rather than the "natural" Gz protein, such that a cyclase-based assay can be established. Thus, for Gi, Gz and Go coupled receptors, we prefer that that when a GPCR Fusion Protein is used and the assay is based upon detection of adenylyl cyclase activity, that the fusion construct be established with Gs (or an equivalent G protein that stimulates the formation of the enzyme adenylyl cyclase).

15 F. Medicinal Chemistry

Generally, but not always, direct identification of candidate compounds is preferably conducted in conjunction with compounds generated via combinatorial chemistry techniques, whereby thousands of compounds are randomly prepared for such analysis. Generally, the results of such screening will be compounds having unique core structures; thereafter, these
20 compounds are preferably subjected to additional chemical modification around a preferred core structure(s) to further enhance the medicinal properties thereof. Such techniques are known to those in the art and will not be addressed in detail in this patent document.

G. Pharmaceutical compositions

Candidate compounds selected for further development can be formulated into pharmaceutical compositions using techniques well known to those in the art. Suitable pharmaceutically-acceptable carriers are available to those in the art; for example, see
5 Remington's Pharmaceutical Sciences, 16th Edition, 1980, Mack Publishing Co., (Oslo et al., eds.)

H. Other Utility

Although a preferred use of the non-endogenous versions the human GPCRs disclosed herein may be for the direct identification of candidate compounds as inverse agonists,
10 agonists or partial agonists (preferably for use as pharmaceutical agents), these versions of human GPCRs can also be utilized in research settings. For example, *in vitro* and *in vivo* systems incorporating GPCRs can be utilized to further elucidate and understand the roles these receptors play in the human condition, both normal and diseased, as well as understanding the role of constitutive activation as it applies to understanding the signaling
15 cascade. The value in non-endogenous human GPCRs is that their utility as a research tool is enhanced in that, because of their unique features, non-endogenous human GPCRs can be used to understand the role of these receptors in the human body before the endogenous ligand therefor is identified. Other uses of the disclosed receptors will become apparent to those in the art based upon, *inter alia*, a review of this patent document.

20

EXAMPLES

The following examples are presented for purposes of elucidation, and not limitation, of the present invention. While specific nucleic acid and amino acid sequences are disclosed herein, those of ordinary skill in the art are credited with the ability to make minor

modifications to these sequences while achieving the same or substantially similar results reported below. The traditional approach to application or understanding of sequence cassettes from one sequence to another (e.g. from rat receptor to human receptor or from human receptor A to human receptor B) is generally predicated upon sequence alignment techniques whereby the sequences are aligned in an effort to determine areas of commonality. The mutational approach disclosed herein does not rely upon this approach but is instead based upon an algorithmic approach and a positional distance from a conserved proline residue located within the TM6 region of human GPCRs. Once this approach is secured, those in the art are credited with the ability to make minor modifications thereto to achieve substantially the same results (*i.e.*, constitutive activation) disclosed herein. Such modified approaches are considered within the purview of this disclosure

Example 1

ENDOGENOUS HUMAN GPCRS

1. Identification of Human GPCRs

Certain of the disclosed endogenous human GPCRs were identified based upon a review of the GenBank™ database information. While searching the database, the following cDNA clones were identified as evidenced below (Table C).

TABLE C

Disclosed Human Orphan GPCRs	Accession Number	Complete DNA Sequence (Base Pairs)	Open Reading Frame (Base Pairs)	Nucleic Acid SEQ.ID. NO.	Amino Acid SEQ.ID. NO.
hARE-3	AL033379	111,389 bp	1,260 bp	1	2
hARE-4	AC006087	226,925 bp	1,119 bp	3	4
hARE-5	AC006255	127,605 bp	1,104 bp	5	6
hRUP3	AL035423	140,094 bp	1,005 bp	7	8

- 24 -

hRUP5	AC005849	169,144 bp	1,413 bp	9	10
hRUP6	AC005871	218,807 bp	1,245 bp	11	12
hRUP7	AC007922	158,858 bp	1,173 bp	13	14

Other disclosed endogenous human GPCRs were identified by conducting a BLAST™
 5 search of EST database (dbest) using the following EST clones as query sequences. The
 following EST clones identified were then used as a probe to screen a human genomic library
 (Table D).

TABLE D

10	Disclosed Human Orphan GPCRs	Query (Sequence)	EST Clone/ Accession No. Identified	Open Reading Frame (Base Pairs)	Nucleic Acid SEQ.ID.NO.	Amino Acid SEQ.ID.NO.
	hGPCR27	Mouse GPCR27	AA775870	1,125 bp	17	18
	hARE-1	TDAG	1689643 AI090920	999 bp	19	20
15	hARE-2	GPCR27	68530 AA359504	1,122 bp	21	22
	hPPR1	Bovine PPR1	238667 H67224	1,053 bp	23	24
	hG2A	Mouse 1179426	<i>See Example 2(a), below</i>	1,113 bp	25	26
	hCHN3	N.A.	EST 36581 (full length)	1,113 bp	27	28
	hCHN4	TDAG	1184934 AA804531	1,077 bp	29	30
20	hCHN6	N.A.	EST 2134670 (full length)	1,503 bp	31	32
	hCHN8	KIAA0001	EST 764455	1,029 bp	33	34
	hCHN 9	1365839	EST 1541536	1,077 bp	35	36
	hCHN10	Mouse EST 1365839	Human 1365839	1,005 bp	37	38
	hRUP4	N.A.	AI307658	1,296 bp	39	40
25		N.A. = "not applicable".				

2. Full Length Cloning

a. Human G2A

Mouse EST clone 1179426 was used to obtain a human genomic clone containing all

- 25 -

but three amino acid G2A coding sequences. The 5' of this coding sequence was obtained by using 5'RACE, and the template for PCR was Clontech's Human Spleen Marathon-Ready™ cDNA. The disclosed human G2A was amplified by PCR using the G2A cDNA specific primers for the first and second round PCR as shown in SEQ.ID.NO.: 41 and SEQ.ID.NO.:42

5 as follows:

5'-CTGTGTACAGCAGTTCGCAGAGTG-3' (SEQ.ID.NO.: 41; 1st round PCR)

5'-GAGTGCCAGGCAGAGCAGGTAGAC-3' (SEQ.ID.NO.: 42; second round PCR).

PCR was performed using Advantage GC Polymerase Kit (Clontech; manufacturing instructions will be followed), at 94°C for 30 sec followed by 5 cycles of 94°C for 5 sec and
10 72°C for 4 min; and 30 cycles of 94° for 5 sec and 70° for 4 min. An approximate 1.3 Kb PCR fragment was purified from agarose gel, digested with Hind III and Xba I and cloned into the expression vector pRC/CMV2 (Invitrogen). The cloned-insert was sequenced using the T7 Sequenase™ kit (USB Amersham; manufacturer instructions followed) and the sequence was compared with the presented sequence. Expression of the human G2A was detected by
15 probing an RNA dot blot (Clontech; manufacturer instructions followed) with the P³²-labeled fragment.

b. CHN9

Sequencing of the EST clone 1541536 showed CHN9 to be a partial cDNA clone having only an initiation codon; *i.e.*, the termination codon was missing. When CHN9
20 was used to blast against data base (nr), the 3' sequence of CHN9 was 100% homologous to the 5' untranslated region of the leukotriene B4 receptor cDNA, which contained a termination codon in the frame with CHN9 coding sequence. To determine whether the 5' untranslated region of LTB4R cDNA was the 3' sequence of CHN9, PCR was performed using primers based upon the 5' sequence flanking the initiation codon found in CHN9 and

- 26 -

the 3' sequence around the termination codon found in the LTB4R 5' untranslated region.

The 5' primer sequence utilized was as follows:

5'-CCCGAATTCCTGCTTGCTCCCAGCTTGGCCC-3' (SEQ.ID.NO.: 43; sense) and

5'-TGTGGATCCTGCTGTCAAAGGTCCCATTCGG-3' (SEQ.ID.NO.: 44; antisense).

- 5 PCR was performed using thymus cDNA as a template and rTth polymerase (Perkin Elmer) with the buffer system provided by the manufacturer, 0.25 uM of each primer, and 0.2 mM of each 4 nucleotides. The cycle condition was 30 cycles of 94°C for 1 min, 65°C for 1min and 72 °C for 1 min and 10 sec. A 1.1kb fragment consistent with the predicted size was obtained from PCR. This PCR fragment was subcloned into pCMV (*see below*) and
- 10 sequenced (*see*, SEQ.ID.NO.: 35).

c. RUP 4

The full length RUP4 was cloned by RT-PCR with human brain cDNA (Clontech) as templates:

5'-TCACAATGCTAGGTGTGGTC-3' (SEQ.ID.NO.: 45; sense) and

- 15 5'-TGCATAGACAATGGGATTACAG-3' (SEQ.ID.NO.: 46; antisense).

PCR was performed using TaqPlus Precision™ polymerase (Stratagene; manufacturing instructions followed) by the following cycles: 94°C for 2 min; 94°C 30 sec; 55°C for 30 sec, 72°C for 45 sec, and 72°C for 10 min. Cycles 2 through 4 were repeated 30 times.

- The PCR products were separated on a 1% agarose gel and a 500 bp PCR fragment
- 20 was isolated and cloned into the pCRII-TOPO™ vector (Invitrogen) and sequenced using the T7 DNA Sequenase™ kit (Amsham) and the SP6/T7 primers (Stratagene). Sequence analysis revealed that the PCR fragment was indeed an alternatively spliced form of AI307658 having a continuous open reading frame with similarity to other GPCRs. The completed sequence of this PCR fragment was as follows:

5'-TCACAATGCTAGGTGTGGTCTGGCTGGTGGCAGTCATCGTAGGATCACCCATGTGGCAC
GTGCAACAACCTTGAGATCAAATATGACTTCCTATATGAAAAGGAACACATCTGCTGCTTAAGA
GTGGACCAGCCCTGTGCACCAGAAGATCTACACCACCTTCATCCTTGTCATCCTCTTCCTCCTGC
CTCTTATGGTGATGCTTATTCTGTACGTAAAATTGGTTATGAACCTTTGGATAAAGAAAAGAGTT
5 GGGGATGGTTTCAGTGCCTCGAACTATTCATGGAAAAGAAATGTCCAAAATAGCCAGGAAGAAG
AAACGAGCTGTCAATTATGATGGTGACAGTGGTGGCTCTCTTTGCTGTGTGCTGGGCACCATTCC
ATGTTGTCCATATGATGATTGAATACAGTAATTTTGAAGAAGGAATATGATGATGTCACAATCAA
GATGATTTTTGCTATCGTGCAAATTATTGGATTTTCCAACCTCCATCTGTAATCCCATTTGTCTATGCA-
3' (SEQ.ID.NO.: 47)

- 10 Based on the above sequence, two sense oligonucleotide primer sets:

5'-CTGCTTAGAAGAGTGGACCAG-3' (SEQ.ID.NO.: 48; oligo 1),

5'-CTGTGCACCAGAAGATCTACAC-3' (SEQ.ID.NO.: 49; oligo 2) and

two antisense oligonucleotide primer sets:

5'-CAAGGATGAAGGTGGTGTAGA-3' (SEQ.ID.NO.: 50; oligo 3)

- 15 5'-GTGTAGATCTTCTGGTGCACAGG-3' (SEQ.ID.NO.: 51; oligo 4)

were used for 3'- and 5'-RACE PCR with a human brain Marathon-Ready™ cDNA (Clontech, Cat# 7400-1) as template, according to manufacture's instructions. DNA fragments generated by the RACE PCR were cloned into the pCRII-TOPO™ vector (Invitrogen) and sequenced using the SP6/T7 primers (Stratagene) and some internal primers.

- 20 The 3' RACE product contained a poly(A) tail and a completed open reading frame ending at a TAA stop codon. The 5' RACE product contained an incomplete 5' end; i.e., the ATG initiation codon was not present.

Based on the new 5' sequence, oligo 3 and the following primer:

5'-GCAATGCAGGTCATAGTGAGC-3' (SEQ.ID.NO.: 52; oligo 5)

- 25 were used for the second round of 5' race PCR and the PCR products were analyzed as above.

A third round of 5' race PCR was carried out utilizing antisense primers:

5'-TGGAGCATGGTGACGGGAATGCAGAAG-3' (SEQ.ID.NO.: 53; oligo 6) and

5'-GTGATGAGCAGGTCACTGAGCGCCAAG-3' (SEQ.ID.NO.: 54; oligo 7).

The sequence of the 5' RACE PCR products revealed the presence of the initiation codon

- 28 -

ATG, and further round of 5' race PCR did not generate any more 5' sequence. The completed 5' sequence was confirmed by RT-PCR using sense primer

5'-GCAATGCAGGCGCTTAACATTAC-3' (SEQ.ID.NO.: 55; oligo 8)

and oligo 4 as primers and sequence analysis of the 650 bp PCR product generated from
5 human brain and heart cDNA templates (Clontech, Cat# 7404-1). The completed 3' sequence was confirmed by RT-PCR using oligo 2 and the following antisense primer:

5'-TTGGGTTACAATCTGAAGGGCA-3' (SEQ.ID.NO.:56; oligo 9)

and sequence analysis of the 670 bp PCR product generated from human brain and heart cDNA templates. (Clontech, Cat# 7404-1).

10

d. RUP5

The full length RUP5 was cloned by RT-PCR using a sense primer upstream from ATG, the initiation codon (SEQ.ID.NO.:57), and an antisense primer containing TCA as the stop codon (SEQ.ID.NO.:58), which had the following sequences:

5'-ACTCCGTGTCCAGCAGGACTCTG-3' (SEQ.ID.NO.: 57)

15 5'-TGCGTGTTTCCTGGACCCTCACGTG-3' (SEQ.ID.NO.: 58)

and human peripheral leukocyte cDNA (Clontech) as a template. Advantage™ cDNA polymerase (Clontech) was used for the amplification in a 50ul reaction by the following cycle with step 2 through step 4 repeated 30 times: 94°C for 30 sec; 94° for 15 sec; 69° for 40 sec; 72°C for 3 min; and 72°C fro 6 min. A 1.4kb PCR fragment was isolated and cloned with
20 the pCRII-TOPO™ vector (Invitrogen) and completely sequenced using the T7 DNA Sequenase™ kit (Amsham). See, SEQ.ID.NO.: 9.

e. RUP6

The full length RUP6 was cloned by RT-PCR using primers:

5'-CAGGCCTTGGATTTTAATGTCAGGGATGG-3' (SEQ.ID.NO.: 59) and

- 29 -

5'-GGAGAGTCAGCTCTGAAAGAATTCAGG-3' (SEQ.ID.NO.: 60);

and human thymus Marathon-Ready™ cDNA (Clontech) as a template. Advantage cDNA polymerase (Clontech, according to manufacturer's instructions) was used for the amplification in a 50ul reaction by the following cycle: 94°C for 30sec; 94°C for 5 sec; 66°C for 40sec; 72°C for 2.5 sec and 72°C for 7 min. Cycles 2 through 4 were repeated 30 times. A 1.3 Kb PCR fragment was isolated and cloned into the pCRII-TOPO™ vector (Invitrogen) and completely sequenced (*see*, SEQ.ID.NO.: 11) using the ABI Big Dye Terminator™ kit (P.E. Biosystem).

f. RUP7

10 The full length RUP7 was cloned by RT-PCR using primers:

5'-TGATGTGATGCCAGATACTAATAGCAC-3' (SEQ.ID.NO.: 61; sense) and

5'-CCTGATTCATTAGGTGAGATTGAGAC-3' (SEQ.ID.NO.: 62; antisense)

and human peripheral leukocyte cDNA (Clontech) as a template. Advantage™ cDNA polymerase (Clontech) was used for the amplification in a 50 ul reaction by the following cycle with step 2 to step 4 repeated 30 times: 94°C for 2 minutes; 94°C for 15 seconds; 60°C for 20 seconds; 72°C for 2 minutes; 72°C for 10 minutes. A 1.25 Kb PCR fragment was isolated and cloned into the pCRII-TOPO™ vector (Invitrogen) and completely sequenced using the ABI Big Dye Terminator™ kit (P.E. Biosystem). *See*, SEQ.ID.NO.: 13.

3. Angiotensin II Type 1 Receptor ("AT1")

20 The endogenous human angiotensin II type 1 receptor ("AT1") was obtained by PCR using genomic DNA as template and rTth polymerase (Perkin Elmer) with the buffer system provided by the manufacturer, 0.25 µM of each primer, and 0.2 mM of each 4 nucleotides. The cycle condition was 30 cycles of 94°C for 1 min, 55°C for 1min and 72°C for 1.5 min. The 5' PCR primer contains a HindIII site with the sequence:

- 30 -

5'-CCCAAGCTTCCCCAGGTGTATTTGAT-3' (SEQ.ID.NO.: 63)

and the 3' primer contains a BamHI site with the following sequence:

5'-GTTGGATCCACATAATGCATTTTCTC-3' (SEQ.ID.NO.: 64).

The resulting 1.3 kb PCR fragment was digested with HindIII and BamHI and cloned into
5 HindIII-BamHI site of pCMV expression vector. The cDNA clone was fully sequenced.
Nucleic acid (SEQ.ID.NO.: 65) and amino acid (SEQ.ID.NO.: 66) sequences for human AT1
were thereafter determined and verified.

4. GPR38

To obtain GPR38, PCR was performed by combining two PCR fragments, using
10 human genomic cDNA as template and rTth polymerase (Perkin Elmer) with the buffer system
provided by the manufacturer, 0.25uM of each primer, and 0.2 mM of each 4 nucleotides.
The cycle condition for each PCR reaction was 30 cycles of 94°C for 1 min, 62°C for 1min
and 72°C for 2 min.

The first fragment was amplified with the 5' PCR primer that contained an end site
15 with the following sequence:

5'-ACCATGGGCAGCCCCTGGAACGGCAGC-3' (SEQ.ID.NO.:67)

and a 3' primer having the following sequence:

5'-AGAACCACCACCAGCAGGACGCGGACGGTCTGCCGGTGG-3' (SEQ.ID.NO.:68).

The second PCR fragment was amplified with a 5' primer having the following sequence:

20 5'-GTCCGCGTCCTGCTGGTGGTGGTTCTGGCATTATAATT-3' (SEQ.ID.NO.: 69)

and a 3' primer that contained a BamHI site and having the following sequence:

5'-CCTGGATCCTTATCCCATCGTCTTCACGTTAGC-3' (SEQ.ID.NO.: 70).

The two fragments were used as templates to amplify GPR38, using SEQ.ID.NO.: 67 and
SEQ.ID.NO.: 70 as primers (using the above-noted cycle conditions). The resulting 1.44kb

- 31 -

PCR fragment was digested with BamHI and cloned into Blunt-BamHI site of pCMV expression vector.

5. MC4

To obtain MC4, PCR was performed using human genomic cDNA as template and rTth polymerase (Perkin Elmer) with the buffer system provided by the manufacturer, 0.25uM of each primer, and 0.2 mM of each 4 nucleotides. The cycle condition for each PCR reaction was 30 cycles of 94°C for 1 min, 54°C for 1min and 72°C for 1.5 min.

The 5' PCR contained an EcoRI site with the sequence:

5'-CTGGAATTCTCTGCCAGCATGGTGA-3' (SEQ.ID.NO.: 71)

and the 3' primer contained a BamHI site with the sequence:

5'-GCAGGATCCTATATTGCGTGCTCTGTCCCC'-3 (SEQ.ID.NO.: 72).

The 1.0 kb PCR fragment was digest with EcoRI and BamHI and cloned into EcoRI-BamHI site of pCMV expression vector. Nucleic acid (SEQ.ID.NO.: 73) and amino acid (SEQ.ID.NO.: 74) sequences for human MC4 were thereafter determined.

6. CCKB

To obtain CCKB, PCR was performed using human stomach cDNA as template and rTth polymerase (Perkin Elmer) with the buffer system provided by the manufacturer, 0.25uM of each primer, and 0.2 mM of each 4 nucleotides. The cycle condition for each PCR reaction was 30 cycles of 94°C for 1 min, 65°C for 1min and 72°C for 1 min and 30 sec.

The 5' PCR contained a HindIII site with the sequence:

5'-CCGAAGCTTCGAGCTGAGTAAGGCGGCGGGCT-3' (SEQ.ID.NO.: 75)

and the 3' primer contained an EcoRI site with the sequence:

5'-GTGGAATTCATTTGCCCTGCCTCAACCCCA-3 (SEQ.ID.NO.: 76).

The resulting 1.44 kb PCR fragment was digest with HindIII and EcoRI and cloned into

HindIII-EcoRI site of pCMV expression vector. Nucleic acid (SEQ.ID.NO.: 77) and amino acid (SEQ.ID.NO.: 78) sequences for human CCKB were thereafter determined.

7. TDAG8

To obtain TDAG8, PCR was performed using genomic DNA as template and rTth polymerase (Perkin Elmer) with the buffer system provided by the manufacturer, 0.25 μ M of each primer, and 0.2 mM of each 4 nucleotides. The cycle condition was 30 cycles of 94°C for 1 min, 56°C for 1min and 72 °C for 1 min and 20 sec. The 5' PCR primer contained a HindIII site with the following sequence:

5'-TGCAAGCTTAAAAAGGAAAAATGAACAGC-3' (SEQ.ID.NO.: 79)

and the 3' primer contained a BamHI site with the following sequence:

5'-TAAGGATCCCTTCCCTTCAAAACATCCTTG-3' (SEQ.ID.NO.: 80).

The resulting 1.1 kb PCR fragment was digested with HindIII and BamHI and cloned into HindIII-BamHI site of pCMV expression vector. Three resulting clones sequenced contained three potential polymorphisms involving changes of amino acid 43 from Pro to Ala, amino acid 97 from Lys to Asn and amino acid 130 from Ile to Phe. Nucleic acid (SEQ.ID.NO.: 81) and amino acid (SEQ.ID.NO.: 82) sequences for human TDAG8 were thereafter determined.

8. H9

To obtain H9, PCR was performed using pituitary cDNA as template and rTth polymerase (Perkin Elmer) with the buffer system provided by the manufacturer, 0.25 μ M of each primer, and 0.2 mM of each 4 nucleotides. The cycle condition was 30 cycles of 94°C for 1 min, 62°C for 1 min and 72°C for 2 min. The 5' PCR primer contained a HindIII site with the following sequence:

5'-GGAAAGCTTAACGATCCCCAGGAGCAACAT-3' (SEQ.ID.NO.:15)

and the 3' primer contained a BamHI site with the following sequence:

5'-CTGGGATCCTACGAGAGCATTTTTCACACAG-3' (SEQ.ID.NO.:16).

The resulting 1.9 kb PCR fragment was digested with HindIII and BamHI and cloned into HindIII-BamHI site of pCMV expression vector. H9 contained three potential polymorphisms involving changes of amino acid P320S, S493N and amino acid G448A. Nucleic acid
5 (SEQ.ID.NO.: 139) and amino acid (SEQ.ID.NO.: 140) sequences for human H9 were thereafter determined and verified.

Example 2

PREPARATION OF NON-ENDOGENOUS, CONSTITUTIVELY ACTIVATED GPCRS

Those skilled in the art are credited with the ability to select techniques for
10 mutation of a nucleic acid sequence. Presented below are approaches utilized to create non-endogenous versions of several of the human GPCRs disclosed above. The mutations disclosed below are based upon an algorithmic approach whereby the 16th amino acid (located in the IC3 region of the GPCR) from a conserved proline residue (located in the TM6 region of the GPCR, near the TM6/IC3 interface) is mutated, most preferably to a
15 lysine amino acid residue.

1. Transformer Site-Directed™ Mutagenesis

Preparation of non-endogenous human GPCRs may be accomplished on human GPCRs using Transformer Site-Directed™ Mutagenesis Kit (Clontech) according to the manufacturer instructions. Two mutagenesis primers are utilized, most preferably a lysine
20 mutagenesis oligonucleotide that creates the lysine mutation, and a selection marker oligonucleotide. For convenience, the codon mutation to be incorporated into the human GPCR is also noted, in standard form (Table E):

- 34 -

TABLE E

	Receptor Identifier	Codon Mutation
	hARE-3	F313K
	hARE-4	V233K
5	hARE-5	A240K
	hGPCR14	L257K
	hGPCR27	C283K
	hARE-1	E232K
	hARE-2	G285K
10	hPPR1	L239K
	hG2A	K232A
	hRUP3	L224K
	hRUP5	A236K
	hRUP6	N267K
15	hRUP7	A302K
	hCHN4	V236K
	hMC4	A244K
	hCHN3	S284K
	hCHN6	L352K
20	hCHN8	N235K
	hCHN9	G223K
	hCHN10	L231K
	hH9	F236K

The following GPCRs were mutated according with the above method using the

25 designated sequence primers (Table F).

- 35 -

TABLE F

Receptor Identifier	Codon Mutation	Lysine Mutagenesis (SEQ.ID.NO.) 5'-3' orientation, mutation sequence underlined	Selection Marker (SEQ.ID.NO.) 5'-3' orientation
hRUP4	V272K	CAGGAAGAAGAA <u>AC</u> GAGC TGTCATTATGATGGTGACA GTG (83)	CACTGTCACCATCATAATG ACAGCTCGTTTCTTCTTCC TG (84)
hAT1	<i>see below</i>	alternative approach; <i>see below</i>	alternative approach; <i>see below</i>
hGPR38	V297K	GGCCACCGGCAGACCA <u>AA</u> C GCGTCCTGCTG (85)	CTCCTTCGGTCCTCCTATC GTTGTCAGAAAGT (86)
hCCKB	V332K	alternative approach; <i>see below</i>	alternative approach; <i>see below</i>
hTDAG8	I225K	GGAAAAGAAGAGAATCAA <u>AAAA</u> CTACTTGTCAGCATC (87)	CTCCTTCGGTCCTCCTATC GTTGTCAGAAAGT (88)
hH9	F236K	GCTGAGGTTTCGAATA <u>AA</u> C TAACCATGTTTGTG (143)	CTCCTTCGGTCCTCCTATC GTTGTCAGAAAGT (144)
hMC4	A244K	GCCAATATGAAGGGAA <u>AA</u> ATTACCTTGACCATC (137)	CTCCTTCGGTCCTCCTATC GTTGTCAGAAAGT (138)

The non-endogenous human GPCRs were then sequenced and the derived and verified nucleic acid and amino acid sequences are listed in the accompanying "Sequence Listing" appendix to this patent document, as summarized in Table G below:

TABLE G

	Non Endogenous Human GPCR	Nucleic Acid Sequence Listing	Amino Acid Sequence Listing
15	hRUP4 (V272K)	SEQ.ID.NO.: 127	SEQ.ID.NO.: 128
20	hAT1 (<i>see alternative approaches below</i>)	(<i>see alternative approaches below</i>)	(<i>see alternative approaches, below</i>)
	hGPR38 (V297K)	SEQ.ID.NO.: 129	SEQ.ID.NO.: 130
25	hCCKB (V332K)	SEQ.ID.NO.: 131	SEQ.ID.NO.: 132
	hTDAG8 (I225K)	SEQ.ID.NO.: 133	SEQ.ID.NO.: 134
	hH9 (F236K)	SEQ.ID.NO.: 141	SEQ.ID.NO.: 142
30	hMC4 (A244K)	SEQ.ID.NO.: 135	SEQ.ID.NO.: 136

2. Alternative Approaches For Creation of Non-Endogenous Human GPCRs

a. AT1

1. F239K Mutation

Preparation of a non-endogenous, constitutively activated human AT1 receptor was accomplished by creating an F239K mutation (see, SEQ.ID.NO.: 89 for nucleic acid sequence, and SEQ.ID.NO.: 90 for amino acid sequence). Mutagenesis was performed using Transformer Site-Directed Mutagenesis™ Kit (Clontech) according to the to manufacturer's instructions. The two mutagenesis primers were used, a lysine mutagenesis oligonucleotide (SEQ.ID.NO.: 91) and a selection marker oligonucleotide (SEQ.ID.NO.: 92), which had the following sequences:

5'-CCAAGAAATGATGATATTAAAAAGATAATTATGGC-3' (SEQ.ID.NO.: 91)

5'-CTCCTTCGGTCCTCCTATCGTTGTCAGAAAGT-3' (SEQ.ID.NO.: 92),

respectively.

2. N111A Mutation

Preparation of a non-endogenous human AT1 receptor was also accomplished by creating an N111A mutation (see, SEQ.ID.NO.:93 for nucleic acid sequence, and SEQ.ID.NO.: 94 for amino acid sequence). Two PCR reactions were performed using pfu polymerase (Stratagene) with the buffer system provided by the manufacturer, supplemented with 10% DMSO, 0.25 μ M of each primer, and 0.5 mM of each 4 nucleotides. The 5' PCR sense primer used had the following sequence:

5'-CCCAAGCTTCCCCAGGTGTATTTGAT-3' (SEQ.ID.NO.: 95)

and the antisense primer had the following sequence:

- 37 -

5'-CCTGCAGGCGAAACTGACTCTGGCTGAAG-3' (SEQ.ID.NO.: 96).

The resulting 400 bp PCR fragment was digested with HindIII site and subcloned into HindIII-SmaI site of pCMV vector (5' construct). The 3' PCR sense primer used had the following sequence:

5 5'-CTGTACGCTAGTGTGTTTCTACTCACGTGTCTCAGCATTGAT-3' (SEQ.ID.NO.: 97)

and the antisense primer had the following sequence:

5'-GTTGGATCCACATAATGCATTTTCTC-3' (SEQ.ID.NO.: 98)

The resulting 880 bp PCR fragment was digested with BamHI and inserted into Pst (blunted by T4 polymerase) and BamHI site of 5' construct to generate the full length

10 N111A construct. The cycle condition was 25 cycles of 94°C for 1 min, 60°C for 1 min and 72°C for 1 min (5' PCR) or 1.5 min (3' PCR).

3. AT2K255IC3 Mutation

Preparation of a non-endogenous, constitutively activated human AT1 was accomplished by creating an AT2K255IC3 "domain swap" mutation (see, SEQ.ID.NO.:99 for nucleic acid sequence, and SEQ.ID.NO.: 100 for amino acid sequence). Restriction sites flanking IC3 of AT1 were generated to facilitate replacement of the IC3 with corresponding IC3 from angiotensin II type 2 receptor (AT2). This was accomplished by performing two PCR reactions. A 5' PCR fragment (Fragment A) encoded from the 5' untranslated region to the beginning of IC3 was generated by utilizing SEQ.ID.NO.: 63 as

20 sense primer and the following sequence:

5'-TCCGAATTCCAAAATAACTTGTAAGAATGATCAGAAA-3' (SEQ.ID.NO.: 101)

as antisense primer. A 3' PCR fragment (Fragment B) encoding from the end of IC3 to the 3' untranslated region was generated by using the following sequence:

5'-AGATCTTAAGAAGATAATTATGGCAATTGTGCT-3' (SEQ.ID.NO.: 102)

- 38 -

as sense primer and SEQ.ID.NO.: 64 as antisense primer. The PCR condition was 30 cycles of 94°C for 1 min, 55°C for 1min and 72°C for 1.5 min using endogenous AT1 cDNA clone as template and pfu polymerase (Stratagene), with the buffer systems provided by the manufacturer, supplemented with 10% DMSO, 0.25 µM of each primer, and 0.5 mM of each 4 nucleotides. Fragment A (720 bp) was digested with HindIII and EcoRI and subcloned. Fragment B was digested with BamHI and subcloned into pCMV vector with an EcoRI site 5' to the cloned PCR fragment.

The DNA fragment (Fragment C) encoding IC3 of AT2 with a L255K mutation and containing an EcoRI cohesive end at 5' and a AflII cohesive end at 3', was generated by annealing 2 synthetic oligonucleotides having the following sequences:

5'AATTCGAAAACACTTACTGAAGACGAATAGCTATGGGAAGAACAGGATAACCCGTGACCAA
G-3' (sense; SEQ.ID.NO.: 103)

5'TTAAGTTGGTCACGGGTTATCCTGTTCTTCCCATAGCTATTCGTCTTCAGT
AAGTGTTTTTCG-3' (antisense; SEQ.ID.NO.: 104).

Fragment C was inserted in front of Fragment B through EcoRI and AflII site. The resulting clone was then ligated with the Fragment A through the EcoRI site to generate AT1 with AT2K255IC3.

4. A243+ Mutation

Preparation of a non-endogenous human AT1 receptor was also accomplished by creating an A243+ mutation (see, SEQ.ID.NO.: 105 for nucleic acid sequence, and SEQ.ID.NO.: 106 for amino acid sequence). An A243+ mutation was constructed using the following PCR based strategy: Two PCR reactions was performed using pfu polymerase (Stratagene) with the buffer system provided by the manufacturer supplemented with 10% DMSO, 0.25 µM of each primer, and 0.5 mM of each 4 nucleotides. The 5' PCR sense primer

- 39 -

utilized had the following sequence:

5'-CCCAAGCTTCCCCAGGTGTATTTGAT-3' (SEQ.ID.NO.: 107)

and the antisense primer had the following sequence:

5'-AAGCACAATTGCTGCATAATTATCTTAAAAATATCATC-3' (SEQ.ID.NO.: 108).

- 5 The 3' PCR sense primer utilized had the following sequence:

5'-AAGATAATTATGGCAGCAATTGTGCTTTTCTTTTCTTT-3' (SEQ.ID.NO.: 109)

containing the Ala insertion and antisense primer:

5'-GTTGGATCCACATAATGCATTTTCTC-3' (SEQ.ID.NO.: 110).

The cycle condition was 25 cycles of 94°C for 1 min, 54°C for 1min and 72°C for 1.5 min.

- 10 An aliquot of the 5' and 3' PCR were then used as co-template to perform secondary PCR using the 5' PCR sense primer and 3' PCR antisense primer. The PCR condition was the same as primary PCR except the extension time was 2.5 min. The resulting PCR fragment was digested with HindIII and BamHI and subcloned into pCMV vector. (See, SEQ.ID.NO.: 105)

15 **4. CCKB**

Preparation of the non-endogenous, constitutively activated human CCKB receptor was accomplished by creating a V322K mutation (see, SEQ.ID.NO.: 111 for nucleic acid sequence and SEQ.ID.NO.: 112 for amino acid sequence). Mutagenesis was performed by PCR via amplification using the wildtype CCKB from Example 1.

- 20 The first PCR fragment (1kb) was amplified by using SEQ.ID.NO.: 75 and an antisense primer comprising a V322K mutation:

5'-CAGCAGCATGCGCTTCACGCGCTTCTTAGCCCAG-3' (SEQ.ID.NO.: 113).

The second PCR fragment (0.44kb) was amplified by using a sense primer comprising the V322K mutation:

5'-AGAAGCGCGTGAAGCGCATGCTGCTGGTGATCGTT-3' (SEQ.ID.NO.: 114) and SEQ.ID.NO.:

76.

The two resulting PCR fragments were then used as template for amplifying CCKB comprising V332K, using SEQ.ID.NO.: 75 and SEQ.ID.NO.: 76 and the above-noted system and conditions. The resulting 1.44kb PCR fragment containing the V332K mutation was digested with HindIII and EcoRI and cloned into HindIII-EcoRI site of pCMV expression vector. (See, SEQ.ID.NO.: 111).

3. QuikChange™ Site-Directed™ Mutagenesis

Preparation of non-endogenous human GPCRs can also be accomplished by using QuikChange™ Site-Directed™ Mutagenesis Kit (Stratagene, according to manufacturer's instructions). Endogenous GPCR is preferably used as a template and two mutagenesis primers utilized, as well as, most preferably, a lysine mutagenesis oligonucleotide and a selection marker oligonucleotide (included in kit). For convenience, the codon mutation incorporated into the human GPCR and the respective oligonucleotides are noted, in standard form (Table H):

- 41 -

TABLE H

Receptor Identifier	Codon Mutation	Lysine Mutagenesis (SEQ.ID.NO.) 5'-3' orientation, mutation underlined	Selection Marker (SEQ.ID.NO.) 5'-3' orientation
hCHN3	S284K	ATGGAGAAAAGAATCA <u>AAA</u> AGAA TGTTCTATATA (115)	TATATAGAACATTCTTTT GATTCTTTCTCCAT (116)
hCHN6	L352K	CGCTCTCTGGCCTTGA <u>AAG</u> CGCAC GCTCAGC (117)	GCTGAGCGTGCGCTTCA AGGCCAGAGAGCG (118)
5 hCHN8	N235K	CCCAGGAAAAAGGTGA <u>AA</u> AGTCA AAGTTTTC (119)	GAAAACTTTGACTTTCAC CTTTTTCCTGGG (120)
hCHN9	G223K	GGGGCGCGGGTGAA <u>AC</u> CGGCTGG TGAGC (121)	GCTCACCGAGCCGTTTCA CCCGCGCCCC (122)
hCHN10	L231K	CCCCTTGAA <u>AA</u> GCCTAAGAACTT GGTCATC (123)	GATGACCAAGTTCTTAG GCTTTTCAAGGGG (124)

Example 3**RECEPTOR EXPRESSION**

10 Although a variety of cells are available to the art for the expression of proteins, it is most preferred that mammalian cells be utilized. The primary reason for this is predicated upon practicalities, *i.e.*, utilization of, *e.g.*, yeast cells for the expression of a GPCR, while possible, introduces into the protocol a non-mammalian cell which may not (indeed, in the case of yeast, does not) include the receptor-coupling, genetic-mechanism and secretary

15 pathways that have evolved for mammalian systems - thus, results obtained in non-mammalian cells, while of potential use, are not as preferred as that obtained from mammalian cells. Of the mammalian cells, COS-7, 293 and 293T cells are particularly preferred, although the specific mammalian cell utilized can be predicated upon the particular needs of the artisan.

On day one, 1×10^7 293T cells per 150mm plate were plated out. On day two, two

20 reaction tubes were prepared (the proportions to follow for each tube are per plate): tube A was prepared by mixing 20 μ g DNA (*e.g.*, pCMV vector; pCMV vector with receptor cDNA, etc.) in 1.2ml serum free DMEM (Irvine Scientific, Irvine, CA); tube B was

- 42 -

prepared by mixing 120 μ l lipofectamine (Gibco BRL) in 1.2ml serum free DMEM. Tubes A and B were admixed by inversions (several times), followed by incubation at room temperature for 30-45min. The admixture is referred to as the "transfection mixture". Plated 293T cells were washed with 1XPBS, followed by addition of 10ml serum free DMEM. 2.4ml of the transfection mixture were added to the cells, followed by incubation for 4hrs at 37°C/5% CO₂. The transfection mixture was removed by aspiration, followed by the addition of 25ml of DMEM/10% Fetal Bovine Serum. Cells were incubated at 37°C/5% CO₂. After 72hr incubation, cells were harvested and utilized for analysis.

Example 4

10 ASSAYS FOR DETERMINATION OF CONSTITUTIVE ACTIVITY OF NON-ENDOGENOUS GPCRS

A variety of approaches are available for assessment of constitutive activity of the non-endogenous human GPCRs. The following are illustrative; those of ordinary skill in the art are credited with the ability to determine those techniques that are preferentially beneficial for the needs of the artisan.

1. Membrane Binding Assays: [³⁵S]GTP γ S Assay

When a G protein-coupled receptor is in its active state, either as a result of ligand binding or constitutive activation, the receptor couples to a G protein and stimulates the release of GDP and subsequent binding of GTP to the G protein. The alpha subunit of the G protein-receptor complex acts as a GTPase and slowly hydrolyzes the GTP to GDP, at which point the receptor normally is deactivated. Constitutively activated receptors continue to exchange GDP for GTP. The non-hydrolyzable GTP analog, [³⁵S]GTP γ S, can be utilized to demonstrate enhanced binding of [³⁵S]GTP γ S to membranes expressing constitutively activated receptors. The advantage of using [³⁵S]GTP γ S binding to measure constitutive

activation is that: (a) it is generically applicable to all G protein-coupled receptors; (b) it is proximal at the membrane surface making it less likely to pick-up molecules which affect the intracellular cascade.

The assay utilizes the ability of G protein coupled receptors to stimulate [³⁵S]GTPγS binding to membranes expressing the relevant receptors. The assay can, therefore, be used in the direct identification method to screen candidate compounds to known, orphan and constitutively activated G protein-coupled receptors. The assay is generic and has application to drug discovery at all G protein-coupled receptors.

The [³⁵S]GTPγS assay can be incubated in 20 mM HEPES and between 1 and about 20mM MgCl₂ (this amount can be adjusted for optimization of results, although 20mM is preferred) pH 7.4, binding buffer with between about 0.3 and about 1.2 nM [³⁵S]GTPγS (this amount can be adjusted for optimization of results, although 1.2 is preferred) and 12.5 to 75 μg membrane protein (*e.g.* COS-7 cells expressing the receptor; this amount can be adjusted for optimization, although 75μg is preferred) and 1 μM GDP (this amount can be changed for optimization) for 1 hour. Wheatgerm agglutinin beads (25 μl; Amersham) should then be added and the mixture incubated for another 30 minutes at room temperature. The tubes are then centrifuged at 1500 x g for 5 minutes at room temperature and then counted in a scintillation counter.

A less costly but equally applicable alternative has been identified which also meets the needs of large scale screening. Flash plates™ and Wallac™ scintistrips may be utilized to format a high throughput [³⁵S]GTPγS binding assay. Furthermore, using this technique, the assay can be utilized for known GPCRs to simultaneously monitor tritiated ligand binding to the receptor at the same time as monitoring the efficacy via [³⁵S]GTPγS binding. This is

- 44 -

possible because the Wallac beta counter can switch energy windows to look at both tritium and ^{35}S -labeled probes. This assay may also be used to detect other types of membrane activation events resulting in receptor activation. For example, the assay may be used to monitor ^{32}P phosphorylation of a variety of receptors (both G protein coupled and tyrosine kinase receptors). When the membranes are centrifuged to the bottom of the well, the bound $[\text{S}]^{\text{35}}\text{GTP}\gamma\text{S}$ or the ^{32}P -phosphorylated receptor will activate the scintillant which is coated of the wells. Scinti[®] strips (Wallac) have been used to demonstrate this principle. In addition, the assay also has utility for measuring ligand binding to receptors using radioactively labeled ligands. In a similar manner, when the radiolabeled bound ligand is centrifuged to the bottom of the well, the scintistrip label comes into proximity with the radiolabeled ligand resulting in activation and detection.

2. Adenylyl Cyclase

A Flash Plate[™] Adenylyl Cyclase kit (New England Nuclear; Cat. No. SMP004A) designed for cell-based assays can be modified for use with crude plasma membranes. The Flash Plate wells contain a scintillant coating which also contains a specific antibody recognizing cAMP. The cAMP generated in the wells was quantitated by a direct competition for binding of radioactive cAMP tracer to the cAMP antibody. The following serves as a brief protocol for the measurement of changes in cAMP levels in membranes that express the receptors.

Transfected cells are harvested approximately three days after transfection. Membranes were prepared by homogenization of suspended cells in buffer containing 20mM HEPES, pH 7.4 and 10mM MgCl_2 . Homogenization is performed on ice using a Brinkman Polytron[™] for approximately 10 seconds. The resulting homogenate is centrifuged at 49,000

- 45 -

X g for 15 minutes at 4°C. The resulting pellet is then resuspended in buffer containing 20mM HEPES, pH 7.4 and 0.1 mM EDTA, homogenized for 10 seconds, followed by centrifugation at 49,000 X g for 15 minutes at 4°C. The resulting pellet can be stored at -80°C until utilized. On the day of measurement, the membrane pellet is slowly thawed at room temperature, resuspended in buffer containing 20mM HEPES, pH 7.4 and 10mM MgCl₂ (these amounts can be optimized, although the values listed herein are preferred), to yield a final protein concentration of 0.60mg/ml (the resuspended membranes were placed on ice until use).

cAMP standards and Detection Buffer (comprising 2 µCi of tracer [¹²⁵I] cAMP (100 µl) to 11 ml Detection Buffer) are prepared and maintained in accordance with the manufacturer's instructions. Assay Buffer is prepared fresh for screening and contained 20mM HEPES, pH 7.4, 10mM MgCl₂, 20mM (Sigma), 0.1 units/ml creatine phosphokinase (Sigma), 50 µM GTP (Sigma), and 0.2 mM ATP (Sigma); Assay Buffer can be stored on ice until utilized. The assay is initiated by addition of 50ul of assay buffer followed by addition of 50ul of membrane suspension to the NEN Flash Plate. The resultant assay mixture is incubated for 60 minutes at room temperature followed by addition of 100ul of detection buffer. Plates are then incubated an additional 2-4 hours followed by counting in a Wallac MicroBeta™ scintillation counter. Values of cAMP/well are extrapolated from a standard cAMP curve that is contained within each assay plate.

20 C. Reporter-Based Assays

1. CREB Reporter Assay (Gs-associated receptors)

A method to detect Gs stimulation depends on the known property of the transcription factor CREB, which is activated in a cAMP-dependent manner. A PathDetect™ CREB trans-

Reporting System (Stratagene, Catalogue # 219010) can utilized to assay for Gs coupled activity in 293 or 293T cells. Cells are transfected with the plasmids components of this above system and the indicated expression plasmid encoding endogenous or mutant receptor using a Mammalian Transfection Kit (Stratagene, Catalogue #200285) according to the manufacturer's instructions. Briefly, 400 ng pFR-Luc (luciferase reporter plasmid containing Gal4 recognition sequences), 40 ng pFA2-CREB (Gal4-CREB fusion protein containing the Gal4 DNA-binding domain), 80 ng pCMV-receptor expression plasmid (comprising the receptor) and 20 ng CMV-SEAP (secreted alkaline phosphatase expression plasmid; alkaline phosphatase activity is measured in the media of transfected cells to control for variations in transfection efficiency between samples) are combined in a calcium phosphate precipitate as per the Kit's instructions. Half of the precipitate is equally distributed over 3 wells in a 96-well plate, kept on the cells overnight, and replaced with fresh medium the following morning. Forty-eight (48) hr after the start of the transfection, cells are treated and assayed for, *e.g.*, luciferase activity

15 2. AP1 reporter assay (Gq-associated receptors)

A method to detect Gq stimulation depends on the known property of Gq-dependent phospholipase C to cause the activation of genes containing AP1 elements in their promoter. A Pathdetect™ AP-1 cis-Reporting System (Stratagene, Catalogue # 219073) can be utilized following the protocol set forth above with respect to the CREB reporter assay, except that the components of the calcium phosphate precipitate were 410 ng pAP1-Luc, 80 ng pCMV-receptor expression plasmid, and 20 ng CMV-SEAP.

3. CRE-LUC Reporter Assay

293 and 293T cells are plated-out on 96 well plates at a density of 2×10^4 cells per

well and were transfected using Lipofectamine Reagent (BRL) the following day according to manufacturer instructions. A DNA/lipid mixture is prepared for each 6-well transfection as follows: 260ng of plasmid DNA in 100 μ l of DMEM were gently mixed with 2 μ l of lipid in 100 μ l of DMEM (the 260ng of plasmid DNA consisted of 200ng of a 8xCRE-Luc reporter plasmid (see below and Figure 1 for a representation of a portion of the plasmid), 50ng of pCMV comprising endogenous receptor or non-endogenous receptor or pCMV alone, and 10ng of a GPRS expression plasmid (GPRS in pcDNA3 (Invitrogen)). The 8XCRE-Luc reporter plasmid was prepared as follows: vector.SRIF- β -gal was obtained by cloning the rat somatostatin promoter (-71/+51) at BglV-HindIII site in the p β gal-Basic Vector (Clontech).

10 Eight (8) copies of cAMP response element were obtained by PCR from an adenovirus template AdpCF126CCRE8 (see, 7 *Human Gene Therapy* 1883 (1996)) and cloned into the SRIF- β -gal vector at the Kpn-BglV site, resulting in the 8xCRE- β -gal reporter vector. The 8xCRE-Luc reporter plasmid was generated by replacing the beta-galactosidase gene in the 8xCRE- β -gal reporter vector with the luciferase gene obtained from the pGL3-basic vector

15 (Promega) at the HindIII-BamHI site. Following 30 min. incubation at room temperature, the DNA/lipid mixture was diluted with 400 μ l of DMEM and 100 μ l of the diluted mixture was added to each well. 100 μ l of DMEM with 10% FCS were added to each well after a 4hr incubation in a cell culture incubator. The following day the transfected cells were changed with 200 μ l/well of DMEM with 10% FCS. Eight (8) hours later, the wells were changed to

20 100 μ l /well of DMEM without phenol red, after one wash with PBS. Luciferase activity were measured the next day using the LucLite™ reporter gene assay kit (Packard) following manufacturer instructions and read on a 1450 MicroBeta™ scintillation and luminescence counter (Wallac).

4. SRF-LUC Reporter Assay

One method to detect Gq stimulation depends on the known property of Gq-dependent phospholipase C to cause the activation of genes containing serum response factors in their promoter. A Pathdetect™ SRF-Luc-Reporting System (Stratagene) can be utilized to assay for Gq coupled activity in, e.g., COS7 cells. Cells are transfected with the plasmid components of the system and the indicated expression plasmid encoding endogenous or non-endogenous GPCR using a Mammalian Transfection™ Kit (Stratagene, Catalogue #200285) according to the manufacturer's instructions. Briefly, 410 ng SRF-Luc, 80 ng pCMV-receptor expression plasmid and 20 ng CMV-SEAP (secreted alkaline phosphatase expression plasmid; alkaline phosphatase activity is measured in the media of transfected cells to control for variations in transfection efficiency between samples) are combined in a calcium phosphate precipitate as per the manufacturer's instructions. Half of the precipitate is equally distributed over 3 wells in a 96-well plate, kept on the cells in a serum free media for 24 hours. The last 5 hours the cells are incubated with 1 μ M Angiotensin, where indicated. Cells are then lysed and assayed for luciferase activity using a LucLite™ Kit (Packard, Cat. # 6016911) and "Trilux 1450 Microbeta" liquid scintillation and luminescence counter (Wallac) as per the manufacturer's instructions. The data can be analyzed using GraphPad Prism™ 2.0a (GraphPad Software Inc.).

5. Intracellular IP₃ Accumulation Assay

On day 1, cells comprising the receptors (endogenous and/or non-endogenous) can be plated onto 24 well plates, usually 1×10^5 cells/well (although this number can be optimized. On day 2 cells can be transfected by firstly mixing 0.25 μ g DNA in 50 μ l serum free DMEM/well and 2 μ l lipofectamine in 50 μ l serumfree DMEM/well. The solutions

are gently mixed and incubated for 15-30 min at room temperature. Cells are washed with 0.5 ml PBS and 400 μ l of serum free media is mixed with the transfection media and added to the cells. The cells are then incubated for 3-4 hrs at 37°C/5%CO₂ and then the transfection media is removed and replaced with 1ml/well of regular growth media. On

5 day 3 the cells are labeled with ³H-myo-inositol. Briefly, the media is removed and the cells are washed with 0.5 ml PBS. Then 0.5 ml inositol-free/serum free media (GIBCO-BRL) is added/well with 0.25 μ Ci of ³H-myo-inositol / well and the cells are incubated for 16-18 hrs o/n at 37°C/5%CO₂. On Day 4 the cells are washed with 0.5 ml PBS and 0.45 ml of assay medium is added containing inositol-free/serum free media 10 μ M pargyline

10 10 mM lithium chloride or 0.4 ml of assay medium and 50 μ l of 10x ketanserin (ket) to final concentration of 10 μ M. The cells are then incubated for 30 min at 37°C. The cells are then washed with 0.5 ml PBS and 200 μ l of fresh/icecold stop solution (1M KOH; 18 mM Na-borate; 3.8 mM EDTA) is added/well. The solution is kept on ice for 5-10 min or until cells were lysed and then neutralized by 200 μ l of fresh/ice cold neutralization sol.

15 (7.5 % HCL). The lysate is then transferred into 1.5 ml eppendorf tubes and 1 ml of chloroform/methanol (1:2) is added/tube. The solution is vortexed for 15 sec and the upper phase is applied to a Biorad AG1-X8™ anion exchange resin (100-200 mesh). Firstly, the resin is washed with water at 1:1.25 W/V and 0.9 ml of upper phase is loaded onto the column. The column is washed with 10 mls of 5 mM myo-inositol and 10 ml of 5

20 mM Na-borate/60mM Na-formate. The inositol tris phosphates are eluted into scintillation vials containing 10 ml of scintillation cocktail with 2 ml of 0.1 M formic acid/ 1 M ammonium formate. The columns are regenerated by washing with 10 ml of 0.1 M formic acid/3M ammonium formate and rinsed twice with dd H₂O and stored at 4°C in water.

Exemplary results are presented below in Table I:

TABLE I

Receptor	Mutation	Assay Utilized	Signal Generated: Endogenous Version (Relative Light Units)	Signal Generated: Non-Endogenous Version (Relative Light Units)	Percent Difference	
5	hAT1	F239K	SRF-LUC	34	137	75% ¹
		AT2K255IC3	SRF-LUC	34	127	73% ¹
	hTDAG8	I225K	CRE-LUC (293 cells)	2,715	14,440	81% ¹
		I225K	CRE-LUC (293T cells)	65,681	185,636	65% ¹
	hH9	F236K	CRE-LUC	1,887	6,096	69% ¹
	hCCKB	V332K	CRE-LUC	785	3,223	76% ¹

C. CELL-BASED DETECTION ASSAY (EXAMPLE -TDAG8)

10 293 cells were plated-out on 150mm plates at a density of 1.3×10^7 cells per plate, and were transfected using 12ug of the respective DNA and 60ul of Lipofectamine Reagent (BRL) per plate. The transfected cells were grown in media containing serum for an assay performed 24 hours post-transfection. For detection assay performed 48 hours post-transfection (assay comparing serum and serum-free media; see Figure 3), the initial media
 15 was changed to either serum or serum-free media. The serum-free media was comprised solely of Dulbecco's Modified Eagle's (DME) High Glucose Medium (Irvine Scientific #9024). In addition to the above DME Medium, the media with serum contained the following: 10% Fetal Bovine Serum (Hyclone #SH30071.03), 1% of 100mM Sodium Pyruvate (Irvine Scientific #9334), 1% of 20mM L-Glutamine (Irvine Scientific #9317), and 1% of Penicillin-

Streptomycin solution (Irvine Scientific #9366).

A 96-well Adenylyl Cyclase Activation Flashplate™ was used (NEN: #SMP004A). First, 50ul of the standards for the assay were added to the plate, in duplicate, ranging from concentrations of 50pmol to zero pmol cAMP per well. The standard cAMP (NEN: #SMP004A) was reconstituted in water, and serial dilutions were made using 1xPBS (Irvine Scientific: #9240). Next, 50ul of the stimulation buffer (NEN: #SMP004A) was added to all wells. In the case of using compounds to measure activation or inactivation of cAMP, 10ul of each compound, diluted in water, was added to its respective well, in triplicate. Various final concentrations used range from 1uM up to 1mM. Adenosine 5'-triphosphate, ATP, (Research Biochemicals International: #A-141) and Adenosine 5'-diphosphate, ADP, (Sigma: #A2754) were used in the assay. Next, the 293 cells transfected with the respective cDNA (CMV or TDAG8) were harvested 24 (assay detection in serum media) or 48 hours post-transfection (assay detection comparing serum and serum-free media). The media was aspirated and the cells washed once with 1xPBS. Then 5ml of 1xPBS was added to the cells along with 3ml of cell dissociation buffer (Sigma: #C-1544). The detached cells were transferred to a centrifuge tube and centrifuged at room temperature for five minutes. The supernatant was removed and the cell pellet was resuspended in an appropriate amount of 1xPBS to obtain a final concentration of 2×10^6 cells per milliliter. To the wells containing the compound, 50ul of the cells in 1xPBS (1×10^5 cells/well) were added. The plate was incubated on a shaker for 15 minutes at room temperature. The detection buffer containing the tracer cAMP was prepared. In 11ml of detection buffer (NEN: #SMP004A), 50ul (equal to 1uCi) of [125 I]cAMP (NEN: #SMP004A) was added. Following incubation, 50ul of this detection buffer containing tracer cAMP was added to each well. The plate was placed on a shaker and

- 52 -

incubated at room temperature for two hours. Finally, the solution from the wells of the plate were aspirated and the flashplate was counted using the Wallac MicroBeta™ scintillation counter.

In Figure 2A, ATP and ADP bind to endogenous TDAG8 resulting in an increase of cAMP of about 59% and about 55% respectively. Figure 2B evidences ATP and ADP binding to endogenous TDAG8 where endogenous TDAG8 was transfected and grown in serum and serum-free medium. ATP binding to endogenous TDAG8 grown in serum media evidences an increase in cAMP of about 65%, compared to the endogenous TDAG8 with no compounds; in serum-free media there was an increase of about 68%. ADP binding to endogenous TDAG8 in serum evidences about a 61% increase, while in serum-free ADP binding evidences an increase of about 62% increase. ATP and ADP bind to endogenous TDAG8 with an EC50 value of 139.8uM and 120.5uM, respectively (data not shown).

Although the results presented in Figure 2B indicate substantially the same results when serum and serum-free media were compared, our choice is to use a serum based media, although a serum-free media can also be utilized.

Example 6 **GPCR FUSION PROTEIN PREPARATION**

The design of the constitutively activated GPCR-G protein fusion construct was accomplished as follows: both the 5' and 3' ends of the rat G protein Gsα (long form; Itoh, H. et al., 83 *PNAS* 3776 (1986)) were engineered to include a HindIII (5'-AAGCTT-3') sequence thereon. Following confirmation of the correct sequence (including the flanking HindIII sequences), the entire sequence was shuttled into pcDNA3.1(-) (Invitrogen, cat. no. V795-20) by subcloning using the HindIII restriction site of that vector. The correct

- 53 -

orientation for the Gs α sequence was determined after subcloning into pcDNA3.1(-). The modified pcDNA3.1(-) containing the rat Gs α gene at HindIII sequence was then verified; this vector was now available as a "universal" Gs α protein vector. The pcDNA3.1(-) vector contains a variety of well-known restriction sites upstream of the HindIII site, thus
5 beneficially providing the ability to insert, upstream of the Gs protein, the coding sequence of an endogenous, constitutively active GPCR. This same approach can be utilized to create other "universal" G protein vectors, and, of course, other commercially available or proprietary vectors known to the artisan can be utilized – the important criteria is that the sequence for the GPCR be upstream and in-frame with that of the G protein.

10 TDAG8 couples via Gs, while H9 couples via Gz. For the following exemplary GPCR Fusion Proteins, fusion to Gs α was accomplished.

A TDAG8(I225K)-Gs α Fusion Protein construct was made as follows: primers were designed as follows:

5'-gatcTCTAGAATGAACAGCACATGTATTGAAG-3' (SEQ.ID.NO.: 125; sense)

15 5'-ctagGGTACCCGCTCAAGGACCTCTAATTCCATAG-3' (SEQ.ID.NO.: 126; antisense).

Nucleotides in lower caps are included as spacers in the restriction sites between the G protein and TDAG8. The sense and anti-sense primers included the restriction sites for XbaI and KpnI, respectively.

PCR was then utilized to secure the respective receptor sequences for fusion within
20 the Gs α universal vector disclosed above, using the following protocol for each: 100ng cDNA for TDAG8 was added to separate tubes containing 2ul of each primer (sense and anti-sense), 3uL of 10mM dNTPs, 10uL of 10XTaqPlus™ Precision buffer, 1uL of TaqPlus™ Precision polymerase (Stratagene: #600211), and 80uL of water. Reaction temperatures and cycle times for TDAG8 were as follows: the initial denaturing step was done at 94 °C for five minutes, and

a cycle of 94°C for 30 seconds; 55°C for 30 seconds; 72°C for two minutes. A final extension time was done at 72°C for ten minutes. PCR product for was run on a 1% agarose gel and then purified (data not shown). The purified product was digested with XbaI and KpnI (New England Biolabs) and the desired inserts purified and ligated into the Gs universal
5 vector at the respective restriction site. The positive clones was isolated following transformation and determined by restriction enzyme digest; expression using 293 cells was accomplished following the protocol set forth *infra*. Each positive clone for TDAG8:Gs – Fusion Protein was sequenced to verify correctness.

GPCR Fusion Proteins comprising non-endogenous, constitutively activated
10 TDAG8(I225K) were analyzed as above and verified for constitutive activation.

An H9(F236K)-Gsa Fusion Protein construct was made as follows: primers were designed as follows:

5'-TTAgatcGGGGCCCCACCCTAGCGGT-3' (SEQ.ID.NO.: 145; sense)

5'-ggtagcCCCACAGCCATTTTCATCAGGATC-3' (SEQ.ID.NO.: 146; antisense).

15 Nucleotides in lower caps are included as spacers in the restriction sites between the G protein and H9. The sense and anti-sense primers included the restriction sites for EcoRV and KpnI, respectively such that spacers (attributed to the restriction sites) exists between the G protein and H9.

PCR was then utilized to secure the respective receptor sequences for fusion within
20 the Gsa universal vector disclosed above, using the following protocol for each: 80ng cDNA for H9 was added to separate tubes containing 100ng of each primer (sense and anti-sense), and 45uL of PCR Supermix™ (Gibco-Brl, LifeTech) (50ul total reaction volume). Reaction temperatures and cycle times for H9 were as follows: the initial denaturing step was done it 94°C for one, and a cycle of 94°C for 30 seconds; 55°C for 30 seconds; 72°C for two

- 55 -

minutes. A final extension time was done at 72°C for seven minutes. PCR product for was run on a 1% agarose gel and then purified (data not shown). The purified product was cloned into pCRII-TOPO™ System followed by identification of positive clones. Positive clones were isolated, digested with EcoRV and KpnI (New England Biolabs) and the desired inserts were isolated, purified and ligated into the Gs universal vector at the respective restriction site. The positive clones was isolated following transformation and determined by restriction enzyme digest; expression using 293 cells was accomplished following the protocol set forth *infra*. Each positive clone for H9(F236K):Gs – Fusion Protein was sequenced to verify correctness. Membranes were frozen (-80°C) until utilized.

To ascertain the ability of measuring a cAMP response mediated by the Gs protein (even though H9 couples with Gz), the following cAMP membrane assay was utilized, based upon an NEN Adenyl Cyclase Activation Flahplate™ Assay kit (96 well format). "Binding Buffer" consisted of 10mM HEPES, 100mM NaCl and 10mM MgCl (ph 7.4). "Regeneration Buffer" was prepared in Binding Buffer and consisted of 20mM phosphocreatine, 20U creatine phosphokinase, 20uM GTP, 0.2mM ATP, and 0.6mM IBMX. "cAMP Standards" were prepared in Binding Buffer as follows:

		cAMP Stock (5,000 pmol/ml in 2ml H ₂ O) in ul	Added to indicted amount of Binding Buffer	Final Assay Concentration (50ul into 100ul) to achieve indicated pmol/well
20	A	250	1ml	50
	B	500 of A	500ul	25
	C	500 of B	500ul	12.5
	D	500 of C	750ul	5.0
	E	500 of D	500ul	2.5
25	F	500 of E	500ul	1.25
	G	500 of F	750ul	0.5

Frozen membranes (both pCMV as control and the non-endogenous H(-Gs Fusion Protein) were thawed (on ice at room temperature until in solution). Membranes were

- 56 -

homogenized with a polytron until in suspension (2 x 15 seconds). Membrane protein concentration was determined using the Bradford Assay Protocol (*see infra*). Membrane concentration was diluted to 0.5mg/ml in Regeneration Buffer (final assay concentration - 25ug/well). Thereafter, 50ul of Binding Buffer was added to each well. For control, 50ul/well of cAMP standard was added to wells 11 and 12 A-G, with Binding Buffer alone to 12H (on the 96-well format). Thereafter, 50ul/well of protein was added to the wells and incubated at room temperature (on shaker) for 60min. 100ul [¹²⁵I]cAMP in Detection Buffer (*see infra*) was added to each well (final - 50ul [¹²⁵I]cAMP into 11ml Detection Buffer). These were incubated for 2hrs at room temperature. Plates were aspirated with an 8 channel manifold and sealed with plate covers. Results (pmoles cAMP bound) were read in a Wallac™ 1450 on "prot #15). Results are presented in Figure 3.

The results presented in Figure 3 indicate that the Gs coupled fusion was able to "drive" the cyclase reaction such that measurement of the constitutive activation of H9(F236K) was viable. Based upon these results, the direct identification of candidate compounds that are inverse agonists, agonists and partial agonists is possible using a cyclase-based assay.

Example 6

Protocol: Direct Identification of Inverse Agonists and Agonists Using [³⁵S]GTPγS

Although we have utilized endogenous, constitutively active GPCRs for the direct identification of candidate compounds as, e.g., inverse agonists, for reasons that are not altogether understood, intra-assay variation can become exacerbated. Preferably, then, a GPCR Fusion Protein, as disclosed above, is also utilized with a non-endogenous, constitutively activated GPCR. We have determined that when such a protein is used, intra-assay variation appears to be substantially stabilized, whereby an effective signal-to-noise ratio is obtained. This has the beneficial result of allowing for a more robust identification

- 57 -

of candidate compounds. Thus, it is preferred that for direct identification, a GPCR Fusion Protein be used and that when utilized, the following assay protocols be utilized.

Membrane Preparation

Membranes comprising the non-endogenous, constitutively active orphan GPCR Fusion Protein of interest and for use in the direct identification of candidate compounds as inverse agonists, agonists or partial agonists are preferably prepared as follows:

a. Materials

"Membrane Scrape Buffer" is comprised of 20mM HEPES and 10mM EDTA, pH 7.4;

"Membrane Wash Buffer" is comprised of 20 mM HEPES and 0.1 mM EDTA, pH 7.4;

10 "Binding Buffer" is comprised of 20mM HEPES, 100 mM NaCl, and 10 mM MgCl₂, pH 7.4

b. Procedure

All materials are kept on ice throughout the procedure. Firstly, the media is aspirated from a confluent monolayer of cells, followed by rinse with 10ml cold PBS, followed by aspiration. Thereafter, 5ml of Membrane Scrape Buffer is added to scrape cells; this is followed by transfer of cellular extract into 50ml centrifuge tubes (centrifuged at 20,000 rpm for 17 minutes at 4°C). Thereafter, the supernatant is aspirated and the pellet is resuspended in 30ml Membrane Wash Buffer followed by centrifuge at 20,000 rpm for 17 minutes at 4°C. The supernatant is then aspirated and the pellet resuspended in Binding Buffer. This is then homogenized using a Brinkman polytron™ homogenizer (15-20 second bursts until the all material is in suspension). This is referred to herein as "Membrane Protein".

Bradford Protein Assay

Following the homogenization, protein concentration of the membranes is determined using the Bradford Protein Assay (protein can be diluted to about 1.5mg/ml, aliquoted and

- 58 -

frozen (-80°C) for later use; when frozen, protocol for use is as follows: on the day of the assay, frozen Membrane Protein is thawed at room temperature, followed by vortex and then homogenized with a polytron at about 12 x 1,000 rpm for about 5-10 seconds; it is noted that for multiple preparations, the homogenizer should be thoroughly cleaned between
5 homogenization of different preparations).

a. Materials

Binding Buffer (as per above); Bradford Dye Reagent; Bradford Protein Standard are utilized, following manufacturer instructions (Biorad, cat. no. 500-0006).

b. Procedure

10 Duplicate tubes are prepared, one including the membrane, and one as a control "blank". Each contained 800ul Binding Buffer. Thereafter, 10ul of Bradford Protein Standard (1mg/ml) is added to each tube, and 10ul of membrane Protein is then added to just one tube (not the blank). Thereafter, 200ul of Bradford Dye Reagent is added to each tube, followed by vortex of each. After five (5) minutes, the tubes were re-vortexed and the material therein
15 is transferred to cuvettes. The cuvettes are then read using a CECIL 3041 spectrophotometer, at wavelength 595.

Direct Identification Assay

a. Materials

GDP Buffer consists of 37.5 ml Binding Buffer and 2mg GDP (Sigma, cat. no. G-
20 7127), followed by a series of dilutions in Binding Buffer to obtain 0.2 uM GDP (final concentration of GDP in each well was 0.1 uM GDP); each well comprising a candidate compound, has a final volume of 200ul consisting of 100ul GDP Buffer (final concentration, 0.1uM GDP), 50ul Membrane Protein in Binding Buffer, and 50ul [^{35}S]GTP γ S (0.6 nM) in

- 59 -

Binding Buffer (2.5 ul [³⁵S]GTPγS per 10ml Binding Buffer).

b. Procedure

Candidate compounds are preferably screened using a 96-well plate format (these can be frozen at -80°C). Membrane Protein (or membranes with expression vector excluding the GPCR Fusion Protein, as control), are homogenized briefly until in suspension. Protein concentration is then determined using the Bradford Protein Assay set forth above. Membrane Protein (and control) is then diluted to 0.25mg/ml in Binding Buffer (final assay concentration, 12.5ug/well). Thereafter, 100 ul GDP Buffer is added to each well of a Wallac Scintistrip™ (Wallac). A 5ul pin-tool is then used to transfer 5 ul of a candidate compound into such well (i.e., 5ul in total assay volume of 200 ul is a 1:40 ratio such that the final screening concentration of the candidate compound is 10uM). Again, to avoid contamination, after each transfer step the pin tool should be rinsed in three reservoirs comprising water (1X), ethanol (1X) and water (2X) – excess liquid should be shaken from the tool after each rinse and dried with paper and kimwipes. Thereafter, 50 ul of Membrane Protein is added to each well (a control well comprising membranes without the GPCR Fusion Protein is also utilized), and pre-incubated for 5-10 minutes at room temperature. Thereafter, 50 ul of [³⁵S]GTPγS (0.6 nM) in Binding Buffer is added to each well, followed by incubation on a shaker for 60 minutes at room temperature (again, in this example, plates were covered with foil). The assay is then stopped by spinning of the plates at 4000 RPM for 15 minutes at 22°C. The plates are then aspirated with an 8 channel manifold and sealed with plate covers. The plates are then read on a Wallac 1450 using setting "Prot. #37" (as per manufacturer instructions).

Example 7

Protocol: Confirmation Assay

Using an independent assay approach to provide confirmation of a directly identified

- 60 -

candidate compound as set forth above, it is preferred that a confirmation assay then be utilized. In this case, the preferred confirmation assay is a cyclase-based assay.

A modified Flash Plate™ Adenylyl Cyclase kit (New England Nuclear; Cat. No. SMP004A) is preferably utilized for confirmation of candidate compounds directly identified
5 as inverse agonists and agonists to non-endogenous, constitutively activated orphan GPCRs in accordance with the following protocol.

Transfected cells are harvested approximately three days after transfection. Membranes are prepared by homogenization of suspended cells in buffer containing 20mM HEPES, pH 7.4 and 10mM MgCl₂. Homogenization is performed on ice using a Brinkman
10 Polytron™ for approximately 10 seconds. The resulting homogenate is centrifuged at 49,000 X g for 15 minutes at 4°C. The resulting pellet is then resuspended in buffer containing 20mM HEPES, pH 7.4 and 0.1 mM EDTA, homogenized for 10 seconds, followed by centrifugation at 49,000 X g for 15 minutes at 4°C. The resulting pellet can be stored at -80°C until utilized. On the day of direct identification screening, the membrane pellet is
15 slowly thawed at room temperature, resuspended in buffer containing 20mM HEPES, pH 7.4 and 10mM MgCL₂, to yield a final protein concentration of 0.60mg/ml (the resuspended membranes are placed on ice until use).

cAMP standards and Detection Buffer (comprising 2 μCi of tracer [¹²⁵I] cAMP (100 μl] to 11 ml Detection Buffer) are prepared and maintained in accordance with the
20 manufacturer's instructions. Assay Buffer is prepared fresh for screening and contained 20mM HEPES, pH 7.4, 10mM MgCl₂, 20mM phosphocreatine (Sigma), 0.1 units/ml creatine phosphokinase (Sigma), 50 μM GTP (Sigma), and 0.2 mM ATP (Sigma); Assay Buffer can be stored on ice until utilized.

- 61 -

Candidate compounds identified as per above (if frozen, thawed at room temperature) are added, preferably, to 96-well plate wells ($3\mu\text{l}$ /well; $12\mu\text{M}$ final assay concentration), together with $40\mu\text{l}$ Membrane Protein ($30\mu\text{g}$ /well) and $50\mu\text{l}$ of Assay Buffer. This admixture is then incubated for 30 minutes at room temperature, with gentle shaking.

5 Following the incubation, $100\mu\text{l}$ of Detection Buffer is added to each well, followed by incubation for 2-24 hours. Plates are then counted in a Wallac MicroBeta™ plate reader using "Prot. #31" (as per manufacturer instructions).

It is intended that each of the patents, applications, and printed publications mentioned in this patent document be hereby incorporated by reference in their entirety.

10 As those skilled in the art will appreciate, numerous changes and modifications may be made to the preferred embodiments of the invention without departing from the spirit of the invention. It is intended that all such variations fall within the scope of the invention.

Although a variety of expression vectors are available to those in the art, for purposes of utilization for both the endogenous and non-endogenous human GPCRs, it is most preferred that the vector utilized be pCMV. This vector was deposited with the American Type Culture Collection (ATCC) on October 13, 1998 (10801 University Blvd., Manassas, VA 20110-2209 USA) under the provisions of the Budapest Treaty for the International Recognition of the Deposit of Microorganisms for the Purpose of Patent Procedure. The DNA was tested by the ATCC and determined to be. The ATCC has
20 assigned the following deposit number to pCMV: ATCC #203351.

CLAIMS

What is claimed is:

1. A cDNA encoding a non-endogenous, constitutively activated version of a human G protein-coupled receptor comprising hARE-3(F313K).
- 5 2. A non-endogenous version of a human G protein-coupled receptor encoded by the cDNA of claim 1.
3. A Plasmid comprising a Vector and the cDNA of claim 1.
4. A Host Cell comprising the Plasmid of claim 3.
5. A cDNA encoding a non-endogenous, constitutively activated version of a human
10 G protein-coupled receptor comprising hARE-4(V233K)
6. A non-endogenous version of a human G protein-coupled receptor encoded by the cDNA of claim 5.
7. A Plasmid comprising a Vector and the cDNA of claim 5.
8. A Host Cell comprising the Plasmid of claim 7.
- 15 9. A cDNA encoding a non-endogenous, constitutively activated version of a human G protein-coupled receptor comprising hARE-5(A240K).
10. A non-endogenous version of a human G protein-coupled receptor encoded by the cDNA of claim 9.
11. A Plasmid comprising a Vector and the cDNA of claim 5.
- 20 12. A Host Cell comprising the Plasmid of claim 11.
13. A cDNA encoding a non-endogenous, constitutively activated version of a human G protein-coupled receptor comprising hGPCR14(L257K).

14. A non-endogenous version of a human G protein-coupled receptor encoded by the cDNA of claim 13.
15. A Plasmid comprising a Vector and the cDNA of claim 13.
- 5 16. A Host Cell comprising the Plasmid of claim 15.
17. A cDNA encoding a non-endogenous, constitutively activated version of a human G protein-coupled receptor comprising hGPCR27(C283K).
18. A non-endogenous version of a human G protein-coupled receptor encoded by the cDNA of claim 17.
- 10 19. A Plasmid comprising a Vector and the cDNA of claim 17.
20. A Host Cell comprising the Plasmid of claim 19.
21. A cDNA encoding a non-endogenous, constitutively activated version of a human G protein-coupled receptor comprising hARE-1(E232K).
22. A non-endogenous version of a human G protein-coupled receptor encoded by the cDNA of claim 21.
- 15 23. A Plasmid comprising a Vector and the cDNA of claim 21.
24. A Host Cell comprising the Plasmid of claim 23.
25. A cDNA encoding a non-endogenous, constitutively activated version of a human G protein-coupled receptor comprising hARE-2(G285K).
- 20 26. A non-endogenous version of a human G protein-coupled receptor encoded by the cDNA of claim 25.
27. A Plasmid comprising a Vector and the cDNA of claim 25.
28. A Host Cell comprising the Plasmid of claim 27.

- 64 -

29. A cDNA encoding a non-endogenous, constitutively activated version of a human G protein-coupled receptor comprising hPPR1(L239K).

30. A non-endogenous version of a human G protein-coupled receptor encoded by the cDNA of claim 29.

5 31. A Plasmid comprising a Vector and the cDNA of claim 29.

32. A Host Cell comprising the Plasmid of claim 31.

33. A cDNA encoding a non-endogenous, constitutively activated version of a human G protein-coupled receptor comprising hG2A(K232A).

10 34. A non-endogenous version of a human G protein-coupled receptor encoded by the cDNA of claim 33.

35. A Plasmid comprising a Vector and the cDNA of claim 33.

36. A Host Cell comprising the Plasmid of claim 35.

37. A cDNA encoding a non-endogenous, constitutively activated version of a human G protein-coupled receptor comprising hRUP3(L224K).

15 38. A non-endogenous version of a human G protein-coupled receptor encoded by the cDNA of claim 37.

39. A Plasmid comprising a Vector and the cDNA of claim 37.

40. A Host Cell comprising the Plasmid of claim 39.

20 41. A cDNA encoding a non-endogenous, constitutively activated version of a human G protein-coupled receptor comprising hRUP5(A236K).

42. A non-endogenous version of a human G protein-coupled receptor encoded by the cDNA of claim 41.

43. A Plasmid comprising a Vector and the cDNA of claim 41.

- 65 -

44. A Host Cell comprising the Plasmid of claim 42.
45. A cDNA encoding a non-endogenous, constitutively activated version of a human
G protein-coupled receptor comprising hRUP6(N267K)
46. A non-endogenous version of a human G protein-coupled receptor encoded by the
5 cDNA of claim 45.
47. A Plasmid comprising a Vector and the cDNA of claim 45.
48. A Host Cell comprising the Plasmid of claim 47.
49. A cDNA encoding a non-endogenous, constitutively activated version of a human
G protein-coupled receptor comprising hRUP7(A302K).
- 10 50. A non-endogenous version of a human G protein-coupled receptor encoded by the
cDNA of claim 49.
51. A Plasmid comprising a Vector and the cDNA of claim 49.
52. A Host Cell comprising the Plasmid of claim 51.
53. A cDNA encoding a non-endogenous, constitutively activated version of a human
15 G protein-coupled receptor comprising hCHN4(V236K).
54. A non-endogenous version of a human G protein-coupled receptor encoded by the
cDNA of claim 53.
55. A Plasmid comprising a Vector and the cDNA of claim 53.
56. A Host Cell comprising the Plasmid of claim 55.
- 20 57. A cDNA encoding a non-endogenous, constitutively activated version of a human
G protein-coupled receptor comprising hMC4(A244K).
58. A non-endogenous version of a human G protein-coupled receptor encoded by the
cDNA of claim 57.

- 66 -

59. A Plasmid comprising a Vector and the cDNA of claim 57.
60. A Host Cell comprising the Plasmid of claim 60.
61. A cDNA encoding a non-endogenous, constitutively activated version of a human G protein-coupled receptor comprising hCHN3(S284K).
- 5 62. A non-endogenous version of a human G protein-coupled receptor encoded by the cDNA of claim 61.
63. A Plasmid comprising a Vector and the cDNA of claim 61.
64. A Host Cell comprising the Plasmid of claim 63.
65. A cDNA encoding a non-endogenous, constitutively activated version of a human
- 10 G protein-coupled receptor comprising hCHN6(L352K).
66. A non-endogenous version of a human G protein-coupled receptor encoded by the cDNA of claim 65.
67. A Plasmid comprising a Vector and the cDNA of claim 65.
68. A Host Cell comprising the Plasmid of claim 67.
- 15 69. A cDNA encoding a non-endogenous, constitutively activated version of a human G protein-coupled receptor comprising hCHN8(N235K).
70. A non-endogenous version of a human G protein-coupled receptor encoded by the cDNA of claim 69.
71. A Plasmid comprising a Vector and the cDNA of claim 69.
- 20 72. A Host Cell comprising the Plasmid of claim 71.
73. A cDNA encoding a non-endogenous, constitutively activated version of a human G protein-coupled receptor comprising hH9(F236K).
74. A non-endogenous version of a human G protein-coupled receptor encoded by the

cDNA of claim 73.

75. A Plasmid comprising a Vector and the cDNA of claim 73.

76. A Host Cell comprising the Plasmid of claim 74.

77. A cDNA encoding a non-endogenous, constitutively activated version of a human

5 G protein-coupled AT1-receptor selected from the group consisting of:

hAT1(F239K); hAT1(N111A); hAT1(AT2K255IC3); and hAT1(A243+).

78. A non-endogenous version of a human G protein-coupled receptor encoded by a
cDNA of claim 77.

79. A Plasmid comprising a Vector and the cDNA of claim 77.

10 80. A Host Cell comprising the Plasmid of claim 79.

1/2




FIG. 1

2/2




FIG. 2A




FIG. 2B




FIG. 3

- 1 -

SEQUENCE LISTING

(1) GENERAL INFORMATION:

- 5 (i) APPLICANT: Behan, Dominic P.
Lehmann-Bruinsma, Karin
Chalmers, Derek T.
Lowitz, Kevin P.
Lin, I-Lin
10 Dang, Huong T.
Chen, Ruoping
Liaw, Chen W.
Gore, Martin J.
White, Carol
- 15 (ii) TITLE OF INVENTION: Non-Endogenous, Constitutively Activated Human G
Protein-Coupled Receptors
- (iii) NUMBER OF SEQUENCES: 146
- (iv) CORRESPONDENCE ADDRESS:
20 (A) ADDRESSEE: Arena Pharmaceuticals, Inc.
(B) STREET: 6166 Nancy Ridge Drive
(C) CITY: San Diego
(D) STATE: CA
(E) COUNTRY: USA
(F) ZIP: 92121
- 25 (v) COMPUTER READABLE FORM:
(A) MEDIUM TYPE: Floppy disk
(B) COMPUTER: IBM PC compatible
(C) OPERATING SYSTEM: PC-DOS/MS-DOS
(D) SOFTWARE: PatentIn Release #1.0, Version #1.30
- 30 (vi) CURRENT APPLICATION DATA:
(A) APPLICATION NUMBER: US
(B) FILING DATE:
(C) CLASSIFICATION:
- 35 (vii) ATTORNEY/AGENT INFORMATION:
(A) NAME: Burgoon, Richard P.
(B) REGISTRATION NUMBER: 34,787
- (ix) TELECOMMUNICATION INFORMATION:
(A) TELEPHONE: (858)453-7200
(B) TELEFAX: (858)453-7210
- 40 (2) INFORMATION FOR SEQ ID NO:1:
- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 1260 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single

- 2 -

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:1:

	ATGGTCTTCT CGGCAGTGTT GACTGCGTTC CATACCGGGA CATCCAACAC AACATTGTC	60
5	GTGTATGAAA ACACCTACAT GAATATTACA CTCCCTCCAC CATTCCAGCA TCCTGACCTC	120
	AGTCCATTGC TTAGATATAG TTTTGAAACC ATGGCTCCCA CTGGTTTGAG TTCCTTGACC	180
	GTGAATAGTA CAGCTGTGCC CACAACACCA GCAGCATTTA AGAGCCTAAA CTGCGCTCTT	240
	CAGATCACCC TTTCTGCTAT AATGATATTC ATTCTGTTTG TGTCTTTTCT TGGGAACCTG	300
	GTTGTTTGCC TCATGGTTTA CCAAAAAGCT GCCATGAGGT CTGCAATTAA CATCCTCCTT	360
10	GCCAGCCTAG CTTTTGCAGA CATGTTGCTT GCAGTGCTGA ACATGCCCTT TGCCCTGGTA	420
	ACTATTCTTA CTACCCGATG GATTTTTGGG AAATTCTTCT GTAGGGTATC TGCTATGTTT	480
	TTCTGGTTAT TTGTGATAGA AGGAGTAGCC ATCCTGCTCA TCATTAGCAT AGATAGGTTT	540
	CTTATTATAG TCCAGAGGCA GGATAAGCTA AACCCATATA GAGCTAAGGT TCTGATTGCA	600
	GTTTCTTGGG CAACTTCCTT TTGTGTAGCT TTTCTTTTGG CCGTAGGAAA CCCCAGCTG	660
15	CAGATACCTT CCCGAGCTCC CCAGTGTGTG TTTGGGTACA CAACCAATCC AGGCTACCAG	720
	GCTTATGTGA TTTTGATTTC TCTCATTCTT TTCTTCATAC CCTTCCTGGT AATACTGTAC	780
	TCATTTATGG GCATACTCAA CACCCTTCGG CACAATGCCT TGAGGATCCA TAGCTACCCT	840
	GAAGGTATAT GCCTCAGCCA GGCCAGCAAA CTGGGTCTCA TGAGTCTGCA GAGACCTTTC	900
	CAGATGAGCA TTGACATGGG CTTTAAACAA CGTGCCTTCA CCACTATTTT GATTCTCTTT	960
20	GCTGTCTTCA TTGTCTGCTG GGCCCCATTC ACCACTTACA GCCTTGTGGC AACATTCAGT	1020
	AAGCACTTTT ACTATCAGCA CAACTTTTTT GAGATTAGCA CCTGGCTACT GTGGCTCTGC	1080
	TACCTCAAGT CTGCATTGAA TCCGCTGATC TACTACTGGA GGATTAAGAA ATTCCATGAT	1140
	GCTTGCTTGG ACATGATGCC TAAGTCCTTC AAGTTTTTGC CGCAGCTCCC TGGTCACACA	1200
	AAGCGACGGA TACGTCTTAG TGCTGTCTAT GTGTGTGGGG AACATCGGAC GGTGGTGTGA	1260

25 (3) INFORMATION FOR SEQ ID NO:2:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 419 amino acids

(B) TYPE: amino acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

30

- 3 -

(ii) MOLECULE TYPE: DNA (genomic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:2:

	Met	Val	Phe	Ser	Ala	Val	Leu	Thr	Ala	Phe	His	Thr	Gly	Thr	Ser	Asn	
	1				5					10					15		
5	Thr	Thr	Phe	Val	Val	Tyr	Glu	Asn	Thr	Tyr	Met	Asn	Ile	Thr	Leu	Pro	
				20					25					30			
	Pro	Pro	Phe	Gln	His	Pro	Asp	Leu	Ser	Pro	Leu	Leu	Arg	Tyr	Ser	Phe	
				35				40					45				
10	Glu	Thr	Met	Ala	Pro	Thr	Gly	Leu	Ser	Ser	Leu	Thr	Val	Asn	Ser	Thr	
		50					55					60					
	Ala	Val	Pro	Thr	Thr	Pro	Ala	Ala	Phe	Lys	Ser	Leu	Asn	Leu	Pro	Leu	
	65					70				75					80		
	Gln	Ile	Thr	Leu	Ser	Ala	Ile	Met	Ile	Phe	Ile	Leu	Phe	Val	Ser	Phe	
				85						90					95		
15	Leu	Gly	Asn	Leu	Val	Val	Cys	Leu	Met	Val	Tyr	Gln	Lys	Ala	Ala	Met	
				100					105						110		
	Arg	Ser	Ala	Ile	Asn	Ile	Leu	Leu	Ala	Ser	Leu	Ala	Phe	Ala	Asp	Met	
				115				120					125				
20	Leu	Leu	Ala	Val	Leu	Asn	Met	Pro	Phe	Ala	Leu	Val	Thr	Ile	Leu	Thr	
		130					135						140				
	Thr	Arg	Trp	Ile	Phe	Gly	Lys	Phe	Phe	Cys	Arg	Val	Ser	Ala	Met	Phe	
	145					150					155				160		
	Phe	Trp	Leu	Phe	Val	Ile	Glu	Gly	Val	Ala	Ile	Leu	Leu	Ile	Ile	Ser	
				165						170					175		
25	Ile	Asp	Arg	Phe	Leu	Ile	Ile	Val	Gln	Arg	Gln	Asp	Lys	Leu	Asn	Pro	
				180					185					190			
	Tyr	Arg	Ala	Lys	Val	Leu	Ile	Ala	Val	Ser	Trp	Ala	Thr	Ser	Phe	Cys	
			195				200						205				
30	Val	Ala	Phe	Pro	Leu	Ala	Val	Gly	Asn	Pro	Asp	Leu	Gln	Ile	Pro	Ser	
		210					215					220					
	Arg	Ala	Pro	Gln	Cys	Val	Phe	Gly	Tyr	Thr	Thr	Asn	Pro	Gly	Tyr	Gln	
	225				230						235				240		
	Ala	Tyr	Val	Ile	Leu	Ile	Ser	Leu	Ile	Ser	Phe	Phe	Ile	Pro	Phe	Leu	
				245						250				255			
35	Val	Ile	Leu	Tyr	Ser	Phe	Met	Gly	Ile	Leu	Asn	Thr	Leu	Arg	His	Asn	
				260					265					270			

- 4 -

Ala Leu Arg Ile His Ser Tyr Pro Glu Gly Ile Cys Leu Ser Gln Ala
 275 280 285

Ser Lys Leu Gly Leu Met Ser Leu Gln Arg Pro Phe Gln Met Ser Ile
 290 295 300

5 Asp Met Gly Phe Lys Thr Arg Ala Phe Thr Thr Ile Leu Ile Leu Phe
 305 310 315 320

Ala Val Phe Ile Val Cys Trp Ala Pro Phe Thr Thr Tyr Ser Leu Val
 325 330 335

10 Ala Thr Phe Ser Lys His Phe Tyr Tyr Gln His Asn Phe Phe Glu Ile
 340 345 350

Ser Thr Trp Leu Leu Trp Leu Cys Tyr Leu Lys Ser Ala Leu Asn Pro
 355 360 365

Leu Ile Tyr Tyr Trp Arg Ile Lys Lys Phe His Asp Ala Cys Leu Asp
 370 375 380

15 Met Met Pro Lys Ser Phe Lys Phe Leu Pro Gln Leu Pro Gly His Thr
 385 390 395 400

Lys Arg Arg Ile Arg Pro Ser Ala Val Tyr Val Cys Gly Glu His Arg
 405 410 415

Thr Val Val

20

(4) INFORMATION FOR SEQ ID NO:3:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 1119 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear
- 25

(ii) MOLECULE TYPE: DNA (genomic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:3:

ATGTTAGCCA ACAGCTCCTC AACCAACAGT TCTGTTCTCC CGTGTCTCTGA CTACCGACCT 60

30 ACCCACC GCC TGCACCTGGT GGTCTACAGC TTGGTGCTGG CTGCCGGGCT CCCCCTCAAC 120

GCGCTAGCCC TCTGGGTCTT CCTGCGCGCG CTGCGCGTGC ACTCGGTGGT GAGCGTGTAC 180

ATGTGTAACC TGGCGGCCAG CGACCTGCTC TTCACCTCT CTGCTGCCCGT TCGTCTCTCC 240

TACTACGCAC TGCACCACTG GCCCTTCCCC GACCTCCTGT GCCAGACGAC GGGCGCCATC 300

TTCCAGATGA ACATGTACGG CAGCTGCATC TTCCTGATGC TCATCAACGT GGACCGCTAC 360

- 5 -

GCCGCCATCG TGCACCCGCT GCGACTGCGC CACCTGCGGC GGCCCCGCGT GGCGCGGCTG 420
 CTCTGCCTGG GCGTGTGGGC GCTCATCCTG GTGTTTGCCG TGCCCGCCGC CCGCGTGCAC 480
 AGGCCCTCGC GTTGCCGCTA CCGGGACCTC GAGGTGCGCC TATGCTTCGA GAGCTTCAGC 540
 GACGAGCTGT GGAAAGGCAG GCTGCTGCCC CTCGTGCTGC TGGCCGAGGC GCTGGGCTTC 600
 5 CTGCTGCCCC TGGCGGCGGT GGTCTACTCG TCGGGCCGAG TCTTCTGGAC GCTGGCGCGC 660
 CCCGACGCCA CGCAGAGCCA GCGGCGGCGG AAGACCGTGC GCCTCCTGCT GGCTAACCTC 720
 GTCATCTTCC TGCTGTGCTT CGTGCCCTAC AACAGCACGC TGGCGGTCTA CGGGCTGCTG 780
 CGGAGCAAGC TGGTGGCGGC CAGCGTGCCT GCCCGCGATC GCGTGCGCGG GGTGCTGATG 840
 GTGATGGTGC TGCTGGCCGG CGCCAACCTG GTGCTGGACC CGCTGGTGTA CTACTTTAGC 900
 10 GCCGAGGGCT TCCGCAACAC CCTGCGCGGC CTGGGCACTC CGCACCAGGC CAGGACCTCG 960
 GCCACCAACG GGACGCGGGC GGCCTCGCG CAATCCGAAA GGTCCGCCGT CACCACCGAC 1020
 GCCACCAGGC CGGATGCCGC CAGTCAGGGG CTGCTCCGAC CCTCCGACTC CCACTCTCTG 1080
 TCTTCCTTCA CACAGTGTCC CCAGGATTCC GCCCTCTGA 1119

(5) INFORMATION FOR SEQ ID NO:4:

- 15 (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 372 amino acids
 (B) TYPE: amino acid
 (C) STRANDEDNESS:
 (D) TOPOLOGY: not relevant

- 20 (ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:4:

Met Leu Ala Asn Ser Ser Ser Thr Asn Ser Ser Val Leu Pro Cys Pro
 1 5 10 15
 Asp Tyr Arg Pro Thr His Arg Leu His Leu Val Val Tyr Ser Leu Val
 25 20 25 30
 Leu Ala Ala Gly Leu Pro Leu Asn Ala Leu Ala Leu Trp Val Phe Leu
 35 40 45
 Arg Ala Leu Arg Val His Ser Val Val Ser Val Tyr Met Cys Asn Leu
 50 55 60
 30 Ala Ala Ser Asp Leu Leu Phe Thr Leu Ser Leu Pro Val Arg Leu Ser
 65 70 75 80
 Tyr Tyr Ala Leu His His Trp Pro Phe Pro Asp Leu Leu Cys Gln Thr

- 6 -

	85	90	95
	Thr Gly Ala Ile Phe Gln Met Asn Met Tyr Gly Ser Cys Ile Phe Leu		
	100	105	110
5	Met Leu Ile Asn Val Asp Arg Tyr Ala Ala Ile Val His Pro Leu Arg		
	115	120	125
	Leu Arg His Leu Arg Arg Pro Arg Val Ala Arg Leu Leu Cys Leu Gly		
	130	135	140
	Val Trp Ala Leu Ile Leu Val Phe Ala Val Pro Ala Ala Arg Val His		
	145	150	155
10	Arg Pro Ser Arg Cys Arg Tyr Arg Asp Leu Glu Val Arg Leu Cys Phe		
	165	170	175
	Glu Ser Phe Ser Asp Glu Leu Trp Lys Gly Arg Leu Leu Pro Leu Val		
	180	185	190
15	Leu Leu Ala Glu Ala Leu Gly Phe Leu Leu Pro Leu Ala Ala Val Val		
	195	200	205
	Tyr Ser Ser Gly Arg Val Phe Trp Thr Leu Ala Arg Pro Asp Ala Thr		
	210	215	220
	Gln Ser Gln Arg Arg Arg Lys Thr Val Arg Leu Leu Leu Ala Asn Leu		
	225	230	235
20	Val Ile Phe Leu Leu Cys Phe Val Pro Tyr Asn Ser Thr Leu Ala Val		
	245	250	255
	Tyr Gly Leu Leu Arg Ser Lys Leu Val Ala Ala Ser Val Pro Ala Arg		
	260	265	270
25	Asp Arg Val Arg Gly Val Leu Met Val Met Val Leu Leu Ala Gly Ala		
	275	280	285
	Asn Cys Val Leu Asp Pro Leu Val Tyr Tyr Phe Ser Ala Glu Gly Phe		
	290	295	300
	Arg Asn Thr Leu Arg Gly Leu Gly Thr Pro His Arg Ala Arg Thr Ser		
	305	310	315
30	Ala Thr Asn Gly Thr Arg Ala Ala Leu Ala Gln Ser Glu Arg Ser Ala		
	325	330	335
	Val Thr Thr Asp Ala Thr Arg Pro Asp Ala Ala Ser Gln Gly Leu Leu		
	340	345	350
35	Arg Pro Ser Asp Ser His Ser Leu Ser Ser Phe Thr Gln Cys Pro Gln		
	355	360	365
	Asp Ser Ala Leu		
	370		

- 7 -

(6) INFORMATION FOR SEQ ID NO:5:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 1107 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:5:

ATGGCCAACCT CCACAGGGCT GAACGCCTCA GAAGTCGCAG GCTCGTTGGG GTTGATCCTG 60
10 GCAGCTGTCTG TGGAGGTGGG GGCAGTCTG GGCACCGCG CGCTGCTGGT CGTGGTGTCTG 120
CGCAGCCCGG GACTGCGCGA CGCGCTCTAC CTGGCGCAC TGTGCGTCGT GGACCTGTCTG 180
GCGGCCCGCT CCATCATGCC GCTGGGCCTG CTGGCCGCAC CGCCGCCCGG GCTGGGCCCGC 240
GTGCGCCTGG GCGCCGCGCC ATGCCGCGCC GCTCGCTTCC TCTCCGCCGC TCTGCTGCCG 300
GCCTGCACGC TCGGGGTGGC CGCACTTGGC CTGGCAGCT ACCGCCTCAT CGTGCACCCG 360
15 CTGCGGCCAG GCTCGCGGCC GCCGCCTGTG CTCGTGCTCA CCGCCGTGTG GGCCGCGGCG 420
GGACTGCTGG GCGCGETCTC CCTGCTCGGC CCGCCGCCCG CACCGCCCC TGCTCCTGCT 480
CGCTGCTCGG TCCTGGCTGG GGCCTCGGG CCCTTCCGGC CGCTCTGGGC CCTGCTGGCC 540
TTCGCGCTGC CCGCCCTCCT GCTGCTCGGC GCCTACGGCG GCATCTTCGT GGTGGCGCGT 600
CGCGCTGCCC TGAGGCCCCC ACGGCCGGCG CGCGGGTCCC GACTCCGCTC GGACTCTCTG 660
20 GATAGCCGCC TTTCCATCTT GCCGCCGCTC CGGCCTCGCC TGCCCGGGGG CAAGGCGGCC 720
CTGGCCCCAG CGCTGGCCGT GGGCCAATTT GCAGCCTGCT GGCTGCCTTA TGGCTGCGCG 780
TGCCTGGCGC CCGCAGCGCG GGCCGCGGAA GCCGAAGCGG CTGTCACCTG GGTGCGCTAC 840
TCGGCCTTCG CGGCTCAGCC CTTCCTGTAC GGGCTGCTGC AGCGCCCCGT GCGCTTGGCA 900
CTGGGCCCGC TCTCTCGCCG TGCAGTGCCT GGACCTGTGC GGGCCTGCAC TCCGAAGCC 960
25 TGGCACCCGC GGGCACTCTT GCAATGCCTC CAGAGACCCC CAGAGGGCCC TGCCGTAGGC 1020
CCTTCTGAGG CTCCAGAACA GACCCCGGAG TTGGCAGGAG GCGGAGCCC CGCATACCAG 1080
GGGCCACCTG AGAGTTCTCT CTCCTGA 1107

(7) INFORMATION FOR SEQ ID NO:6:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 368 amino acids

- 8 -

- (B) TYPE: amino acid
 (C) STRANDEDNESS:
 (D) TOPOLOGY: not relevant

(ii) MOLECULE TYPE: protein.

5 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:6:

	Met	Ala	Asn	Ser	Thr	Gly	Leu	Asn	Ala	Ser	Glu	Val	Ala	Gly	Ser	Leu	
	1				5					10					15		
	Gly	Leu	Ile	Leu	Ala	Ala	Val	Val	Glu	Val	Gly	Ala	Leu	Leu	Gly	Asn	
			20						25					30			
10	Gly	Ala	Leu	Leu	Val	Val	Val	Leu	Arg	Thr	Pro	Gly	Leu	Arg	Asp	Ala	
		35						40					45				
	Leu	Tyr	Leu	Ala	His	Leu	Cys	Val	Val	Asp	Leu	Leu	Ala	Ala	Ala	Ser	
		50					55					60					
	Ile	Met	Pro	Leu	Gly	Leu	Leu	Ala	Ala	Pro	Pro	Pro	Gly	Leu	Gly	Arg	
15	65					70				75					80		
	Val	Arg	Leu	Gly	Pro	Ala	Pro	Cys	Arg	Ala	Ala	Arg	Phe	Leu	Ser	Ala	
					85					90					95		
	Ala	Leu	Leu	Pro	Ala	Cys	Thr	Leu	Gly	Val	Ala	Ala	Leu	Gly	Leu	Ala	
			100						105					110			
20	Arg	Tyr	Arg	Leu	Ile	Val	His	Pro	Leu	Arg	Pro	Gly	Ser	Arg	Pro	Pro	
		115						120					125				
	Pro	Val	Leu	Val	Leu	Thr	Ala	Val	Trp	Ala	Ala	Ala	Gly	Leu	Leu	Gly	
		130					135					140					
	Ala	Leu	Ser	Leu	Leu	Gly	Pro	Pro	Pro	Ala	Pro	Pro	Pro	Ala	Pro	Ala	
25	145					150				155					160		
	Arg	Cys	Ser	Val	Leu	Ala	Gly	Gly	Leu	Gly	Pro	Phe	Arg	Pro	Leu	Trp	
				165						170					175		
	Ala	Leu	Leu	Ala	Phe	Ala	Leu	Pro	Ala	Leu	Leu	Leu	Leu	Gly	Ala	Tyr	
			180					185						190			
30	Gly	Gly	Ile	Phe	Val	Val	Ala	Arg	Arg	Ala	Ala	Leu	Arg	Pro	Pro	Arg	
			195					200					205				
	Pro	Ala	Arg	Gly	Ser	Arg	Leu	Arg	Ser	Asp	Ser	Leu	Asp	Ser	Arg	Leu	
		210					215					220					
	Ser	Ile	Leu	Pro	Pro	Leu	Arg	Pro	Arg	Leu	Pro	Gly	Gly	Lys	Ala	Ala	
35	225					230					235				240		
	Leu	Ala	Pro	Ala	Leu	Ala	Val	Gly	Gln	Phe	Ala	Ala	Cys	Trp	Leu	Pro	

- 9 -

	245	250	255
	Tyr Gly Cys Ala Cys Leu Ala Pro Ala Ala Arg Ala Ala Glu Ala Glu		
	260	265	270
5	Ala Ala Val Thr Trp Val Ala Tyr Ser Ala Phe Ala Ala His Pro Phe		
	275	280	285
	Leu Tyr Gly Leu Leu Gln Arg Pro Val Arg Leu Ala Leu Gly Arg Leu		
	290	295	300
	Ser Arg Arg Ala Leu Pro Gly Pro Val Arg Ala Cys Thr Pro Gln Ala		
	305	310	315
10	Trp His Pro Arg Ala Leu Leu Gln Cys Leu Gln Arg Pro Pro Glu Gly		
	325	330	335
	Pro Ala Val Gly Pro Ser Glu Ala Pro Glu Gln Thr Pro Glu Leu Ala		
	340	345	350
15	Gly Gly Arg Ser Pro Ala Tyr Gln Gly Pro Pro Glu Ser Ser Leu Ser		
	355	360	365

(8) INFORMATION FOR SEQ ID NO:7:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 1008 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:7:

	ATGGAATCAT CTTTCTCATT TGGAGTGATC CTTGCTGTCC TGGCCTCCCT CATCATTGCT	60
25	ACTAACACAC TAGTGGCTGT GGCTGTGCTG CTGTTGATCC ACAAGAATGA TGGTGTCACT	120
	CTCTGCTTCA CCTTGAATCT GGCTGTGGCT GACACCTTGA TTGGTGTGGC CATCTCTGGC	180
	CTACTCACAG ACCAGCTCTC CAGCCCTTCT CGGCCACAC AGAAGACCCT GTGCAGCCTG	240
	CGGATGGCAT TTGTCACTTC CTCCGAGCT GCCTCTGTCC TCACGGTCAT GCTGATCACC	300
	TTTGACAGGT ACCTTGCCAT CAAGCAGCCC TTCGCTACT TGAAGATCAT GAGTGGGTTC	360
30	GTGGCCGGGG CCTGCATTGC CGGGCTGTGG TTAGTGTCTT ACCTCATTGG CTTCCTCCCA	420
	CTCGGAATCC CCATGTTCCA GCAGACTGCC TACAAAGGGC AGTGCAGCTT CTTTGCTGTA	480
	TTTACCCCTC ACTTCGTGCT GACCCTCTCC TCGTGTGGCT TCTTCCCAGC CATGCTCCTC	540
	TTTGTCTTCT TCTACTGCGA CATGCTCAAG ATTGCCTCCA TGCACAGCCA GCAGATTCCA	600

- 10 -

AAGATGGAAC ATGCAGGAGC CATGGCTGGA GGTATCGAT CCCACGGAC TCCCAGCGAC 660
 TTCAAAGCTC TCCGTACTGT GTCTGTTCTC ATTGGGAGCT TTGCTCTATC CTGGACCCCC 720
 TTCCTTATCA CTGGCATTGT GCAGGTGGCC TGCCAGGAGT GTCACCTCTA CCTAGTGCTG 780
 GAACGGTACC TGTGGCTGCT CGGCGTGGGC AACTCCCTGC TCAACCCACT CATCTATGCC 840
 5 TATTGGCAGA AGGAGGTGCG ACTGCAGCTC TACCACATGG CCCTAGGAGT GAAGAAGGTG 900
 CTCACCTCAT TCCTCCTCTT TCTCTCGGCC AGGAATTGTG GCCCAGAGAG GCCCAGGGAA 960
 AGTTCCTGTC ACATCGTCAC TATCTCCAGC TCAGAGTTTG ATGGCTAA 1008

(9) INFORMATION FOR SEQ ID NO:8:

(i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 335 amino acids
 (B) TYPE: amino acid
 (C) STRANDEDNESS:
 (D) TOPOLOGY: not relevant

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:8:

Met Glu Ser Ser Phe Ser Phe Gly Val Ile Leu Ala Val Leu Ala Ser
 1 5 10 15
 Leu Ile Ile Ala Thr Asn Thr Leu Val Ala Val Ala Val Leu Leu Leu
 20 25 30
 Ile His Lys Asn Asp Gly Val Ser Leu Cys Phe Thr Leu Asn Leu Ala
 35 40 45
 Val Ala Asp Thr Leu Ile Gly Val Ala Ile Ser Gly Leu Leu Thr Asp
 50 55 60
 Gln Leu Ser Ser Pro Ser Arg Pro Thr Gln Lys Thr Leu Cys Ser Leu
 25 65 70 75 80
 Arg Met Ala Phe Val Thr Ser Ser Ala Ala Ala Ser Val Leu Thr Val
 85 90 95
 Met Leu Ile Thr Phe Asp Arg Tyr Leu Ala Ile Lys Gln Pro Phe Arg
 100 105 110
 Tyr Leu Lys Ile Met Ser Gly Phe Val Ala Gly Ala Cys Ile Ala Gly
 30 115 120 125
 Leu Trp Leu Val Ser Tyr Leu Ile Gly Phe Leu Pro Leu Gly Ile Pro
 130 135 140
 Met Phe Gln Gln Thr Ala Tyr Lys Gly Gln Cys Ser Phe Phe Ala Val

- 11 -

	145		150		155		160									
	Phe	His	Pro	His	Phe	Val	Leu	Thr	Leu	Ser	Cys	Val	Gly	Phe	Phe	Pro
					165					170						175
5	Ala	Met	Leu	Leu	Phe	Val	Phe	Phe	Tyr	Cys	Asp	Met	Leu	Lys	Ile	Ala
				180					185					190		
	Ser	Met	His	Ser	Gln	Gln	Ile	Arg	Lys	Met	Glu	His	Ala	Gly	Ala	Met
			195					200					205			
	Ala	Gly	Gly	Tyr	Arg	Ser	Pro	Arg	Thr	Pro	Ser	Asp	Phe	Lys	Ala	Leu
	210						215					220				
10	Arg	Thr	Val	Ser	Val	Leu	Ile	Gly	Ser	Phe	Ala	Leu	Ser	Trp	Thr	Pro
	225					230					235					240
	Phe	Leu	Ile	Thr	Gly	Ile	Val	Gln	Val	Ala	Cys	Gln	Glu	Cys	His	Leu
					245					250					255	
15	Tyr	Leu	Val	Leu	Glu	Arg	Tyr	Leu	Trp	Leu	Leu	Gly	Val	Gly	Asn	Ser
				260					265					270		
	Leu	Leu	Asn	Pro	Leu	Ile	Tyr	Ala	Tyr	Trp	Gln	Lys	Glu	Val	Arg	Leu
			275					280					285			
	Gln	Leu	Tyr	His	Met	Ala	Leu	Gly	Val	Lys	Lys	Val	Leu	Thr	Ser	Phe
	290						295					300				
20	Leu	Leu	Phe	Leu	Ser	Ala	Arg	Asn	Cys	Gly	Pro	Glu	Arg	Pro	Arg	Glu
	305					310				315						320
	Ser	Ser	Cys	His	Ile	Val	Thr	Ile	Ser	Ser	Ser	Glu	Phe	Asp	Gly	
				325					330					335		

(10) INFORMATION FOR SEQ ID NO:9:

- 25 (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 1413 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

- 30 (ii) MOLECULE TYPE: DNA (genomic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:9:

	ATGGACACTA	CCATGGAAGC	TGACCTGGGT	GCCACTGGCC	ACAGGCCCCG	CACAGAGCTT	60
	GATGATGAGG	ACTCCTACCC	CCAAGGTGGC	TGGGACACGG	TCTTCCTGGT	GGCCCTGCTG	120
	CTCCTTGGGC	TGCCAGCCAA	TGGGTTGATG	GCGTGGCTGG	CCGGCTCCCA	GGCCCGGCAT	180
35	GGAGCTGGCA	CGCGTCTGGC	GCTGCTCCTG	CTCAGCCTGG	CCCTCTCTGA	CTTCTTGTTT	240

- 12 -

CTGGCAGCAG CGGCCTTCCA GATCCTAGAG ATCCGGCATG GGGGACACTG GCCGCTGGGG 300
 ACAGCTGCCT GCCGCTTCTA CTACTTCCTA TGGGGCGTGT CCTACTCCTC CGGCCTCTTC 360
 CTGCTGGCCG CCCTCAGCCT CGACCGCTGC CTGCTGGCGC TGTGCCCACA CTGGTACCCT 420
 GGGCACC GCC CAGTCCGCCT GCCCCTCTGG GTCTGCGCGG GTGTCTGGGT GCTGGCCACA 480
 5 CTCTTCAGCG TGCCCTGGCT GGTCTTCCCC GAGGCTGCCG TCTGGTGGTA CGACCTGGTC 540
 ATCTGCCTGG ACTTCTGGGA CAGCGAGGAG CTGTCGCTGA GGATGCTGGA GGTCTGGGG 600
 GGCTTCCTGC CTTTCCTCCT GCTGCTCGTC TGCCACGTGC TCACCCAGGC CACAGCCTGT 660
 CGCAGCTGCC ACCGCCAACA GCAGCCCGCA GCCTGCCGGG GCTTCGCCCG TGTGGCCAGG 720
 ACCATTCTGT CAGCCTATGT GGTCTGAGG CTGCCCTACC AGCTGGCCCA GCTGCTCTAC 780
 10 CTGGCCTTCC TGTGGGACGT CTACTCTGGC TACCTGCTCT GGGAGGCCCT GGTCTACTCC 840
 GACTACCTGA TCCTACTCAA CAGCTGCCTC AGCCCCTTCC TCTGCCTCAT GGCCAGTGCC 900
 GACCTCCGGA CCCTGCTGCG CTCCGTGCTC TCGTCTTCG CGGCAGCTCT CTGCGAGGAG 960
 CGGCCGGGCA GCTTCACGCC CACTGAGCCA CAGACCCAGC TAGATTCTGA GGGTCCAAC 1020
 CTGCCAGAGC CGATGGCAGA GGCCAGTCA CAGATGGATC CTGTGGCCCA GCCTCAGGTG 1080
 15 AACCCACAC TCCAGCCAG ATCGGATCCC ACAGCTCAGC CACAGCTGAA CCCTACGGCC 1140
 CAGCCACAGT CGGATCCAC AGCCAGCCA CAGCTGAACC TCATGGCCCA GCCACAGTCA 1200
 GATTCTGTGG CCCAGCCACA GGCAGACACT AACGTCCAGA CCCCTGCACC TGCTGCCAGT 1260
 TCTGTGCCCA GTCCCTGTGA TGAAGCTTCC CCAACCCCAT CCTCGCATCC TACCCAGGG 1320
 GCCCTTGAGG ACCCAGCCAC ACCTCCTGCC TCTGAAGGAG AAAGCCCCAG CAGCAGCCCG 1380
 20 CCAGAGGCGG CCCCGGGCGC AGGCCCCACG TGA 1413

(11) INFORMATION FOR SEQ ID NO:10:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 468 amino acids

(B) TYPE: amino acid

(C) STRANDEDNESS:

(D) TOPOLOGY: not relevant

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:10:

Met Asp Thr Thr Met Glu Ala Asp Leu Gly Ala Thr Gly His Arg Pro
 1 5 10 15

- 13 -

Arg Thr Glu Leu Asp Asp Glu Asp Ser Tyr Pro Gln Gly Gly Trp Asp
 20 25 30
 Thr Val Phe Leu Val Ala Leu Leu Leu Leu Gly Leu Pro Ala Asn Gly
 35 40 45
 5 Leu Met Ala Trp Leu Ala Gly Ser Gln Ala Arg His Gly Ala Gly Thr
 50 55 60
 Arg Leu Ala Leu Leu Leu Leu Ser Leu Ala Leu Ser Asp Phe Leu Phe
 65 70 75 80
 10 Leu Ala Ala Ala Ala Phe Gln Ile Leu Glu Ile Arg His Gly Gly His
 85 90 95
 Trp Pro Leu Gly Thr Ala Ala Cys Arg Phe Tyr Tyr Phe Leu Trp Gly
 100 105 110
 Val Ser Tyr Ser Ser Gly Leu Phe Leu Leu Ala Ala Leu Ser Leu Asp
 115 120 125
 15 Arg Cys Leu Leu Ala Leu Cys Pro His Trp Tyr Pro Gly His Arg Pro
 130 135 140
 Val Arg Leu Pro Leu Trp Val Cys Ala Gly Val Trp Val Leu Ala Thr
 145 150 155 160
 20 Leu Phe Ser Val Pro Trp Leu Val Phe Pro Glu Ala Ala Val Trp Trp
 165 170 175
 Tyr Asp Leu Val Ile Cys Leu Asp Phe Trp Asp Ser Glu Glu Leu Ser
 180 185 190
 Leu Arg Met Leu Glu Val Leu Gly Gly Phe Leu Pro Phe Leu Leu Leu
 195 200 205
 25 Leu Val Cys His Val Leu Thr Gln Ala Thr Arg Thr Cys His Arg Gln
 210 215 220
 Gln Gln Pro Ala Ala Cys Arg Gly Phe Ala Arg Val Ala Arg Thr Ile
 225 230 235 240
 30 Leu Ser Ala Tyr Val Val Leu Arg Leu Pro Tyr Gln Leu Ala Gln Leu
 245 250 255
 Leu Tyr Leu Ala Phe Leu Trp Asp Val Tyr Ser Gly Tyr Leu Leu Trp
 260 265 270
 Glu Ala Leu Val Tyr Ser Asp Tyr Leu Ile Leu Leu Asn Ser Cys Leu
 275 280 285
 35 Ser Pro Phe Leu Cys Leu Met Ala Ser Ala Asp Leu Arg Thr Leu Leu
 290 295 300
 Arg Ser Val Leu Ser Ser Phe Ala Ala Ala Leu Cys Glu Glu Arg Pro

- 14 -

305 310 315 320
 Gly Ser Phe Thr Pro Thr Glu Pro Gln Thr Gln Leu Asp Ser Glu Gly
 325 330 335
 Pro Thr Leu Pro Glu Pro Met Ala Glu Ala Gln Ser Gln Met Asp Pro
 5 340 345 350
 Val Ala Gln Pro Gln Val Asn Pro Thr Leu Gln Pro Arg Ser Asp Pro
 355 360 365
 Thr Ala Gln Pro Gln Leu Asn Pro Thr Ala Gln Pro Gln Ser Asp Pro
 370 375 380
 10 Thr Ala Gln Pro Gln Leu Asn Leu Met Ala Gln Pro Gln Ser Asp Ser
 385 390 395 400
 Val Ala Gln Pro Gln Ala Asp Thr Asn Val Gln Thr Pro Ala Pro Ala
 405 410 415
 Ala Ser Ser Val Pro Ser Pro Cys Asp Glu Ala Ser Pro Thr Pro Ser
 15 420 425 430
 Ser His Pro Thr Pro Gly Ala Leu Glu Asp Pro Ala Thr Pro Pro Ala
 435 440 445
 Ser Glu Gly Glu Ser Pro Ser Ser Thr Pro Pro Glu Ala Ala Pro Gly
 450 455 460
 20 Ala Gly Pro Thr
 465

(12) INFORMATION FOR SEQ ID NO:11:

(i) SEQUENCE CHARACTERISTICS:

- 25 (A) LENGTH: 1248 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:11:

30 ATGTCAGGGA TGGAAAACT TCAGAAATGCT TCCTGGATCT ACCAGCAGAA ACTAGAAGAT 60
 CCATTCCAGA AACACCTGAA CAGCACCGAG GAGTATCTGG CCTTCCTCTG CGGACCTCGG 120
 CGCAGCCACT TCTTCCTCCC CGTGTCTGTG GTGTATGTGC CAATTTTTGT GGTGGGGGTC 180
 ATTGGCAATG TCCTGGTGTG CCTGGTGATT CTGCAGCACC AGGCTATGAA GACGCCCACC 240
 AACTACTACC TCTTCAGCCT GCGGGTCTCT GACCTCCTGG TCCTGCTCCT TGAATGCC 300

- 15 -

CTGGAGGTCT ATGAGATGTG GCGCAACTAC CCTTTCTTGT TCGGGCCCGT GGGCTGCTAC 360
 TTCAAGACGG CCCTCTTTGA GACCGTGTGC TTCGCCTCCA TCCTCAGCAT CACCACCGTC 420
 AGCGTGGAGC GCTACGTGGC CATCCTACAC CCGTTCCGCG CCAAAGTGA GAGCACCCGG 480
 CGCCGGGCCC TCAGGATCCT CGGCATCGTC TGGGGCTTCT CCGTGCTCTT CTCCCTGCCC 540
 5 AACACCAGCA TCCATGGCAT CAAGTTCCAC TACTTCCCCA ATGGGTCCCT GGTCACAGGT 600
 TCGGCCACCT GTACGGTCAT CAAGCCCAGT TGGATCTACA ATTTTCATCAT CCAGGTCACC 660
 TCCTTCCTAT TCTACCTCCT CCCCATGACT GTCATCAGTG TCCTCTACTA CCTCATGGCA 720
 CTCAGACTAA AGAAAGACAA ATCTCTTGAG GCAGATGAAG GGAATGCAA TATTCAAAGA 780
 CCCTGCAGAA AATCAGTCAA CAAGATGCTG TTTGTCTTGG TCTTAGTGTT TGCTATCTGT 840
 10 TGGGCCCCGT TCCACATTGA CCGACTCTTC TTCAGCTTGG TGGAGGAGTG GAGTGAATCC 900
 CTGGCTGCTG GTTCAACCT CGTCCATGTG GTGTCAAGTG TCTTCTTCTA CCTGAGCTCA 960
 GCTGTCAACC CCATTATCTA TAACCTACTG TCTCGCCGCT TCCAGGCAGC ATTCCAGAAT 1020
 GTGATCTCTT CTTTCCACAA ACAGTGGCAC TCCCAGCATG ACCCAGAGTT GCCACCTGCC 1080
 CAGCGGAACA TCTTCTGAC AGAATGCCAC TTTGTGGAGC TGACCGAAGA TATAGGTCCC 1140
 15 CAATTCCCAT GTCAGTCATC CATGCACAAC TCTCACCTCC CAACAGCCCT CTCTAGTGAA 1200
 CAGATGTCAA GAACAACTA TCAAAGCTTC CACTTTAACA AAACCTGA 1248

(13) INFORMATION FOR SEQ ID NO:12:

(i) SEQUENCE CHARACTERISTICS:

- 20 (A) LENGTH: 415 amino acids
 (B) TYPE: amino acid
 (C) STRANDEDNESS:
 (D) TOPOLOGY: not relevant

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:12:

25 Met Ser Gly Met Glu Lys Leu Gln Asn Ala Ser Trp Ile Tyr Gln Gln
 1 5 10 15
 Lys Leu Glu Asp Pro Phe Gln Lys His Leu Asn Ser Thr Glu Glu Tyr
 20 25 30
 30 Leu Ala Phe Leu Cys Gly Pro Arg Arg Ser His Phe Phe Leu Pro Val
 35 40 45
 Ser Val Val Tyr Val Pro Ile Phe Val Val Gly Val Ile Gly Asn Val

- 16 -

	50		55		60
	Leu Val Cys Leu Val Ile Leu Gln His Gln Ala Met Lys Thr Pro Thr				
	65		70		75 80
5	Asn Tyr Tyr Leu Phe Ser Leu Ala Val Ser Asp Leu Leu Val Leu Leu				
		85		90	95
	Leu Gly Met Pro Leu Glu Val Tyr Glu Met Trp Arg Asn Tyr Pro Phe				
		100		105	110
	Leu Phe Gly Pro Val Gly Cys Tyr Phe Lys Thr Ala Leu Phe Glu Thr				
		115		120	125
10	Val Cys Phe Ala Ser Ile Leu Ser Ile Thr Thr Val Ser Val Glu Arg				
		130		135	140
	Tyr Val Ala Ile Leu His Pro Phe Arg Ala Lys Leu Gln Ser Thr Arg				
		145		150	155 160
15	Arg Arg Ala Leu Arg Ile Leu Gly Ile Val Trp Gly Phe Ser Val Leu				
		165		170	175
	Phe Ser Leu Pro Asn Thr Ser Ile His Gly Ile Lys Phe His Tyr Phe				
		180		185	190
	Pro Asn Gly Ser Leu Val Pro Gly Ser Ala Thr Cys Thr Val Ile Lys				
		195		200	205
20	Pro Met Trp Ile Tyr Asn Phe Ile Ile Gln Val Thr Ser Phe Leu Phe				
		210		215	220
	Tyr Leu Leu Pro Met Thr Val Ile Ser Val Leu Tyr Tyr Leu Met Ala				
		225		230	235 240
25	Leu Arg Leu Lys Lys Asp Lys Ser Leu Glu Ala Asp Glu Gly Asn Ala				
		245		250	255
	Asn Ile Gln Arg Pro Cys Arg Lys Ser Val Asn Lys Met Leu Phe Val				
		260		265	270
	Leu Val Leu Val Phe Ala Ile Cys Trp Ala Pro Phe His Ile Asp Arg				
		275		280	285
30	Leu Phe Phe Ser Phe Val Glu Glu Trp Ser Glu Ser Leu Ala Ala Val				
		290		295	300
	Phe Asn Leu Val His Val Val Ser Gly Val Phe Phe Tyr Leu Ser Ser				
		305		310	315 320
35	Ala Val Asn Pro Ile Ile Tyr Asn Leu Leu Ser Arg Arg Phe Gln Ala				
		325		330	335
	Ala Phe Gln Asn Val Ile Ser Ser Phe His Lys Gln Trp His Ser Gln				
		340		345	350

- 17 -

His Asp Pro Gln Leu Pro Pro Ala Gln Arg Asn Ile Phe Leu Thr Glu
 355 360 365

Cys His Phe Val Glu Leu Thr Glu Asp Ile Gly Pro Gln Phe Pro Cys
 370 375 380

5 Gln Ser Ser Met His Asn Ser His Leu Pro Thr Ala Leu Ser Ser Glu
 385 390 395 400

Gln Met Ser Arg Thr Asn Tyr Gln Ser Phe His Phe Asn Lys Thr
 405 410 415

(14) INFORMATION FOR SEQ ID NO:13:

- 10 (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 1173 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

- 15 (ii) MOLECULE TYPE: DNA (genomic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:13:

ATGCCAGATA CTAATAGCAC AATCAATTTA TCACTAAGCA CTCGTGTTAC TTTAGCATTT 60
 TTTATGTCCT TAGTAGCTTT TGCTATAATG CTAGGAAATG CTTTGGTCAT TTTAGCTTTT 120
 GTGGTGGACA AAAACCTTAG ACATCGAAGT AGTTATTTTT TTCTTAAC TT GGCCATCTCT 180
 20 GACTTCTTTG TGGGTGTGAT CTCCATTCCT TTGTACATCC CTCACACGCT GTTCGAATGG 240
 GATTTTGGAA AGGAAATCTG TGTATTTTGG CTCACTACTG ACTATCTGTT ATGTACAGCA 300
 TCTGTATATA ACATTGTCCT CATCAGCTAT GATCGATACC TGTCAGTCTC AAATGCTGTG 360
 TCTTATAGAA CTCAACATAC TGGGGTCTTG AAGATTGTTA CTCTGATGGT GGCCGTTTGG 420
 GTGCTGGCCT TCTTAGTGAA TGGGCCAATG ATTCTAGTTT CAGAGTCTTG GAAGGATGAA 480
 25 GGTAGTGAAT GTGAACCTGG ATTTTTTTTCG GAATGGTACA TCCTTGCCAT CACATCATTC 540
 TTGGAATTCG TGATCCCAGT CATCTTAGTC GCTTATTTC AATGAATAT TTATTGGAGC 600
 CTGTGGAAGC GTGATCATCT CAGTAGGTGC CAAAGCCATC CTGGACTGAC TGCTGTCTCT 660
 TCCAACATCT GTGGACACTC ATTCAGAGGT AGACTATCTT CAAGGAGATC TCTTTCTGCA 720
 TCGACAGAAG TTCCTGCATC CTTTCATTCA GAGAGACAGA GGAGAAAGAG TAGTCTCATG 780
 30 TTTTCTCAA GAACCAAGAT GAATAGCAAT ACAATTGCTT CCAAAATGGG TTCCTTCTCC 840
 CAATCAGATT CTGTAGCTCT TCACCAAAGG GAACATGTTG AACTGCTTAG AGCCAGGAGA 900

- 18 -

TTAGCCAAGT CACTGGCCAT TCTCTTAGGG GTTTTGTCTG TTTGCTGGGC TCCATATTCT 960
 CTGTTTACAA TTGTCCTTTC ATTTTATTCC TCAGCAACAG GTCCTAAATC AGTTTGGTAT 1020
 AGAATTGCAT TTTGGCTTCA GTGGTTCAAT TCCTTTGTCA ATCCTCTTTT GTATCCATTG 1080
 TGTCAACAAGC GCTTTCAAAA GGCTTCTTG AAAATATTTT GTATAAAAAA GCAACCTCTA 1140
 5 CCATCACAAC ACAGTCGGTC AGTATCTTCT TAA 1173

(15) INFORMATION FOR SEQ ID NO:14:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 390 amino acids
 (B) TYPE: amino acid
 (C) STRANDEDNESS:
 (D) TOPOLOGY: not relevant

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:14:

Met Pro Asp Thr Asn Ser Thr Ile Asn Leu Ser Leu Ser Thr Arg Val
 1 5 10 15
 Thr Leu Ala Phe Phe Met Ser Leu Val Ala Phe Ala Ile Met Leu Gly
 20 25 30
 Asn Ala Leu Val Ile Leu Ala Phe Val Val Asp Lys Asn Leu Arg His
 35 40 45
 Arg Ser Ser Tyr Phe Phe Leu Asn Leu Ala Ile Ser Asp Phe Phe Val
 50 55 60
 Gly Val Ile Ser Ile Pro Leu Tyr Ile Pro His Thr Leu Phe Glu Trp
 65 70 75 80
 Asp Phe Gly Lys Glu Ile Cys Val Phe Trp Leu Thr Thr Asp Tyr Leu
 85 90 95
 Leu Cys Thr Ala Ser Val Tyr Asn Ile Val Leu Ile Ser Tyr Asp Arg
 100 105 110
 Tyr Leu Ser Val Ser Asn Ala Val Ser Tyr Arg Thr Gln His Thr Gly
 115 120 125
 Val Leu Lys Ile Val Thr Leu Met Val Ala Val Trp Val Leu Ala Phe
 130 135 140
 Leu Val Asn Gly Pro Met Ile Leu Val Ser Glu Ser Trp Lys Asp Glu
 145 150 155 160
 Gly Ser Glu Cys Glu Pro Gly Phe Phe Ser Glu Trp Tyr Ile Leu Ala
 165 170 175

- 19 -

Ile Thr Ser Phe Leu Glu Phe Val Ile Pro Val Ile Leu Val Ala Tyr
 180 185 190
 Phe Asn Met Asn Ile Tyr Trp Ser Leu Trp Lys Arg Asp His Leu Ser
 195 200 205
 5 Arg Cys Gln Ser His Pro Gly Leu Thr Ala Val Ser Ser Asn Ile Cys
 210 215 220
 Gly His Ser Phe Arg Gly Arg Leu Ser Ser Arg Arg Ser Leu Ser Ala
 225 230 235 240
 Ser Thr Glu Val Pro Ala Ser Phe His Ser Glu Arg Gln Arg Arg Lys
 10 245 250 255
 Ser Ser Leu Met Phe Ser Ser Arg Thr Lys Met Asn Ser Asn Thr Ile
 260 265 270
 Ala Ser Lys Met Gly Ser Phe Ser Gln Ser Asp Ser Val Ala Leu His
 275 280 285
 15 Gln Arg Glu His Val Glu Leu Leu Arg Ala Arg Arg Leu Ala Lys Ser
 290 295 300
 Leu Ala Ile Leu Leu Gly Val Phe Ala Val Cys Trp Ala Pro Tyr Ser
 305 310 315 320
 Leu Phe Thr Ile Val Leu Ser Phe Tyr Ser Ser Ala Thr Gly Pro Lys
 20 325 330 335
 Ser Val Trp Tyr Arg Ile Ala Phe Trp Leu Gln Trp Phe Asn Ser Phe
 340 345 350
 Val Asn Pro Leu Leu Tyr Pro Leu Cys His Lys Arg Phe Gln Lys Ala
 355 360 365
 25 Phe Leu Lys Ile Phe Cys Ile Lys Lys Gln Pro Leu Pro Ser Gln His
 370 375 380
 Ser Arg Ser Val Ser Ser
 385 390

(16) INFORMATION FOR SEQ ID NO:15:

- 30 (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 30 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear
 35 (ii) MOLECULE TYPE: DNA (genomic)
 (iv) ANTI-SENSE: NO
 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:15:

- 20 -

GGAAAGCTTA ACGATCCCCA GGAGCAACAT

30

(17) INFORMATION FOR SEQ ID NO:16:

(i) SEQUENCE CHARACTERISTICS:

- 5 (A) LENGTH: 31 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(iv) ANTI-SENSE: YES

10 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:16:

CTGGGATCCT ACGAGAGCAT TTTTCACACA G
31

(18) INFORMATION FOR SEQ ID NO:17:

(i) SEQUENCE CHARACTERISTICS:

- 15 (A) LENGTH: 1128 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

20 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:17:

ATGGCGAACG CGAGCGAGCC GGGTGGCAGC GCGGCGGCG AGGCGGCCGC CCTGGGCCTC 60
AAGCTGGCCA CGCTCAGCCT GCTGCTGTGC GTGAGCCTAG CGGGCAACGT GCTGTTCGCG 120
CTGCTGATCG TCGGGGAGCG CAGCCTGCAC CGCGCCCCGT ACTACCTGCT GCTCGACCTG 180
TGCCTGGCCG ACGGGCTGCG CGCGCTCGCC TGCCTCCCG CCGTCATGCT GCGGCGCGCG 240
25 CGTGCGGCGG CCGCGGCGGG GCGCGCGCCG GCGCGCTGG GCTGCAAGCT GCTCGCCTTC 300
CTGGCCGCGC TCTTCTGCTT CCACGCCGCC TTCCTGCTGC TGGGCGTGGG CGTCACCCGC 360
TACCTGGCCA TCGCGACCA CCGCTTCTAT GCAGAGCGCC TGGCCGGCTG GCCGTGCGCC 420
GCCATGCTGG TGTGCGCCGC CTGGGCGCTG GCGCTGGCCG CGGCCTTCCC GCCAGTGCTG 480
GACGGCGGTG GCGACGACGA GGACGCGCCG TCGGCCCTGG AGCAGCGGCC CGACGGCGCC 540
30 CCCGGCGCGC TGGGCTTCCT GCTGCTGCTG GCCGTGGTGG TGGGCGCCAC GCACCTCGTC 600
TACCTCCGCC TGCTCTTCTT CATCCACGAC CGCCGCAAGA TCGGCGCCGC GCGCCTGGTG 660

- 21 -

CCCGCCGTCA GCCACGACTG GACCTTCCAC GGCCCGGGCG CCACCGGCCA GGCGGCCGCC 720
 AACTGGACGG CGGGCTTCGG CCGCGGGCCC ACGCCGCCCG CGCTTGTGGG CATCCGGCCC 780
 GCAGGGCCGG GCCCGGGCGC GCGCCGCCTC CTCGTGCTGG AAGAATTCAA GACGGAGAAG 840
 AGGCTGTGCA AGATGTTCTA CGCCGTCACG CTGCTCTTCC TGCTCCTCTG GGGGCCCTAC 900
 5 GTCGTGGCCA GCTACCTGCG GGTCTGGTG CGGCCCGGCG CCGTCCCCCA GGCCTACCTG 960
 ACGGCCTCCG TGTGGCTGAC CTTGCGCAG GCCGGCATCA ACCCCGTCGT GTGCTTCCTC 1020
 TTCAACAGGG AGCTGAGGGA CTGCTTCAGG GCCCAGTTCC CCTGCTGCCA GAGCCCCCGG 1080
 ACCACCCAGG CGACCCATCC CTGCGACCTG AAAGGCATTG GTTTATGA 1128

(19) INFORMATION FOR SEQ ID NO:18:

- 10 (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 375 amino acids
 - (B) TYPE: amino acid
 - (C) STRANDEDNESS:
 - (D) TOPOLOGY: not relevant

- 15 (ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:18:

Met Ala Asn Ala Ser Glu Pro Gly Gly Ser Gly Gly Gly Glu Ala Ala
 1 5 10 15
 Ala Leu Gly Leu Lys Leu Ala Thr Leu Ser Leu Leu Leu Cys Val Ser
 20 20 25 30
 Leu Ala Gly Asn Val Leu Phe Ala Leu Leu Ile Val Arg Glu Arg Ser
 35 40 45
 Leu His Arg Ala Pro Tyr Tyr Leu Leu Leu Asp Leu Cys Leu Ala Asp
 50 55 60
 25 Gly Leu Arg Ala Leu Ala Cys Leu Pro Ala Val Met Leu Ala Ala Arg
 65 70 75 80
 Arg Ala Ala Ala Ala Ala Gly Ala Pro Pro Gly Ala Leu Gly Cys Lys
 85 90 95
 30 Leu Leu Ala Phe Leu Ala Ala Leu Phe Cys Phe His Ala Ala Phe Leu
 100 105 110
 Leu Leu Gly Val Gly Val Thr Arg Tyr Leu Ala Ile Ala His His Arg
 115 120 125
 Phe Tyr Ala Glu Arg Leu Ala Gly Trp Pro Cys Ala Ala Met Leu Val
 130 135 140

- 22 -

	Cys	Ala	Ala	Trp	Ala	Leu	Ala	Leu	Ala	Ala	Ala	Phe	Pro	Pro	Val	Leu	
	145					150					155					160	
	Asp	Gly	Gly	Gly	Asp	Asp	Glu	Asp	Ala	Pro	Cys	Ala	Leu	Glu	Gln	Arg	
					165					170					175		
5	Pro	Asp	Gly	Ala	Pro	Gly	Ala	Leu	Gly	Phe	Leu	Leu	Leu	Leu	Ala	Val	
					180				185						190		
	Val	Val	Gly	Ala	Thr	His	Leu	Val	Tyr	Leu	Arg	Leu	Leu	Phe	Phe	Ile	
			195					200						205			
	His	Asp	Arg	Arg	Lys	Met	Arg	Pro	Ala	Arg	Leu	Val	Pro	Ala	Val	Ser	
10							210						220				
	His	Asp	Trp	Thr	Phe	His	Gly	Pro	Gly	Ala	Thr	Gly	Gln	Ala	Ala	Ala	
	225					230					235					240	
	Asn	Trp	Thr	Ala	Gly	Phe	Gly	Arg	Gly	Pro	Thr	Pro	Pro	Ala	Leu	Val	
					245					250					255		
15	Gly	Ile	Arg	Pro	Ala	Gly	Pro	Gly	Arg	Gly	Ala	Arg	Arg	Leu	Leu	Val	
				260					265						270		
	Leu	Glu	Glu	Phe	Lys	Thr	Glu	Lys	Arg	Leu	Cys	Lys	Met	Phe	Tyr	Ala	
				275				280					285				
	Val	Thr	Leu	Leu	Phe	Leu	Leu	Leu	Trp	Gly	Pro	Tyr	Val	Val	Ala	Ser	
20				290				295					300				
	Tyr	Leu	Arg	Val	Leu	Val	Arg	Pro	Gly	Ala	Val	Pro	Gln	Ala	Tyr	Leu	
	305					310					315					320	
	Thr	Ala	Ser	Val	Trp	Leu	Thr	Phe	Ala	Gln	Ala	Gly	Ile	Asn	Pro	Val	
					325					330					335		
25	Val	Cys	Phe	Leu	Phe	Asn	Arg	Glu	Leu	Arg	Asp	Cys	Phe	Arg	Ala	Gln	
				340					345					350			
	Phe	Pro	Cys	Cys	Gln	Ser	Pro	Arg	Thr	Thr	Gln	Ala	Thr	His	Pro	Cys	
				355				360						365			
30	Asp	Leu	Lys	Gly	Ile	Gly	Leu										
	370					375											

(20) INFORMATION FOR SEQ ID NO:19:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 1002 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

- 23 -

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:19:

ATGAACACCA CAGTGATGCA AGGCTTCAAC AGATCTGAGC GGTGCCCCAG AGACACTCGG 60
 ATAGTACAGC TGGTATTCCC AGCCCTCTAC ACAGTGGTTT TCTTGACCGG CATCCTGCTG 120
 AATACTTTGG CTCTGTGGGT GTTGTGTCAC ATCCCCAGCT CCTCCACCTT CATCATCTAC 180
 5 CTCAAAAACA CTTTGGTGGC CGACTTGATA ATGACACTCA TGCTTCCTTT CAAAATCCTC 240
 TCTGACTCAC ACCTGGCACC CTGGCAGCTC AGAGCTTTTG TGTGTCGTTT TTCTTCGGTG 300
 ATATTTTATG AGACCATGTA TGTGGGCATC GTGCTGTTAG GGCTCATAGC CTTTGACAGA 360
 TTCCTCAAGA TCATCAGACC TTTGAGAAAT ATTTTCTAA AAAAACCTGT TTTTGCAAAA 420
 ACGGTCTCAA TCTTCATCTG GTTCTTTTTG TTCTTCATCT CCCTGCCAAA TACGATCTTG 480
 10 AGCAACAAGG AAGCAACACC ATCGTCTGTG AAAAAGTGTG CTTCTTAAA GGGGCCTCTG 540
 GGGCTGAAAT GGCATCAAAT GGTAAATAAC ATATGCCAGT TTATTTTCTG GACTGTTTTT 600
 ATCCTAATGC TTGTGTTTFA TGTGGTTATT GCAAAAAAAG TATATGATTC TTATAGAAAG 660
 TCCAAAAGTA AGGACAGAAA AAACAACAAA AAGCTGGAAG GCAAAGTATT TGTGTGCGTG 720
 GCTGTCTTCT TTGTGTGTTT TGCTCCATT CATTTTGCCA GAGTTCCATA TACTCACAGT 780
 15 CAAACCAACA ATAAGACTGA CTGTAGACTG CAAAATCAAC TGTTTATTGC TAAAGAAACA 840
 ACTCTCTTTT TGGCAGCAAC TAACATTTGT ATGGATCCCT TAATATACAT ATTCTTATGT 900
 AAAAAATTCA CAGAAAAGCT ACCATGTATG CAAGGGAGAA AGACCACAGC ATCAAGCCAA 960
 GAAAATCATA GCAGTCAGAC AGACAACATA ACCTTAGGCT GA 1002

(21) INFORMATION FOR SEQ ID NO:20:

- 20 (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 333 amino acids
 (B) TYPE: amino acid
 (C) STRANDEDNESS:
 (D) TOPOLOGY: not relevant

- 25 (ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:20:

Met Asn Thr Thr Val Met Gln Gly Phe Asn Arg Ser Glu Arg Cys Pro
 1 5 10 15
 Arg Asp Thr Arg Ile Val Gln Leu Val Phe Pro Ala Leu Tyr Thr Val
 20 25 30

- 24 -

Val Phe Leu Thr Gly Ile Leu Leu Asn Thr Leu Ala Leu Trp Val Phe
 35 40 45
 Val His Ile Pro Ser Ser Ser Thr Phe Ile Ile Tyr Leu Lys Asn Thr
 50 55 60
 5 Leu Val Ala Asp Leu Ile Met Thr Leu Met Leu Pro Phe Lys Ile Leu
 65 70 75 80
 Ser Asp Ser His Leu Ala Pro Trp Gln Leu Arg Ala Phe Val Cys Arg
 85 90 95
 10 Phe Ser Ser Val Ile Phe Tyr Glu Thr Met Tyr Val Gly Ile Val Leu
 100 105 110
 Leu Gly Leu Ile Ala Phe Asp Arg Phe Leu Lys Ile Ile Arg Pro Leu
 115 120 125
 Arg Asn Ile Phe Leu Lys Lys Pro Val Phe Ala Lys Thr Val Ser Ile
 130 135 140
 15 Phe Ile Trp Phe Phe Leu Phe Phe Ile Ser Leu Pro Asn Thr Ile Leu
 145 150 155 160
 Ser Asn Lys Glu Ala Thr Pro Ser Ser Val Lys Lys Cys Ala Ser Leu
 165 170 175
 20 Lys Gly Pro Leu Gly Leu Lys Trp His Gln Met Val Asn Asn Ile Cys
 180 185 190
 Gln Phe Ile Phe Trp Thr Val Phe Ile Leu Met Leu Val Phe Tyr Val
 195 200 205
 Val Ile Ala Lys Lys Val Tyr Asp Ser Tyr Arg Lys Ser Lys Ser Lys
 210 215 220
 25 Asp Arg Lys Asn Asn Lys Lys Leu Glu Gly Lys Val Phe Val Val Val
 225 230 235 240
 Ala Val Phe Phe Val Cys Phe Ala Pro Phe His Phe Ala Arg Val Pro
 245 250 255
 30 Tyr Thr His Ser Gln Thr Asn Asn Lys Thr Asp Cys Arg Leu Gln Asn
 260 265 270
 Gln Leu Phe Ile Ala Lys Glu Thr Thr Leu Phe Leu Ala Ala Thr Asn
 275 280 285
 Ile Cys Met Asp Pro Leu Ile Tyr Ile Phe Leu Cys Lys Lys Phe Thr
 290 295 300
 35 Glu Lys Leu Pro Cys Met Gln Gly Arg Lys Thr Thr Ala Ser Ser Gln
 305 310 315 320
 Glu Asn His Ser Ser Gln Thr Asp Asn Ile Thr Leu Gly

- 25 -

325

330

(22) INFORMATION FOR SEQ ID NO:21:

(i) SEQUENCE CHARACTERISTICS:

- 5 (A) LENGTH: 1122 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:21:

```

10 ATGGCCAACA CTACCGGAGA GCCTGAGGAG GTGAGCGGCG CTCTGTCCCC ACCGTCCGCA    60
   TCAGCTTATG TGAAGCTGGT ACTGCTGGGA CTGATTATGT GCGTGAGCCT GCGGGGTAAC    120
   GCCATCTTGT CCCTGCTGGT GCTCAAGGAG CGTGCCCTGC ACAAGGCTCC TTACTACTTC    180
   CTGCTGGACC TGTGCCTGGC CGATGGCATA CGCTCTGCCG TCTGCTTCCC CTTTGTGCTG    240
   GCTTCTGTGC GCCACGGCTC TTCATGGACC TTCAGTGACAC TCAGCTGCAA GATTGTGGCC    300
15 TTTATGGCCG TGCTCTTTTG CTTCCATGCG GCCTTCATGC TGTTCATGCAT CAGCGTCACC    360
   CGCTACATGG CCATCGCCCA CCACCGCTTC TACGCCAAGC GCATGACACT CTGGACATGC    420
   GCGGCTGTCA TCTGCATGGC CTGGACCCTG TCTGTGGCCA TGGCCTTCCC ACCTGTCTTT    480
   GACGTGGGCA CCTACAAGTT TATTCGGGAG GAGGACCAGT GCATCTTTGA GCATCGCTAC    540
   TTCAAGGCCA ATGACACGCT GGGCTTCATG CTTATGTTGG CTGTGCTCAT GGCAGCTACC    600
20 CATGCTGTCT ACGGCAAGCT GTCCTCTTC GAGTATCGTC ACCGCAAGAT GAAGCCAGTG    660
   CAGATGGTGC CAGCCATCAG CCAGAACTGG ACATTCCATG GTCCCGGGGC CACCGGCCAG    720
   GCTGCTGCCA ACTGGATCGC CGGCTTTGGC CGTGGGCCCA TGCCACCAAC CCTGCTGGGT    780
   ATCCGGCAGA ATGGGCATGC AGCCAGCCGG CGGCTACTGG GCATGGACGA GGTCAAGGGT    840
   GAAAAGCAGC TGGGCCGCAT GTTCTACGCG ATCACACTGC TCTTTCTGCT CCTCTGGTCA    900
25 CCCTACATCG TGGCCTGCTA CTGGCGAGTG TTTGTGAAAG CCTGTGCTGT GCCCCACCGC    960
   TACCTGGCCA CTGCTGTTTG GATGAGCTTC GCCCAGGCTG CCGTCAACCC AATTGTCTGC   1020
   TTCTGCTCA ACAAGGACCT CAAGAAGTGC CTGACCACTC ACGCCCCCTG CTGGGGCACA   1080
   GGAGGTGCCC CGGCTCCCAG AGAACCTAC TGTGTCATGT GA                               1122

```

(23) INFORMATION FOR SEQ ID NO:22:

- 26 -

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 373 amino acids
 (B) TYPE: amino acid
 (C) STRANDEDNESS:
 (D) TOPOLOGY: not relevant

5

(ii) MOLECULE TYPE: DNA (genomic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:22:

	Met	Ala	Asn	Thr	Thr	Gly	Glu	Pro	Glu	Glu	Val	Ser	Gly	Ala	Leu	Ser	
	1				5					10					15		
10	Pro	Pro	Ser	Ala	Ser	Ala	Tyr	Val	Lys	Leu	Val	Leu	Leu	Gly	Leu	Ile	
				20					25					30			
	Met	Cys	Val	Ser	Leu	Ala	Gly	Asn	Ala	Ile	Leu	Ser	Leu	Leu	Val	Leu	
				35				40						45			
15	Lys	Glu	Arg	Ala	Leu	His	Lys	Ala	Pro	Tyr	Tyr	Phe	Leu	Leu	Asp	Leu	
				50			55					60					
	Cys	Leu	Ala	Asp	Gly	Ile	Arg	Ser	Ala	Val	Cys	Phe	Pro	Phe	Val	Leu	
				65			70				75					80	
	Ala	Ser	Val	Arg	His	Gly	Ser	Ser	Trp	Thr	Phe	Ser	Ala	Leu	Ser	Cys	
					85					90					95		
20	Lys	Ile	Val	Ala	Phe	Met	Ala	Val	Leu	Phe	Cys	Phe	His	Ala	Ala	Phe	
				100					105					110			
	Met	Leu	Phe	Cys	Ile	Ser	Val	Thr	Arg	Tyr	Met	Ala	Ile	Ala	His	His	
				115				120					125				
25	Arg	Phe	Tyr	Ala	Lys	Arg	Met	Thr	Leu	Trp	Thr	Cys	Ala	Ala	Val	Ile	
				130			135						140				
	Cys	Met	Ala	Trp	Thr	Leu	Ser	Val	Ala	Met	Ala	Phe	Pro	Pro	Val	Phe	
				145			150				155					160	
	Asp	Val	Gly	Thr	Tyr	Lys	Phe	Ile	Arg	Glu	Glu	Asp	Gln	Cys	Ile	Phe	
					165					170					175		
30	Glu	His	Arg	Tyr	Phe	Lys	Ala	Asn	Asp	Thr	Leu	Gly	Phe	Met	Leu	Met	
				180					185					190			
	Leu	Ala	Val	Leu	Met	Ala	Ala	Thr	His	Ala	Val	Tyr	Gly	Lys	Leu	Leu	
				195				200					205				
35	Leu	Phe	Glu	Tyr	Arg	His	Arg	Lys	Met	Lys	Pro	Val	Gln	Met	Val	Pro	
				210			215					220					
	Ala	Ile	Ser	Gln	Asn	Trp	Thr	Phe	His	Gly	Pro	Gly	Ala	Thr	Gly	Gln	
				225			230				235					240	

- 27 -

	Ala Ala Ala Asn Trp Ile Ala Gly Phe Gly Arg Gly Pro Met Pro Pro	245	250	255
	Thr Leu Leu Gly Ile Arg Gln Asn Gly His Ala Ala Ser Arg Arg Leu	260	265	270
5	Leu Gly Met Asp Glu Val Lys Gly Glu Lys Gln Leu Gly Arg Met Phe	275	280	285
	Tyr Ala Ile Thr Leu Leu Phe Leu Leu Leu Trp Ser Pro Tyr Ile Val	290	295	300
10	Ala Cys Tyr Trp Arg Val Phe Val Lys Ala Cys Ala Val Pro His Arg	305	310	315
	Tyr Leu Ala Thr Ala Val Trp Met Ser Phe Ala Gln Ala Ala Val Asn	325	330	335
	Pro Ile Val Cys Phe Leu Leu Asn Lys Asp Leu Lys Lys Cys Leu Thr	340	345	350
15	Thr His Ala Pro Cys Trp Gly Thr Gly Gly Ala Pro Ala Pro Arg Glu	355	360	365
	Pro Tyr Cys Val Met	370		

(24) INFORMATION FOR SEQ ID NO:23:

20 (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 1053 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

25 (ii) MOLECULE TYPE: DNA (genomic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:23:

	ATGGCTTTGG	AACAGAACCA	GTCAACAGAT	TATTATTATG	AGGAAAATGA	AATGAATGGC	60
	ACTTATGACT	ACAGTCAATA	TGAATTGATC	TGTATCAAAG	AAGATGTCAG	AGAATTTGCA	120
	AAAGTTTTC	TCCCTGTATT	CCTCACAATA	GCTTTCGTCA	TTGGACTTGC	AGGCAATTCC	180
30	ATGGTAGTGG	CAATTTATGC	CTATTACAAG	AAACAGAGAA	CCAAAACAGA	TGTGTACATC	240
	CTGAATTTGG	CTGTAGCAGA	TTTACTCCTT	CTATTCACTC	TGCCTTTTTG	GGCTGTTAAT	300
	GCAGTTCATG	GGTGGGTTTT	AGGGAAAATA	ATGTGCAAAA	TAACTTCAGC	CTTGTTACACA	360
	CTAAACTTTG	TCTCTGGAAT	GCAGTTTCTG	GCTTGCATCA	GCATAGACAG	ATATGTGGCA	420
	GTAACATAATG	TCCCCAGCCA	ATCAGGAGTG	GGAAAACCAT	GCTGGATCAT	CTGTTTCTGT	480

- 28 -

GTCTGGATGG CTGCCATCTT GCTGAGCATA CCCCAGCTGG TTTTATATAC AGTAAATGAC 540
 AATGCTAGGT GCATTCCCAT TTTCCCCCGC TACCTAGGAA CATCAATGAA AGCATTGATT 600
 CAAATGCTAG AGATCTGCAT TGGATTTGTA GTACCCTTTC TTATTATGGG GGTGTGCTAC 660
 TTTATCACGG CAAGGACACT CATGAAGATG CCAAACATTA AAATATCTCG ACCCCTAAAA 720
 5 GTTCTGCTCA CAGTCGTTAT AGTTTTCATT GTCACTCAAC TGCCTTATAA CATTGTCAAG 780
 TTCTGCCGAG CCATAGACAT CATCTACTCC CTGATCACCA GCTGCAACAT GAGCAAACGC 840
 ATGGACATCG CCATCCAAGT CACAGAAAGC ATTGCACTCT TTCACAGCTG CCTCAACCCA 900
 ATCCTTTATG TTTTATGGG AGCATCTTTC AAAAATACTG TTATGAAAGT GGCCAAGAAA 960
 TATGGGTCTT GGAGAAGACA GAGACAAAGT GTGGAGGAGT TTCCTTTTGA TTCTGAGGGT 1020
 10 CCTACAGAGC CAACCAGTAC TTTTAGCATT TAA 1053

(25) INFORMATION FOR SEQ ID NO:24:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 350 amino acids
 (B) TYPE: amino acid
 15 (C) STRANDEDNESS:
 (D) TOPOLOGY: not relevant

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:24:

20 Met Ala Leu Glu Gln Asn Gln Ser Thr Asp Tyr Tyr Tyr Glu Glu Asn
 1 5 10 15
 Glu Met Asn Gly Thr Tyr Asp Tyr Ser Gln Tyr Glu Leu Ile Cys Ile
 20 25 30
 Lys Glu Asp Val Arg Glu Phe Ala Lys Val Phe Leu Pro Val Phe Leu
 35 40 45
 25 Thr Ile Ala Phe Val Ile Gly Leu Ala Gly Asn Ser Met Val Val Ala
 50 55 60
 Ile Tyr Ala Tyr Tyr Lys Lys Gln Arg Thr Lys Thr Asp Val Tyr Ile
 65 70 75 80
 30 Leu Asn Leu Ala Val Ala Asp Leu Leu Leu Leu Phe Thr Leu Pro Phe
 85 90 95
 Trp Ala Val Asn Ala Val His Gly Trp Val Leu Gly Lys Ile Met Cys
 100 105 110
 Lys Ile Thr Ser Ala Leu Tyr Thr Leu Asn Phe Val Ser Gly Met Gln

- 29 -

	115	120	125
	Phe Leu Ala Cys Ile Ser Ile Asp Arg Tyr Val Ala Val Thr Asn Val		
	130	135	140
5	Pro Ser Gln Ser Gly Val Gly Lys Pro Cys Trp Ile Ile Cys Phe Cys		
	145	150	155
	Val Trp Met Ala Ala Ile Leu Leu Ser Ile Pro Gln Leu Val Phe Tyr		
		165	170
			175
	Thr Val Asn Asp Asn Ala Arg Cys Ile Pro Ile Phe Pro Arg Tyr Leu		
		180	185
			190
10	Gly Thr Ser Met Lys Ala Leu Ile Gln Met Leu Glu Ile Cys Ile Gly		
		195	200
			205
	Phe Val Val Pro Phe Leu Ile Met Gly Val Cys Tyr Phe Ile Thr Ala		
		210	215
			220
15	Arg Thr Leu Met Lys Met Pro Asn Ile Lys Ile Ser Arg Pro Leu Lys		
		225	230
			235
			240
	Val Leu Leu Thr Val Val Ile Val Phe Ile Val Thr Gln Leu Pro Tyr		
		245	250
			255
	Asn Ile Val Lys Phe Cys Arg Ala Ile Asp Ile Ile Tyr Ser Leu Ile		
		260	265
			270
20	Thr Ser Cys Asn Met Ser Lys Arg Met Asp Ile Ala Ile Gln Val Thr		
		275	280
			285
	Glu Ser Ile Ala Leu Phe His Ser Cys Leu Asn Pro Ile Leu Tyr Val		
		290	295
			300
25	Phe Met Gly Ala Ser Phe Lys Asn Tyr Val Met Lys Val Ala Lys Lys		
		305	310
			315
			320
	Tyr Gly Ser Trp Arg Arg Gln Arg Gln Ser Val Glu Glu Phe Pro Phe		
		325	330
			335
	Asp Ser Glu Gly Pro Thr Glu Pro Thr Ser Thr Phe Ser Ile		
		340	345
			350

30 (26) INFORMATION FOR SEQ ID NO:25:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 1116 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

35

(ii) MOLECULE TYPE: DNA (genomic)

- 30 -

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:25:

ATGCCAGGAA ACGCCACCCC AGTGACCACC ACTGCCCCGT GGGCCTCCCT GGGCCTCTCC 60
 GCCAAGACCT GCAACAACGT GTCCTTCGAA GAGAGCAGGA TAGTCCTGGT CGTGGTGTAC 120
 AGCGCGGTGT GCACGCTGGG GGTGCCGGCC AACTGCCTGA CTGCGTGGCT GGCCTGCTG 180
 5 CAGGTACTGC AGGGCAACGT GCTGGCCGTC TACCTGCTCT GCCTGGCACT CTGCGAACTG 240
 CTGTACACAG GCACGCTGCC ACTCTGGGTC ATCTATATCC GCAACCAGCA CCGCTGGACC 300
 CTAGGCCTGC TGGCCTCGAA GGTGACCGCC TACATCTTCT TCTGCAACAT CTACGTCAGC 360
 ATCCTCTTCC TGTGCTGCAT CTCCTGCGAC CGCTTCGTGG CCGTGGTGTA CGCGCTGGAG 420
 AGTCGGGGCC GCCGCCGCCG GAGGACCGCC ATCTCATCT CCGCCTGCAT CTTTCATCTC 480
 10 GTCGGGATCG TTCACTACCC GGTGTTCCAG ACGGAAGACA AGGAGACCTG CTTTGACATG 540
 CTGCAGATGG ACAGCAGGAT TGCCGGGTAC TACTACGCCA GGTTACCGT TGGCTTTGCC 600
 ATCCCTCTCT CCATCATCGC CTTACCAAC CACCGGATTT TCAGGAGCAT CAAGCAGAGC 660
 ATGGGCTTAA GCGCTGCCCA GAAGGCCAAG GTGAAGCACT CGGCCATCGC GGTGGTTGTC 720
 ATCTTCCTAG TCTGCTTCGC CCCGTACCAC CTGGTTCTCC TCGTCAAAGC CGCTGCCTTT 780
 15 TCCTACTACA GAGGAGACAG GAACGCCATG TCGCGCTTGG AGGAAAGGCT GTACACAGCC 840
 TCTGTGGTGT TTCTGTGCCT GTCCACGGTG AACGGCGTGG CTGACCCCAT TATCTACGTG 900
 CTGGCCACGG ACCATTCCCG CCAAGAAGTG TCCAGAATCC ATAAGGGGTG GAAAGAGTGG 960
 TCCATGAAGA CAGACGTCAC CAGGCTCACC CACAGCAGGG ACACCGAGGA GCTGCAGTCG 1020
 CCCGTGGCCC TTGCAGACCA CTACACCTTC TCCAGGCCCG TGCACCCACC AGGGTCACCA 1080
 20 TGCCCTGCAA AGAGGCTGAT TGAGGAGTCC TGCTGA 1116

(28) INFORMATION FOR SEQ ID NO:26:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 371 amino acids
 (B) TYPE: amino acid
 25 (C) STRANDEDNESS:
 (D) TOPOLOGY: not relevant

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:26:

30 Met Pro Gly Asn Ala Thr Pro Val Thr Thr Thr Ala Pro Trp Ala Ser
 1 5 10 15

- 31 -

	Leu Gly Leu Ser Ala Lys Thr Cys Asn Asn Val Ser Phe Glu Glu Ser	
	20	25 30
	Arg Ile Val Leu Val Val Val Tyr Ser Ala Val Cys Thr Leu Gly Val	
	35	40 45
5	Pro Ala Asn Cys Leu Thr Ala Trp Leu Ala Leu Leu Gln Val Leu Gln	
	50	55 60
	Gly Asn Val Leu Ala Val Tyr Leu Leu Cys Leu Ala Leu Cys Glu Leu	
	65	70 75 80
10	Leu Tyr Thr Gly Thr Leu Pro Leu Trp Val Ile Tyr Ile Arg Asn Gln	
	85	90 95
	His Arg Trp Thr Leu Gly Leu Leu Ala Ser Lys Val Thr Ala Tyr Ile	
	100	105 110
	Phe Phe Cys Asn Ile Tyr Val Ser Ile Leu Phe Leu Cys Cys Ile Ser	
	115	120 125
15	Cys Asp Arg Phe Val Ala Val Val Tyr Ala Leu Glu Ser Arg Gly Arg	
	130	135 140
	Arg Arg Arg Arg Thr Ala Ile Leu Ile Ser Ala Cys Ile Phe Ile Leu	
	145	150 155 160
20	Val Gly Ile Val His Tyr Pro Val Phe Gln Thr Glu Asp Lys Glu Thr	
	165	170 175
	Cys Phe Asp Met Leu Gln Met Asp Ser Arg Ile Ala Gly Tyr Tyr Tyr	
	180	185 190
	Ala Arg Phe Thr Val Gly Phe Ala Ile Pro Leu Ser Ile Ile Ala Phe	
	195	200 205
25	Thr Asn His Arg Ile Phe Arg Ser Ile Lys Gln Ser Met Gly Leu Ser	
	210	215 220
	Ala Ala Gln Lys Ala Lys Val Lys His Ser Ala Ile Ala Val Val Val	
	225	230 235 240
30	Ile Phe Leu Val Cys Phe Ala Pro Tyr His Leu Val Leu Leu Val Lys	
	245	250 255
	Ala Ala Ala Phe Ser Tyr Tyr Arg Gly Asp Arg Asn Ala Met Cys Gly	
	260	265 270
	Leu Glu Glu Arg Leu Tyr Thr Ala Ser Val Val Phe Leu Cys Leu Ser	
	275	280 285
35	Thr Val Asn Gly Val Ala Asp Pro Ile Ile Tyr Val Leu Ala Thr Asp	
	290	295 300

- 32 -

His Ser Arg Gln Glu Val Ser Arg Ile His Lys Gly Trp Lys Glu Trp
 305 310 315 320
 Ser Met Lys Thr Asp Val Thr Arg Leu Thr His Ser Arg Asp Thr Glu
 325 330 335
 5 Glu Leu Gln Ser Pro Val Ala Leu Ala Asp His Tyr Thr Phe Ser Arg
 340 345 350
 Pro Val His Pro Pro Gly Ser Pro Cys Pro Ala Lys Arg Leu Ile Glu
 355 360 365
 Glu Ser Cys
 10 370

(28) INFORMATION FOR SEQ ID NO:27:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 1113 base pairs

(B) TYPE: nucleic acid

15 (C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:27:

ATGGCGAACT ATAGCCATGC AGCTGACAAC ATTTTGCAAA ATCTCTCGCC TCTAACAGCC 60
 20 TTTCTGAAAC TGACTTCCTT GGGTTTCATA ATAGGAGTCA GCGTGGTGGG CAACCTCCTG 120
 ATCTCCATTT TGCTAGTGAA AGATAAGACC TTGCATAGAG CACCTTACTA CTCCTGTGTG 180
 GATCTTTGCT GTTCAGATAT CCTCAGATCT GCAATTTGTT TCCCATTTGT GTTCAACTCT 240
 GTCAAAAATG GCTCTACCTG GACTTATGGG ACTCTGACTT GCAAAGTGAT TGCCTTTCTG 300
 GGGGTTTTGT CCTGTTTCCA CACTGCTTTC ATGCTCTTCT GCATCAGTGT CACCAGATAC 360
 25 TTAGCTATCG CCCATCACCG CTTCTATACA AAGAGGCTGA CCTTTTGGAC GTGTCTGGCT 420
 GTGATCTGTA TGGTGTGGAC TCTGTCTGTG GCCATGGCAT TTCCCCCGGT TTTAGACGTG 480
 GGCACCTACT CATTCATTAG GGAGGAAGAT CAATGCACCT TCCAACACCG CTCCTTCAGG 540
 GCTAATGATT CCTTAGGATT TATGCTGCTT CTTGCTCTCA TCCTCCTAGC CACACAGCTT 600
 GTCTACCTCA AGCTGATATT TTTCGTCCAC GATCGAAGAA AAATGAAGCC AGTCCAGTTT 660
 30 GTAGCAGCAG TCAGCCAGAA CTGGACTTTT CATGGTCCTG GAGCCAGTGG CCAGGCAGCT 720
 GCCAATTGGC TAGCAGGATT TGGAAGGGGT CCCACACCAC CCACCTTGCT GGGCATCAGG 780
 CAAAATGCAA ACACCACAGG CAGAAGAAGG CTATTGGTCT TAGACGAGTT CAAAATGGAG 840

- 33 -

AAAAGAATCA GCAGAATGTT CTATATAATG ACTTTTCTGT TTCTAACCTT GTGGGGCCCC 900
 TACCTGGTGG CCTGTTATTG GAGAGTTTTT GCAAGAGGGC CTGTAGTACC AGGGGGATT 960
 CTAACAGCTG CTGTCTGGAT GAGTTTTGCC CAAGCAGGAA TCAATCCTTT TGTCTGCATT 1020
 TTCTCAAACA GGGAGCTGAG GCGCTGTTTC AGCACAACCC TTCTTTACTG CAGAAAATCC 1080
 5 AGGTTACCAA GGGAACCTTA CTGTGTTATA TGA 1113

(29) INFORMATION FOR SEQ ID NO:28:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 370 amino acids
 (B) TYPE: amino acid
 10 (C) STRANDEDNESS:
 (D) TOPOLOGY: not relevant

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:28:

15 Met Ala Asn Tyr Ser His Ala Ala Asp Asn Ile Leu Gln Asn Leu Ser
 1 5 10 15
 Pro Leu Thr Ala Phe Leu Lys Leu Thr Ser Leu Gly Phe Ile Ile Gly
 20 25 30
 Val Ser Val Val Gly Asn Leu Leu Ile Ser Ile Leu Leu Val Lys Asp
 35 40 45
 20 Lys Thr Leu His Arg Ala Pro Tyr Tyr Phe Leu Leu Asp Leu Cys Cys
 50 55 60
 Ser Asp Ile Leu Arg Ser Ala Ile Cys Phe Pro Phe Val Phe Asn Ser
 65 70 75 80
 25 Val Lys Asn Gly Ser Thr Trp Thr Tyr Gly Thr Leu Thr Cys Lys Val
 85 90 95
 Ile Ala Phe Leu Gly Val Leu Ser Cys Phe His Thr Ala Phe Met Leu
 100 105 110
 Phe Cys Ile Ser Val Thr Arg Tyr Leu Ala Ile Ala His His Arg Phe
 115 120 125
 30 Tyr Thr Lys Arg Leu Thr Phe Trp Thr Cys Leu Ala Val Ile Cys Met
 130 135 140
 Val Trp Thr Leu Ser Val Ala Met Ala Phe Pro Pro Val Leu Asp Val
 145 150 155 160
 Gly Thr Tyr Ser Phe Ile Arg Glu Glu Asp Gln Cys Thr Phe Gln His

- 34 -

	165	170	175
	Arg Ser Phe Arg Ala Asn Asp Ser Leu Gly Phe Met Leu Leu Leu Ala		
	180	185	190
5	Leu Ile Leu Leu Ala Thr Gln Leu Val Tyr Leu Lys Leu Ile Phe Phe		
	195	200	205
	Val His Asp Arg Arg Lys Met Lys Pro Val Gln Phe Val Ala Ala Val		
	210	215	220
	Ser Gln Asn Trp Thr Phe His Gly Pro Gly Ala Ser Gly Gln Ala Ala		
	225	230	235 240
10	Ala Asn Trp Leu Ala Gly Phe Gly Arg Gly Pro Thr Pro Pro Thr Leu		
	245	250	255
	Leu Gly Ile Arg Gln Asn Ala Asn Thr Thr Gly Arg Arg Arg Leu Leu		
	260	265	270
15	Val Leu Asp Glu Phe Lys Met Glu Lys Arg Ile Ser Arg Met Phe Tyr		
	275	280	285
	Ile Met Thr Phe Leu Phe Leu Thr Leu Trp Gly Pro Tyr Leu Val Ala		
	290	295	300
	Cys Tyr Trp Arg Val Phe Ala Arg Gly Pro Val Val Pro Gly Gly Phe		
	305	310	315 320
20	Leu Thr Ala Ala Val Trp Met Ser Phe Ala Gln Ala Gly Ile Asn Pro		
	325	330	335
	Phe Val Cys Ile Phe Ser Asn Arg Glu Leu Arg Arg Cys Phe Ser Thr		
	340	345	350
25	Thr Leu Leu Tyr Cys Arg Lys Ser Arg Leu Pro Arg Glu Pro Tyr Cys		
	355	360	365
	Val Ile		
	370		

(30) INFORMATION FOR SEQ ID NO:29:

- (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 1080 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

- 35 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:29:

ATGCAGGTCC CGAACAGCAC CGGCCCGGAC AACGCGACGC TGCAGATGCT GCGGAACCCG 60

- 35 -

GCGATCGCGG TGGCCCTGCC CGTGGTGTAC TCGCTGGTGG CGGCGGTCAG CATCCCGGGC 120
 AACCTCTTCT CTCTGTGGGT GCTGTGCCGG CGCATGGGGC CCAGATCCCC GTCGGTCATC 180
 TTCATGATCA ACCTGAGCGT CACGGACCTG ATGCTGGCCA GCGTGTGGCC TTTCCAAATC 240
 TACTACCATT GCAACCGCCA CCACTGGGTA TTCGGGGTGC TGCTTTGCAA CGTGGTGACC 300
 5 GTGGCCTTTT ACGCAAACAT GTATTCCAGC ATCCTCACCA TGACCTGTAT CAGCGTGGAG 360
 CGCTTCCTGG GGGTCTGTGA CCCGCTCAGC TCCAAGCGCT GCGCCCGCCG TCGTTACGCG 420
 GTGGCCGCGT GTGCAGGGAC CTGGCTGCTG CTCTGACCG CCCTGTGCCC GCTGGCGCGC 480
 ACCGATCTCA CCTACCCGGT GCACGCCCTG GGCATCATCA CCTGCTTCGA CGTCCTCAAG 540
 TGGACGATGC TCCCCAGCGT GGCCATGTGG GCCGTGTTCC TCTTCACCAT CTTCATCCTG 600
 10 CTGTTCTCTA TCCCGTTTCG GATCACCGTG GCTTGTTACA CGGCCACCAT CCTCAAGCTG 660
 TTGCGCACGG AGGAGGCGCA CGGCCGGGAG CAGCGGAGGC GCGCGGTGGG CCTGGCCGCG 720
 GTGGTCTTGC TGGCCTTTGT CACCTGCTTC GCCCCAACA ACTTCGTGCT CCTGGCGCAC 780
 ATCGTGAGCC GCCTGTTCTA CGGCAAGAGC TACTACCACG TGTACAAGCT CACGCTGTGT 840
 CTCAGCTGCC TCAACAAC TGCTGGACCCG TTTGTTTATT ACTTTGCGTC CCGGGAATTC 900
 15 CAGCTGCGCC TGCGGGAATA TTTGGGCTGC CGCCGGGTGC CCAGAGACAC CCTGGACACG 960
 CGCCGCGAGA GCCTCTTCTC CGCCAGGACC ACGTCCGTGC GCTCCGAGGC CGGTGCGCAC 1020
 CCTGAAGGGA TGGAGGGAGC CACCAGGCCC GGCTCCAGA GGCAGGAGAG TGTGTTCTGA 1080

(31) INFORMATION FOR SEQ ID NO:30:

- 20 (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 359 amino acids
 (B) TYPE: amino acid
 (C) STRANDEDNESS:
 (D) TOPOLOGY: not relevant

(ii) MOLECULE TYPE: protein

- 25 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:30:

Met Gln Val Pro Asn Ser Thr Gly Pro Asp Asn Ala Thr Leu Gln Met
 1 5 10 15

Leu Arg Asn Pro Ala Ile Ala Val Ala Leu Pro Val Val Tyr Ser Leu
 20 25 30

30 Val Ala Ala Val Ser Ile Pro Gly Asn Leu Phe Ser Leu Trp Val Leu

- 36 -

	35	40	45
	Cys Arg Arg Met Gly Pro Arg Ser Pro Ser Val Ile Phe Met Ile Asn 50 55 60		
5	Leu Ser Val Thr Asp Leu Met Leu Ala Ser Val Leu Pro Phe Gln Ile 65 70 75 80		
	Tyr Tyr His Cys Asn Arg His His Trp Val Phe Gly Val Leu Leu Cys 85 90 95		
	Asn Val Val Thr Val Ala Phe Tyr Ala Asn Met Tyr Ser Ser Ile Leu 100 105 110		
10	Thr Met Thr Cys Ile Ser Val Glu Arg Phe Leu Gly Val Leu Tyr Pro 115 120 125		
	Leu Ser Ser Lys Arg Trp Arg Arg Arg Arg Tyr Ala Val Ala Ala Cys 130 135 140		
15	Ala Gly Thr Trp Leu Leu Leu Thr Ala Leu Cys Pro Leu Ala Arg 145 150 155 160		
	Thr Asp Leu Thr Tyr Pro Val His Ala Leu Gly Ile Ile Thr Cys Phe 165 170 175		
	Asp Val Leu Lys Trp Thr Met Leu Pro Ser Val Ala Met Trp Ala Val 180 185 190		
20	Phe Leu Phe Thr Ile Phe Ile Leu Leu Phe Leu Ile Pro Phe Val Ile 195 200 205		
	Thr Val Ala Cys Tyr Thr Ala Thr Ile Leu Lys Leu Leu Arg Thr Glu 210 215 220		
25	Glu Ala His Gly Arg Glu Gln Arg Arg Arg Ala Val Gly Leu Ala Ala 225 230 235 240		
	Val Val Leu Leu Ala Phe Val Thr Cys Phe Ala Pro Asn Asn Phe Val 245 250 255		
	Leu Leu Ala His Ile Val Ser Arg Leu Phe Tyr Gly Lys Ser Tyr Tyr 260 265 270		
30	His Val Tyr Lys Leu Thr Leu Cys Leu Ser Cys Leu Asn Asn Cys Leu 275 280 285		
	Asp Pro Phe Val Tyr Tyr Phe Ala Ser Arg Glu Phe Gln Leu Arg Leu 290 295 300		
35	Arg Glu Tyr Leu Gly Cys Arg Arg Val Pro Arg Asp Thr Leu Asp Thr 305 310 315 320		
	Arg Arg Glu Ser Leu Phe Ser Ala Arg Thr Thr Ser Val Arg Ser Glu 325 330 335		

- 37 -

Ala Gly Ala His Pro Glu Gly Met Glu Gly Ala Thr Arg Pro Gly Leu
 340 345 350

Gln Arg Gln Glu Ser Val Phe
 355

5 (32) INFORMATION FOR SEQ ID NO:31:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 1503 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 10 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:31:

ATGGAGCGTC CCTGGGAGGA CAGCCCAGGC CCGGAGGGGG CAGCTGAGGG CTCGCCTGTG 60
 CCAGTCGCCG CCGGGGCGCG CTCCGGTGCC GCGGCGAGTG GCACAGGCTG GCAGCCATGG 120
 15 GCTGAGTGCC CGGGACCCAA GGGGAGGGGG CAACTGCTGG CGACCGCCGG CCCTTTGCGT 180
 CGCTGGCCCCG CCCCCTCGCC TGCCAGCTCC AGCCCCGCCC CCGGAGCGGC GTCCGCTCAC 240
 TCGGTTCAAG GCAGCGCGAC TGCGGGTGGC GCACGACCAG GGCGCAGACC TTGGGGCGCG 300
 CGGCCCATGG AGTCGGGGCT GTCGCGGCCG GCGCCGGTGA GCGAGGTCAT CGTCCTGCAT 360
 TACAACTACA CCGGCAAGCT CCGCGGTGCG AGCTACCAGC CGGGTGCCGG CCTGCGCGCC 420
 20 GACGCCGTGG TGTGCCTGGC GGTGTGCGCC TTCATCGTGC TAGAGAATCT AGCCGTGTTG 480
 TTGGTGCTCG GACGCCACCC GCGCTTCCAC GCTCCCATGT TCCTGCTCCT GGGCAGCCTC 540
 ACGTTGTGCG ATCTGCTGGC AGGCGCCGCC TACGCCGCCA ACATCCTACT GTCGGGGCCG 600
 CTCACGCTGA AACTGTCCCC CGCGCTCTGG TTCGCACGGG AGGGAGGCGT CTTCTGGCA 660
 CTCCTGCGT CCGTGCTGAG CTTCTGGCC ATCGCGCTGG AGCGCAGCCT CACCATGGCG 720
 25 CGCAGGGGGC CCGCGCCCGT CTCCAGTCGG GGGCGCACGC TGGCGATGGC AGCCGCGGCC 780
 TGGGGCGTGT CGCTGCTCCT CGGGCTCCTG CCAGCGCTGG GCTGGAATTG CCTGGGTGCG 840
 CTGGACGCTT GCTCCACTGT CTTGCCGCTC TACGCCAAGG CCTACGTGCT CTTCTGCGTG 900
 CTCGCCTTCG TGGGCATCCT GGCCGCGATC TGTGCACTCT ACGCGCGCAT CTA CTGCGCAG 960
 GTACGCGCCA ACGCGCGGCG CTTGCCGGCA CGGCCCGGGA CTGCGGGGAC CACCTCGACC 1020
 30 CGGGCGCGTC GCAAGCCGCG CTCTCTGGCC TTGCTGCGCA CGCTCAGCGT GGTGCTCCTG 1080

- 38 -

GCCTTTGTGG CATGTTGGGG CCCCTCTTC CTGCTGCTGT TGCTCGACGT GGC GTGCCCG 1140
 GCGCGCACCT GTCCTGTACT CCTGCAGGCC GATCCCTTCC TGGGACTGGC CATGGCCAAC 1200
 TCACTTCTGA ACCCCATCAT CTACAGCTC ACCAACCGCG ACCTGCGCCA CGCGTCTCTG 1260
 CGCCTGGTCT GCTGCGGACG CCACTCCTGC GGCAGAGACC CGAGTGGCTC CCAGCAGTCG 1320
 5 GCGAGCGCGG CTGAGGCTTC CGGGGGCCTG CGCCGCTGCC TGCCCCCGGG CCTTGATGGG 1380
 AGCTTCAGCG GCTCGGAGCG CTCATCGCCC CAGCGCGACG GGCTGGACAC CAGCGGCTCC 1440
 ACAGGCAGCC CCGGTGCACC CACAGCCGCC CGGACTCTGG TATCAGAACC GGCTGCAGAC 1500
 TGA 1503

(33) INFORMATION FOR SEQ ID NO:32:

- 10 (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 500 amino acids
 (B) TYPE: amino acid
 (C) STRANDEDNESS:
 (D) TOPOLOGY: not relevant

- 15 (ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:32:

Met Glu Arg Pro Trp Glu Asp Ser Pro Gly Pro Glu Gly Ala Ala Glu
 1 5 10 15
 Gly Ser Pro Val Pro Val Ala Ala Gly Ala Arg Ser Gly Ala Ala Ala
 20 20 25 30
 Ser Gly Thr Gly Trp Gln Pro Trp Ala Glu Cys Pro Gly Pro Lys Gly
 35 40 45
 Arg Gly Gln Leu Leu Ala Thr Ala Gly Pro Leu Arg Arg Trp Pro Ala
 50 55 60
 25 Pro Ser Pro Ala Ser Ser Ser Pro Ala Pro Gly Ala Ala Ser Ala His
 65 70 75 80
 Ser Val Gln Gly Ser Ala Thr Ala Gly Gly Ala Arg Pro Gly Arg Arg
 85 90 95
 30 Pro Trp Gly Ala Arg Pro Met Glu Ser Gly Leu Leu Arg Pro Ala Pro
 100 105 110
 Val Ser Glu Val Ile Val Leu His Tyr Asn Tyr Thr Gly Lys Leu Arg
 115 120 125
 Gly Ala Ser Tyr Gln Pro Gly Ala Gly Leu Arg Ala Asp Ala Val Val
 130 135 140

- 39 -

	Cys	Leu	Ala	Val	Cys	Ala	Phe	Ile	Val	Leu	Glu	Asn	Leu	Ala	Val	Leu	145	150	155	160
	Leu	Val	Leu	Gly	Arg	His	Pro	Arg	Phe	His	Ala	Pro	Met	Phe	Leu	Leu	165	170	175	
5	Leu	Gly	Ser	Leu	Thr	Leu	Ser	Asp	Leu	Leu	Ala	Gly	Ala	Ala	Tyr	Ala	180	185	190	
	Ala	Asn	Ile	Leu	Leu	Ser	Gly	Pro	Leu	Thr	Leu	Lys	Leu	Ser	Pro	Ala	195	200	205	
10	Leu	Trp	Phe	Ala	Arg	Glu	Gly	Gly	Val	Phe	Val	Ala	Leu	Thr	Ala	Ser	210	215	220	
	Val	Leu	Ser	Leu	Leu	Ala	Ile	Ala	Leu	Glu	Arg	Ser	Leu	Thr	Met	Ala	225	230	235	240
	Arg	Arg	Gly	Pro	Ala	Pro	Val	Ser	Ser	Arg	Gly	Arg	Thr	Leu	Ala	Met	245	250	255	
15	Ala	Ala	Ala	Ala	Trp	Gly	Val	Ser	Leu	Leu	Leu	Gly	Leu	Leu	Pro	Ala	260	265	270	
	Leu	Gly	Trp	Asn	Cys	Leu	Gly	Arg	Leu	Asp	Ala	Cys	Ser	Thr	Val	Leu	275	280	285	
20	Pro	Leu	Tyr	Ala	Lys	Ala	Tyr	Val	Leu	Phe	Cys	Val	Leu	Ala	Phe	Val	290	295	300	
	Gly	Ile	Leu	Ala	Ala	Ile	Cys	Ala	Leu	Tyr	Ala	Arg	Ile	Tyr	Cys	Gln	305	310	315	320
	Val	Arg	Ala	Asn	Ala	Arg	Arg	Leu	Pro	Ala	Arg	Pro	Gly	Thr	Ala	Gly	325	330	335	
25	Thr	Thr	Ser	Thr	Arg	Ala	Arg	Arg	Lys	Pro	Arg	Ser	Leu	Ala	Leu	Leu	340	345	350	
	Arg	Thr	Leu	Ser	Val	Val	Leu	Leu	Ala	Phe	Val	Ala	Cys	Trp	Gly	Pro	355	360	365	
30	Leu	Phe	Leu	Leu	Leu	Leu	Leu	Asp	Val	Ala	Cys	Pro	Ala	Arg	Thr	Cys	370	375	380	
	Pro	Val	Leu	Leu	Gln	Ala	Asp	Pro	Phe	Leu	Gly	Leu	Ala	Met	Ala	Asn	385	390	395	400
	Ser	Leu	Leu	Asn	Pro	Ile	Ile	Tyr	Thr	Leu	Thr	Asn	Arg	Asp	Leu	Arg	405	410	415	
35	His	Ala	Leu	Leu	Arg	Leu	Val	Cys	Cys	Gly	Arg	His	Ser	Cys	Gly	Arg	420	425	430	
	Asp	Pro	Ser	Gly	Ser	Gln	Gln	Ser	Ala	Ser	Ala	Ala	Glu	Ala	Ser	Gly				

435

440

445

Gly Leu Arg Arg Cys Leu Pro Pro Gly Leu Asp Gly Ser Phe Ser Gly

450

455.

460

Ser Glu Arg Ser Ser Pro Gln Arg Asp Gly Leu Asp Thr Ser Gly Ser

465

470

475

480

Thr Gly Ser Pro Gly Ala Pro Thr Ala Ala Arg Thr Leu Val Ser Glu

485

490

495

Pro Ala Ala Asp

500

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 1029 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:33:

ATGCAAGCCG TCGACAATCT CACCTCTGCG CCTGGGAACA CCAGTCTGTG CACCAGAGAC 60

TACAAAATCA CCCAGGTCCT CTTCCCACTG CTCTACACTG TCCTGTTTTT TGTTCGACTT 120

20 ATCACAAATG GCCTGGCGAT GAGGATTTTC TTTCAAATCC GGAGTAAATC AAACCTTTATT 180

ATTTTCTTA AGAACACAGT CATTTCTGAT CTTCTCATGA TTCTGACTTT TCCATTCAAA 240

ATTCTTAGTG ATGCCAACT GGAACAGGA CCACTGAGAA CTTTGTGTG TCAAGTTACC 300

TCCGTCATAT TTTATTTTAC AATGTATATC AGTATTTTCAT TCCTGGGACT GATAACTATC 360

GATCGCTACC AGAAGACCAC CAGGCCATTT AAAACATCCA ACCCCAAAAA TCTCTTGGGG 420

25 GCTAAGATTC TCTCTGTTGT CATCTGGGCA TTCATGTTCT TACTCTCTTT GCCTAACATG 480

ATTCTGACCA ACAGGCAGCC GAGAGACAAG AATGTGAAGA AATGCTCTTT CCTTAAATCA 540

GAGTTCGGTC TAGTCTGGCA TGAAATAGTA AATTACATCT GTCAAGTCAT TTTCTGGATT 600

AATTTCTTAA^{*} TTGTTATTGT ATGTTATACA CTCATTACAA AAGAACTGTA CCGGTCATAC 660

GTAAGAACGA GGGGTGTAGG TAAAGTCCCC AGGAAAAAGG TGAACGTCAA AGTTTTTCATT 720

30 ATCATTGCTG TATTCTTTAT TTGTTTTGTT CCTTTCATT TTGCCCGAAT TCCTTACACC 780

CTGAGCCAAA CCCGGGATGT CTTTGACTGC ACTGCTGAAA ATACTCTGTT CTATGTGAAA 840

- 41 -

GAGAGCACTC TGTGGTTAAC TTCCTTAAAT GCATGCCTGG ATCCGTTTCAT CTATTTTTTC 900
 CTTTGCAAGT CCTTCAGAAA TTCCTTGATA AGTATGCTGA AGTGCCCCAA TTCTGCAACA 960
 TCTCTGTCCC AGGACAATAG GAAAAAAGAA CAGGATGGTG GTGACCCAAA TGAAGAGACT 1020
 CCAATGTAA 1029

5 (35) INFORMATION FOR SEQ ID NO:34:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 342 amino acids
 (B) TYPE: amino acid
 (C) STRANDEDNESS:
 (D) TOPOLOGY: not relevant

10

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:34:

Met Gln Ala Val Asp Asn Leu Thr Ser Ala Pro Gly Asn Thr Ser Leu
 1 5 10 15
 15 Cys Thr Arg Asp Tyr Lys Ile Thr Gln Val Leu Phe Pro Leu Leu Tyr
 20 25 30
 Thr Val Leu Phe Phe Val Gly Leu Ile Thr Asn Gly Leu Ala Met Arg
 35 40 45
 20 Ile Phe Phe Gln Ile Arg Ser Lys Ser Asn Phe Ile Ile Phe Leu Lys
 50 55 60
 Asn Thr Val Ile Ser Asp Leu Leu Met Ile Leu Thr Phe Pro Phe Lys
 65 70 75 80
 Ile Leu Ser Asp Ala Lys Leu Gly Thr Gly Pro Leu Arg Thr Phe Val
 85 90 95
 25 Cys Gln Val Thr Ser Val Ile Phe Tyr Phe Thr Met Tyr Ile Ser Ile
 100 105 110
 Ser Phe Leu Gly Leu Ile Thr Ile Asp Arg Tyr Gln Lys Thr Thr Arg
 115 120 125
 30 Pro Phe Lys Thr Ser Asn Pro Lys Asn Leu Leu Gly Ala Lys Ile Leu
 130 135 140
 Ser Val Val Ile Trp Ala Phe Met Phe Leu Leu Ser Leu Pro Asn Met
 145 150 155 160
 Ile Leu Thr Asn Arg Gln Pro Arg Asp Lys Asn Val Lys Lys Cys Ser
 165 170 175
 35 Phe Leu Lys Ser Glu Phe Gly Leu Val Trp His Glu Ile Val Asn Tyr

- 42 -

	180	185	190
	Ile Cys Gln Val Ile Phe Trp	Ile Asn Phe Leu Ile Val	Ile Val Cys
	195	200	205
5	Tyr Thr Leu Ile Thr Lys Glu Leu Tyr Arg Ser Tyr Val Arg Thr Arg		
	210	215	220
	Gly Val Gly Lys Val Pro Arg Lys Lys Val Asn Val Lys Val Phe Ile		
	225	230	235 240
	Ile Ile Ala Val Phe Phe Ile Cys Phe Val Pro Phe His Phe Ala Arg		
	245	250	255
10	Ile Pro Tyr Thr Leu Ser Gln Thr Arg Asp Val Phe Asp Cys Thr Ala		
	260	265	270
	Glu Asn Thr Leu Phe Tyr Val Lys Glu Ser Thr Leu Trp Leu Thr Ser		
	275	280	285
15	Leu Asn Ala Cys Leu Asp Pro Phe Ile Tyr Phe Phe Leu Cys Lys Ser		
	290	295	300
	Phe Arg Asn Ser Leu Ile Ser Met Leu Lys Cys Pro Asn Ser Ala Thr		
	305	310	315 320
	Ser Leu Ser Gln Asp Asn Arg Lys Lys Glu Gln Asp Gly Gly Asp Pro		
	325	330	335
20	Asn Glu Glu Thr Pro Met		
	340		

(36) INFORMATION FOR SEQ ID NO:35:

(i) SEQUENCE CHARACTERISTICS:

- 25 (A) LENGTH: 1077 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:35:

30	ATGTCGGTCT GCTACCGTCC CCCAGGGAAC GAGACACTGC TGAGCTGGAA GACTTCGCGG	60
	GCCACAGGCA CAGCCTTCCT GCTGCTGGCG GCGCTGCTGG GGCTGCCTGG CAACGGCTTC	120
	GTGGTGTGGA GCTTGGCGGG CTGGCGGCCT GCACGGGGGC GACCGCTGGC GGCCACGCTT	180
	GTGCTGCACC TGGCGCTGGC CGACGGCGCG GTGCTGCTGC TCACGCCGCT CTTTGTGGCC	240
	TTCTTGACCC GGCAGGCCTG GCCGCTGGGC CAGGCGGGCT GCAAGGCGGT GTACTACGTG	300

- 43 -

TGC GCGCTCA GCATGTACGC CAGCGTGCTG CTCACCGGCC TGCTCAGCCT GCAGCGCTGC 360
 CTCGCACTCA CCCGCCCCCTT CCTGGCGCCT CGGCTGCGCA GCCCGGCCCT GGCCCGCCGC 420
 CTGCTGCTGG CGGTCTGGCT GGCCGCCCTG TTGCTCGCCG TCCCGGCCGC CGTCTACCGC 480
 CACCTGTGGA GGGACCGCGT ATGCCAGCTG TGCCACCCGT CGCCGGTCCA CGCCGCCGCC 540
 5 CACCTGAGCC TGGAGACTCT GACCGCTTTC GTGCTTCCTT TCGGGCTGAT GCTCGGCTGC 600
 TACAGCGTGA CGCTGGCACC GCTGCGGGGC GCCCGCTGGG GCTCCGGGCG GCACGGGGCG 660
 CGGGTGGGCC GGCTGGTGAG CGCCATCGTG CTGCTCTCG GCTTGCTCTG GGCCCCCTAC 720
 CACGCACTCA ACCTTCTGCA GGCGGTCGCA GCGCTGGCTC CACCGGAAGG GGCCTTGGCG 780
 AAGCTGGGCG GAGCCGGCCA GGCGGCGCGA GCGGGAATA CGGCCTTGGC CTTCTTCAGT 840
 10 TCTAGCGTCA ACCCGGTGCT CTACGTCTTC ACCGCTGGAG ATCTGCTGCC CCGGGCAGGT 900
 CCCCGTTTCC TCACCGGCT CTTCGAAGGC TCTGGGGAGG CCCGAGGGGG CGGCCGCTCT 960
 AGGGAAGGGA CCATGGAGCT CCGAACTACC CCTCAGCTGA AAGTGGTGGG GCAGGGCCGC 1020
 GGCAATGGAG ACCCGGGGGG TGGGATGGAG AAGGACGGTC CGGAATGGGA CCTTTGA 1077

(37) INFORMATION FOR SEQ ID NO:36:

- 15 (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 358 amino acids
 (B) TYPE: amino acid
 (C) STRANDEDNESS:
 (D) TOPOLOGY: not relevant

20 (ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:36:

Met Ser Val Cys Tyr Arg Pro Pro Gly Asn Glu Thr Leu Leu Ser Trp
 1 5 10 15
 Lys Thr Ser Arg Ala Thr Gly Thr Ala Phe Leu Leu Leu Ala Ala Leu
 25 20 25 30
 Leu Gly Leu Pro Gly Asn Gly Phe Val Val Trp Ser Leu Ala Gly Trp
 35 40 45
 Arg Pro Ala Arg Gly Arg Pro Leu Ala Ala Thr Leu Val Leu His Leu
 50 55 60
 30 Ala Leu Ala Asp Gly Ala Val Leu Leu Leu Thr Pro Leu Phe Val Ala
 65 70 75 80
 Phe Leu Thr Arg Gln Ala Trp Pro Leu Gly Gln Ala Gly Cys Lys Ala

- 44 -

	85	90	95
	Val Tyr Tyr Val Cys Ala Leu Ser Met Tyr Ala Ser Val Leu Leu Thr		
	100	105	110
5	Gly Leu Leu Ser Leu Gln Arg Cys Leu Ala Val Thr Arg Pro Phe Leu		
	115	120	125
	Ala Pro Arg Leu Arg Ser Pro Ala Leu Ala Arg Arg Leu Leu Leu Ala		
	130	135	140
	Val Trp Leu Ala Ala Leu Leu Leu Ala Val Pro Ala Ala Val Tyr Arg		
	145	150	155
10	His Leu Trp Arg Asp Arg Val Cys Gln Leu Cys His Pro Ser Pro Val		
	165	170	175
	His Ala Ala Ala His Leu Ser Leu Glu Thr Leu Thr Ala Phe Val Leu		
	180	185	190
15	Pro Phe Gly Leu Met Leu Gly Cys Tyr Ser Val Thr Leu Ala Arg Leu		
	195	200	205
	Arg Gly Ala Arg Trp Gly Ser Gly Arg His Gly Ala Arg Val Gly Arg		
	210	215	220
	Leu Val Ser Ala Ile Val Leu Ala Phe Gly Leu Leu Trp Ala Pro Tyr		
	225	230	235
20	His Ala Val Asn Leu Leu Gln Ala Val Ala Ala Leu Ala Pro Pro Glu		
	245	250	255
	Gly Ala Leu Ala Lys Leu Gly Gly Ala Gly Gln Ala Ala Arg Ala Gly		
	260	265	270
25	Thr Thr Ala Leu Ala Phe Phe Ser Ser Ser Val Asn Pro Val Leu Tyr		
	275	280	285
	Val Phe Thr Ala Gly Asp Leu Leu Pro Arg Ala Gly Pro Arg Phe Leu		
	290	295	300
	Thr Arg Leu Phe Glu Gly Ser Gly Glu Ala Arg Gly Gly Gly Arg Ser		
	305	310	315
30	Arg Glu Gly Thr Met Glu Leu Arg Thr Thr Pro Gln Leu Lys Val Val		
	325	330	335
	Gly Gln Gly Arg Gly Asn Gly Asp Pro Gly Gly Gly Met Glu Lys Asp		
	340	345	350
35	Gly Pro Glu Trp Asp Leu		
	355		

(38) INFORMATION FOR SEQ ID NO:37:

- 45 -

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 1005 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:37:

ATGCTGGGGA TCATGGCATG GAATGCAACT TGCAAAACT GGCTGGCAGC AGAGGCTGCC 60
 CTGGAAAAGT ACTACCTTTC CATTTTTTAT GGGATTGAGT TCGTTGTGGG AGTCCTTGGA 120
 10 AATACCATTG TTGTTTACGG CTACATCTTC TCTCTGAAGA ACTGGAACAG CAGTAATATT 180
 TATCTCTTTA ACCTCTCTGT CTCTGACTTA GCTTTTCTGT GCACCCTCCC CATGCTGATA 240
 AGGAGTTATG CCAATGGAAA CTGGATATAT GGAGACGTGC TCTGCATAAG CAACCGATAT 300
 GTGCTTCATG CCAACCTCTA TACCAGCATT CTCTTCTCA CTTTATCAG CATAGATCGA 360
 TACTTGATAA TTAAGTATCC TTTCCGAGAA CACCTTCTGC AAAAGAAAGA GTTTGCTATT 420
 15 TTAATCTCCT TGGCCATTG GGTTTTAGTA ACCTTAGAGT TACTACCCAT ACTTCCCCTT 480
 ATAAATCCTG TTATAACTGA CAATGGCACC ACCTGTAATG ATTTTGCAAG TTCTGGAGAC 540
 CCAACTACA ACCTCATTTA CAGCATGTGT CTAACACTGT TGGGGTTCCT TATTCCTCTT 600
 TTTGTGATGT GTTCTTTTAA TTACAAGATT GCTCTCTTCC TAAAGCAGAG GAATAGGCAG 660
 GTTGCTACTG CTCTGCCCCT TGAAAAGCCT CTCAACTTGG TCATCATGGC AGTGGTAATC 720
 20 TTCTCTGTGC TTTTACACC CTATCAGTC ATGCGGAATG TGAGGATCGC TTCACGCCTG 780
 GGGAGTTGGA AGCAGTATCA GTGCACTCAG GTCGTCATCA ACTCCTTTTA CATTGTGACA 840
 CGGCCTTTGG CCTTTCTGAA CAGTGTATC AACCTGTCT TCTATTTTCT TTTGGGAGAT 900
 CACTTCAGGG ACATGCTGAT GAATCAACTG AGACACAACT TCAAATCCCT TACATCCTTT 960
 AGCAGATGGG CTCATGAACT CCTACTTTCA TTCAGAGAAA AGTGA 1005

25 (39) INFORMATION FOR SEQ ID NO:38:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 334 amino acids
 (B) TYPE: amino acid
 (C) STRANDEDNESS:
 (D) TOPOLOGY: not relevant

30 (ii) MOLECULE TYPE: protein

- 46 -

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:38:

	Met	Leu	Gly	Ile	Met	Ala	Trp	Asn	Ala	Thr	Cys	Lys	Asn	Trp	Leu	Ala	
	1				5					10					15		
5	Ala	Glu	Ala	Ala	Leu	Glu	Lys	Tyr	Tyr	Leu	Ser	Ile	Phe	Tyr	Gly	Ile	
				20					25					30			
	Glu	Phe	Val	Val	Gly	Val	Leu	Gly	Asn	Thr	Ile	Val	Val	Tyr	Gly	Tyr	
			35				40						45				
	Ile	Phe	Ser	Leu	Lys	Asn	Trp	Asn	Ser	Ser	Asn	Ile	Tyr	Leu	Phe	Asn	
		50					55					60					
10	Leu	Ser	Val	Ser	Asp	Leu	Ala	Phe	Leu	Cys	Thr	Leu	Pro	Met	Leu	Ile	
	65					70					75				80		
	Arg	Ser	Tyr	Ala	Asn	Gly	Asn	Trp	Ile	Tyr	Gly	Asp	Val	Leu	Cys	Ile	
					85					90					95		
15	Ser	Asn	Arg	Tyr	Val	Leu	His	Ala	Asn	Leu	Tyr	Thr	Ser	Ile	Leu	Phe	
				100					105					110			
	Leu	Thr	Phe	Ile	Ser	Ile	Asp	Arg	Tyr	Leu	Ile	Ile	Lys	Tyr	Pro	Phe	
		115						120					125				
	Arg	Glu	His	Leu	Leu	Gln	Lys	Lys	Glu	Phe	Ala	Ile	Leu	Ile	Ser	Leu	
		130					135					140					
20	Ala	Ile	Trp	Val	Leu	Val	Thr	Leu	Glu	Leu	Leu	Pro	Ile	Leu	Pro	Leu	
	145					150					155					160	
	Ile	Asn	Pro	Val	Ile	Thr	Asp	Asn	Gly	Thr	Thr	Cys	Asn	Asp	Phe	Ala	
				165					170						175		
25	Ser	Ser	Gly	Asp	Pro	Asn	Tyr	Asn	Leu	Ile	Tyr	Ser	Met	Cys	Leu	Thr	
				180					185					190			
	Leu	Leu	Gly	Phe	Leu	Ile	Pro	Leu	Phe	Val	Met	Cys	Phe	Phe	Tyr	Tyr	
			195					200					205				
	Lys	Ile	Ala	Leu	Phe	Leu	Lys	Gln	Arg	Asn	Arg	Gln	Val	Ala	Thr	Ala	
		210						215					220				
30	Leu	Pro	Leu	Glu	Lys	Pro	Leu	Asn	Leu	Val	Ile	Met	Ala	Val	Val	Ile	
	225					230					235					240	
	Phe	Ser	Val	Leu	Phe	Thr	Pro	Tyr	His	Val	Met	Arg	Asn	Val	Arg	Ile	
				245						250					255		
35	Ala	Ser	Arg	Leu	Gly	Ser	Trp	Lys	Gln	Tyr	Gln	Cys	Thr	Gln	Val	Val	
				260					265					270			
	Ile	Asn	Ser	Phe	Tyr	Ile	Val	Thr	Arg	Pro	Leu	Ala	Phe	Leu	Asn	Ser	

- 47 -

275 280 285

Val Ile Asn Pro Val Phe Tyr Phe Leu Leu Gly Asp His Phe Arg Asp
290 295 300

Met Leu Met Asn Gln Leu Arg His Asn Phe Lys Ser Leu Thr Ser Phe
5 305 310 315 320

Ser Arg Trp Ala His Glu Leu Leu Leu Ser Phe Arg Glu Lys
325 330

(40) INFORMATION FOR SEQ ID NO:39:

10 (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 1296 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

15 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:39:

ATGCAGGCGC TTAACATTAC CCCGGAGCAG TTCTCTCGGC TGCTGCGGGA CCACAACCTG 60

ACGCGGGAGC AGTTCATCGC TCTGTACCGG CTGCGACCGC TCGTCTACAC CCCAGAGCTG 120

CCGGGACGCG CCAAGCTGGC CCTCGTGCTC ACCGGCGTGC TCATCTTCGC CCTGGCGCTC 180

TTTGGAATG CTCTGGTGTT CTACGTGGTG ACCCGCAGCA AGGCCATGCG CACCGTCACC 240

20 AACATCTTTA TCTGCTCCTT GCGGCTCAGT GACCTGCTCA TCACCTTCTT CTGCATTCCC 300

GTCACCATGC TCCAGAACAT TTCCGACAAC TGGCTGGGGG GTGCTTTCAT TTGCAAGATG 360

GTGCCATTG TCCAGTCTAC CGCTGTTGTG ACAGAAATGC TCACTATGAC CTGCATTGCT 420

GTGGAAAGGC ACCAGGGACT TGTGCATCCT TTAAAATGA AGTGGCAATA CACCAACCGA 480

AGGGCTTTCA CAATGCTAGG TGTGGTCTGG CTGGTGGCAG TCATCGTAGG ATCACCCATG 540

25 TGGCACGTGC AACAACTTGA GATCAAATAT GACTTCCTAT ATGAAAAGGA ACACATCTGC 600

TGCTTAGAAG AGTGGACCAG CCCTGTGCAC CAGAAGATCT ACACCACCTT CATCCTTGTC 660

ATCCTCTTCC TCCTGCCTCT TATGGTGATG CTTATTCTGT ACAGTAAAAT TGGTTATGAA 720

CTTTGGATAA AGAAAAGAGT TGGGGATGGT TCAGTGCTTC GAACTATTCA TGGAAAAGAA 780

ATGTCCAAAA TAGCCAGGAA GAAGAAACGA GCTGTCATTA TGATGGTGAC AGTGGTGGCT 840

30 CTCTTTGCTG TGTGCTGGGC ACCATTCCAT GTTGTCATA TGATGATTGA ATACAGTAAT 900

TTTGAAAAGG AATATGATGA TGTACAATC AAGATGATTT TTGCTATCGT GCAAATTATT 960

- 48 -

GGATTTTCCA ACTCCATCTG TAATCCCATT GTCTATGCAT TTATGAATGA AAACCTTCAAA 1020
 AAAAAATGTTT TGTCTGCAGT TTGTTATTGC ATAGTAAATA AAACCTTCTC TCCAGCACAA 1080
 AGGCATGGAA ATTCAGGAAT TACAATGATG CGGAAGAAAG CAAAGTTTTC CCTCAGAGAG 1140
 AATCCAGTGG AGGAAACCAA AGGAGAAGCA TTCAGTGATG GCAACATTGA AGTCAAATTG 1200
 5 TGTGAACAGA CAGAGGAGAA GAAAAAGCTC AAACGACATC TTGCTCTCTT TAGGTCTGAA 1260
 CTGGCTGAGA ATTCTCCTTT AGACAGTGGG CATTAA 1296

(41) INFORMATION FOR SEQ ID NO:40:

(i) SEQUENCE CHARACTERISTICS:

10

- (A) LENGTH: 431 amino acids
 (B) TYPE: amino acid
 (C) STRANDEDNESS:
 (D) TOPOLOGY: not relevant

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:40:

15 Met Gln Ala Leu Asn Ile Thr Pro Glu Gln Phe Ser Arg Leu Leu Arg
 1 5 10 15
 Asp His Asn Leu Thr Arg Glu Gln Phe Ile Ala Leu Tyr Arg Leu Arg
 20 20 25 30
 Pro Leu Val Tyr Thr Pro Glu Leu Pro Gly Arg Ala Lys Leu Ala Leu
 20 35 40 45
 Val Leu Thr Gly Val Leu Ile Phe Ala Leu Ala Leu Phe Gly Asn Ala
 50 55 60
 Leu Val Phe Tyr Val Val Thr Arg Ser Lys Ala Met Arg Thr Val Thr
 65 70 75 80
 25 Asn Ile Phe Ile Cys Ser Leu Ala Leu Ser Asp Leu Leu Ile Thr Phe
 85 90 95
 Phe Cys Ile Pro Val Thr Met Leu Gln Asn Ile Ser Asp Asn Trp Leu
 100 105 110
 30 Gly Gly Ala Phe Ile Cys Lys Met Val Pro Phe Val Gln Ser Thr Ala
 115 120 125
 Val Val Thr Glu Met Leu Thr Met Thr Cys Ile Ala Val Glu Arg His
 130 135 140
 Gln Gly Leu Val His Pro Phe Lys Met Lys Trp Gln Tyr Thr Asn Arg
 145 150 155 160

- 49 -

Arg Ala Phe Thr Met Leu Gly Val Val Trp Leu Val Ala Val Ile Val
 165 170 175
 Gly Ser Pro Met Trp His Val Gln Gln Leu Glu Ile Lys Tyr Asp Phe
 180 185 190
 5 Leu Tyr Glu Lys Glu His Ile Cys Cys Leu Glu Glu Trp Thr Ser Pro
 195 200 205
 Val His Gln Lys Ile Tyr Thr Thr Phe Ile Leu Val Ile Leu Phe Leu
 210 215 220
 10 Leu Pro Leu Met Val Met Leu Ile Leu Tyr Ser Lys Ile Gly Tyr Glu
 225 230 235 240
 Leu Trp Ile Lys Lys Arg Val Gly Asp Gly Ser Val Leu Arg Thr Ile
 245 250 255
 His Gly Lys Glu Met Ser Lys Ile Ala Arg Lys Lys Lys Arg Ala Val
 260 265 270
 15 Ile Met Met Val Thr Val Val Ala Leu Phe Ala Val Cys Trp Ala Pro
 275 280 285
 Phe His Val Val His Met Met Ile Glu Tyr Ser Asn Phe Glu Lys Glu
 290 295 300
 20 Tyr Asp Asp Val Thr Ile Lys Met Ile Phe Ala Ile Val Gln Ile Ile
 305 310 315 320
 Gly Phe Ser Asn Ser Ile Cys Asn Pro Ile Val Tyr Ala Phe Met Asn
 325 330 335
 Glu Asn Phe Lys Lys Asn Val Leu Ser Ala Val Cys Tyr Cys Ile Val
 340 345 350
 25 Asn Lys Thr Phe Ser Pro Ala Gln Arg His Gly Asn Ser Gly Ile Thr
 355 360 365
 Met Met Arg Lys Lys Ala Lys Phe Ser Leu Arg Glu Asn Pro Val Glu
 370 375 380
 30 Glu Thr Lys Gly Glu Ala Phe Ser Asp Gly Asn Ile Glu Val Lys Leu
 385 390 395 400
 Cys Glu Gln Thr Glu Glu Lys Lys Lys Leu Lys Arg His Leu Ala Leu
 405 410 415
 Phe Arg Ser Glu Leu Ala Glu Asn Ser Pro Leu Asp Ser Gly His
 420 425 430

35 (42) INFORMATION FOR SEQ ID NO:41:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 24 base pairs

- 50 -

- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

5 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:41:

CTGTGTACAG CAGTTCGCAG AGTG

24

(43) INFORMATION FOR SEQ ID NO:42:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 24 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:42:

15 GAGTGCCAGG CAGAGCAGGT AGAC

24

(44) INFORMATION FOR SEQ ID NO:43:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 31 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(iv) ANTI-SENSE: NO

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:43:

25 CCCGAATTCC TGCTTGCTCC CAGCTTGGCC C

31

(45) INFORMATION FOR SEQ ID NO:44:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 32 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(iv) ANTI-SENSE: YES

- 51 -

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:44:

TGTGGATCCT GCTGTCAAAG GTCCCATTCC GG

32

(46) INFORMATION FOR SEQ ID NO:45:

(i) SEQUENCE CHARACTERISTICS:

- 5 (A) LENGTH: 20 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

10 (iv) ANTI-SENSE: NO

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:45:

TCACAATGCT AGGTGTGGTC

20

(47) INFORMATION FOR SEQ ID NO:46:

(i) SEQUENCE CHARACTERISTICS:

- 15 (A) LENGTH: 22 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

20 (iv) ANTI-SENSE: YES

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:46:

TGCATAGACA ATGGGATTAC AG

22

(48) INFORMATION FOR SEQ ID NO:47:

(i) SEQUENCE CHARACTERISTICS:

- 25 (A) LENGTH: 511 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

30 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:47:

TCACAATGCT AGGTGTGGTC TGGCTGGTGG CAGTCATCGT AGGATCACCC ATGTGGCACG

60

TGCAACAAC TGAATCAAA TATGACTTCC TATATGAAAA GGAACACATC TGCTGCTTAG

120

- 52 -

AAGAGTGGAC CAGCCCTGTG CACCAGAAGA TCTACACCAC CTTTCATCCTT GTCATCCTCT 180
TCCTCCTGCC TCTTATGGTG ATGCTTATTC TGTACGTAAA ATTGGTTATG AACTTTGGAT 240
AAAGAAAAGA GTTGGGGATG GTTCAGTGCT TCGAACTATT CATGGAAAAG AAATGTCCAA 300
AATAGCCAGG AAGAAGAAAC GAGCTGTCAT TATGATGGTG ACAGTGGTGG CTCTCTTTGC 360
5 TGTGTGCTGG GCACCATTC CATTGTGTCCA TATGATGATT GAATACAGTA ATTTTGAAAA 420
GGAATATGAT GATGTCACAA TCAAGATGAT TTTTGCTATC GTGCAAATTA TTGGATTTTC 480
CAACTCCATC TGTAATCCCA TTGTCTATGC A 511

(49) INFORMATION FOR SEQ ID NO:48:

10 (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 21 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

15 (iv) ANTI-SENSE: NO

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:48:

CTGCTTAGAA GAGTGGACCA G 21

(50) INFORMATION FOR SEQ ID NO:49:

20 (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 22 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

25 (iv) ANTI-SENSE: NO

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:49:

CTGTGCACCA GAAGATCTAC AC 22

(51) INFORMATION FOR SEQ ID NO:50:

30 (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 21 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

- 53 -

(ii) MOLECULE TYPE: DNA (genomic)

(iv) ANTI-SENSE: YES

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:50:

CAAGGATGAA GGTGGTGTAG A

21

5 (52) INFORMATION FOR SEQ ID NO:51:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 23 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

10 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(iv) ANTI-SENSE: YES

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:51:

GTGTAGATCT TCTGGTGCAC AGG

23

15 (53) INFORMATION FOR SEQ ID NO:52:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 21 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

20 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:52:

GCAATGCAGG TCATAGTGAG C

21

(54) INFORMATION FOR SEQ ID NO:53:

25 (i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 27 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

30 (ii) MOLECULE TYPE: DNA (genomic)

(iii) HYPOTHETICAL: YES

(iv) ANTI-SENSE: YES

- 54 -

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:53:

TGGAGCATGG TGACGGGAAT GCAGAAG

27

(55) INFORMATION FOR SEQ ID NO:54:

(i) SEQUENCE CHARACTERISTICS:

- 5 (A) LENGTH: 27 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

10 (iv) ANTI-SENSE: YES

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:54:

GTGATGAGCA GGTCACTGAG CGCCAAG

27

(56) INFORMATION FOR SEQ ID NO:55:

(i) SEQUENCE CHARACTERISTICS:

- 15 (A) LENGTH: 23 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

20 (iv) ANTI-SENSE: NO

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:55:

GCAATGCAGG CGCTTAACAT TAC

23

(57) INFORMATION FOR SEQ ID NO:56:

(i) SEQUENCE CHARACTERISTICS:

- 25 (A) LENGTH: 22 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

30 (iv) ANTI-SENSE: YES

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:56:

TTGGGTTACA ATCTGAAGGG CA

22

- 55 -

(58) INFORMATION FOR SEQ ID NO:57:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 23 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(iv) ANTI-SENSE: NO

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:57:

10 ACTCCGTGTC CAGCAGGACT CTG

23

(58) INFORMATION FOR SEQ ID NO:58:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 24 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(iv) ANTI-SENSE: YES

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:58:

20 TGGCGTGTTC TGGACCCTCA CGTG

24

(58) INFORMATION FOR SEQ ID NO:59:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 29 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(iv) ANTI-SENSE: NO

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:59:

30 CAGGCCTTGG ATTTTAATGT CAGGGATGG

29

(61) INFORMATION FOR SEQ ID NO:60:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 27 base pairs

- 56 -

- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

5 (iv) ANTI-SENSE: YES

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:60:

GGAGAGTCAG CTCTGAAAGA ATTCAGG

27

(62) INFORMATION FOR SEQ ID NO:61:

- 10 (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 27 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

15 (iv) ANTI-SENSE: NO

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:61:

TGATGTGATG CCAGATACTA ATAGCAC

27

(63) INFORMATION FOR SEQ ID NO:62:

- 20 (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 27 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

25 (iv) ANTI-SENSE: YES

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:62:

CCTGATTCAT TTAGGTGAGA TTGAGAC

27

(64) INFORMATION FOR SEQ ID NO:63:

- 30 (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 26 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

- 57 -

(ii) MOLECULE TYPE: DNA (genomic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:63:

CCCAAGCTTC CCCAGGTGTA TTTGAT

26

(3) INFORMATION FOR SEQ ID NO:63:

- 5 (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 26 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

10 (ii) MOLECULE TYPE: DNA (genomic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:64:

GTTGGATCCA CATAATGCAT TTTCTC

26

(66) INFORMATION FOR SEQ ID NO:65:

- 15 (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 1080 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

20 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:65:

ATGATTCTCA ACTCTTCTAC TGAAGATGGT ATTAAAAGAA TCCAAGATGA TTGTCCCAAA 60
GCTGGAAGGC ATAATTACAT ATTTGTTCATG ATTCTACTT TATACAGTAT CATCTTTGTG 120
GTGGGAATAT TTGGAAACAG CTTGGTGGTG ATAGTCATTT ACTTTTATAT GAAGCTGAAG 180
ACTGTGGCCA GTGTTTTTCT TTTGAATTGA GCACTGGCTG ACTTATGCTT TTTACTGACT 240
25 TTGCCACTAT GGGCTGTCTA CACAGCTATG GAATACCGCT GGCCCTTTGG CAATTACCTA 300
TGTAAGATTG CTTCAGCCAG CGTCAGTTTC AACCTGTACG CTAGTGTGTT TCTACTCAG 360
TGTCTCAGCA TTGATCGATA CCTGGCTATT GTTCACCCAA TGAAGTCCCG CCTTCGACGC 420
ACAATGCTTG TAGCCAAAGT CACCTGCATC ATCATTGGC TGCTGGCAGG CTTGGCCAGT 480
TTGCCAGCTA TAATCCATCG AAATGTATTT TTCATTGAGA ACACCAATAT TACAGTTTGT 540
30 GCTTTCCATT ATGAGTCCCA AAATCAACC CTTCCGATAG GGCTGGGCCT GACCAAAAAT 600

- 58 -

ATACTGGGTT TCCTGTTTCC TTTTCTGATC ATTCTTACAA GTTATACTCT TATTGGAAG 660
 GCCCTAAAGA AGGCTTATGA AATTCAGAAG AACAAACCAA GAAATGATGA TATTTTAAAG 720
 ATAATTATGG CAATTGTGCT TTTCTTTTTC TTTTCCTGGA TTCCCCACCA AATATTCACT 780
 TTTCTGGATG TATTGATTCA ACTAGGCATC ATACGTGACT GTAGAATTGC AGATATTGTG 840
 5 GACACGGCCA TGCCTATCAC CATTTGTATA GCTTATTTTA ACAATTGCCT GAATCCTCTT 900
 TTTTATGGCT TTCTGGGGAA AAAATTTAAA AGATATTTTC TCCAGCTTCT AAAATATATT 960
 CCCCCAAAAG CCAAATCCCA CTCAAACCTT TCAACAAAAA TGAGCACGCT TTCCTACCGC 1020
 CCCTCAGATA ATGTAAGCTC ATCCACCAAG AAGCCTGCAC CATGTTTGA GGTGAGTGA 1080

(67) INFORMATION FOR SEQ ID NO:66:

- 10 (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 359 amino acids
 (B) TYPE: amino acid
 (C) STRANDEDNESS:
 (D) TOPOLOGY: not relevant

- 15 (ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:66:

Met Ile Leu Asn Ser Ser Thr Glu Asp Gly Ile Lys Arg Ile Gln Asp
 1 5 10 15
 Asp Cys Pro Lys Ala Gly Arg His Asn Tyr Ile Phe Val Met Ile Pro
 20 20 25 30
 Thr Leu Tyr Ser Ile Ile Phe Val Val Gly Ile Phe Gly Asn Ser Leu
 35 40 45
 Val Val Ile Val Ile Tyr Phe Tyr Met Lys Leu Lys Thr Val Ala Ser
 50 55 60
 25 Val Phe Leu Leu Asn Leu Ala Leu Ala Asp Leu Cys Phe Leu Leu Thr
 65 70 75 80
 Leu Pro Leu Trp Ala Val Tyr Thr Ala Met Glu Tyr Arg Trp Pro Phe
 85 90 95
 30 Gly Asn Tyr Leu Cys Lys Ile Ala Ser Ala Ser Val Ser Phe Asn Leu
 100 105 110
 Tyr Ala Ser Val Phe Leu Leu Thr Cys Leu Ser Ile Asp Arg Tyr Leu
 115 120 125
 Ala Ile Val His Pro Met Lys Ser Arg Leu Arg Arg Thr Met Leu Val

- 59 -

	130		135		140
	Ala Lys Val Thr Cys Ile Ile Ile Trp Leu Leu Ala Gly Leu Ala Ser				
	145		150		155 160
5	Leu Pro Ala Ile Ile His Arg Asn Val Phe Phe Ile Glu Asn Thr Asn				
		165		170	175
	Ile Thr Val Cys Ala Phe His Tyr Glu Ser Gln Asn Ser Thr Leu Pro				
		180		185	190
	Ile Gly Leu Gly Leu Thr Lys Asn Ile Leu Gly Phe Leu Phe Pro Phe				
		195		200	205
10	Leu Ile Ile Leu Thr Ser Tyr Thr Leu Ile Trp Lys Ala Leu Lys Lys				
		210		215	220
	Ala Tyr Glu Ile Gln Lys Asn Lys Pro Arg Asn Asp Asp Ile Phe Lys				
		225		230	235 240
15	Ile Ile Met Ala Ile Val Leu Phe Phe Phe Phe Ser Trp Ile Pro His				
		245		250	255
	Gln Ile Phe Thr Phe Leu Asp Val Leu Ile Gln Leu Gly Ile Ile Arg				
		260		265	270
	Asp Cys Arg Ile Ala Asp Ile Val Asp Thr Ala Met Pro Ile Thr Ile				
		275		280	285
20	Cys Ile Ala Tyr Phe Asn Asn Cys Leu Asn Pro Leu Phe Tyr Gly Phe				
		290		295	300
	Leu Gly Lys Lys Phe Lys Arg Tyr Phe Leu Gln Leu Leu Lys Tyr Ile				
		305		310	315 320
25	Pro Pro Lys Ala Lys Ser His Ser Asn Leu Ser Thr Lys Met Ser Thr				
		325		330	335
	Leu Ser Tyr Arg Pro Ser Asp Asn Val Ser Ser Ser Thr Lys Lys Pro				
		340		345	350
	Ala Pro Cys Phe Glu Val Glu				
		355			

30 (68) INFORMATION FOR SEQ ID NO:67:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 27 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

35 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

- 60 -

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:67:

ACCATGGGCA GCCCCTGGAA CGGCAGC

27

(69) INFORMATION FOR SEQ ID NO:68:

(i) SEQUENCE CHARACTERISTICS:

- 5 (A) LENGTH: 39 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

10 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:68:

AGAACCACCA CCAGCAGGAC GCGGACGGTC TGCCGGTGG

39

(70) INFORMATION FOR SEQ ID NO:69:

(i) SEQUENCE CHARACTERISTICS:

- 15 (A) LENGTH: 39 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:69:

20 GTCCGCGTCC TGCTGGTGGT GGTCTGGCA TTTATAATT

39

(71) INFORMATION FOR SEQ ID NO:70:

(i) SEQUENCE CHARACTERISTICS:

- 25 (A) LENGTH: 33 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: not relevant

(ii) MOLECULE TYPE: DNA (genomic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:70:

CCTGGATCCT TATCCCATCG TCTTCAGTT AGC

33

30 (72) INFORMATION FOR SEQ ID NO:71:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 26 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single

- 61 -

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(iv) ANTI-SENSE: NO

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:71:

5 CTGGAATTCT CCTGCCAGCA TGGTGA
26

(73) INFORMATION FOR SEQ ID NO:72:

(i) SEQUENCE CHARACTERISTICS:

10 (A) LENGTH: 30 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(iv) ANTI-SENSE: YES

15 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:72:

GCAGGATCCT ATATTGCGTG CTCTGTCCCC
30

(74) INFORMATION FOR SEQ ID NO:73:

(i) SEQUENCE CHARACTERISTICS:

20 (A) LENGTH: 999 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

25 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:73:

ATGGTGAACT CCACCCACCG TGGGATGCAC ACTTCTCTGC ACCTCTGGAA CCGCAGCAGT 60
TACAGACTGC ACAGCAATGC CAGTGAGTCC CTTGGAAAAG GCTACTCTGA TGGAGGGTGC 120
TACGAGCAAC TTTTGTCTC TCCTGAGGTG TTTGTGACTC TGGGTGTCAT CAGCTTGTTC 180
GAGAATATCT TAGTGATTGT GGCAATAGCC AAGAACAAGA ATCTGCATTC ACCCATGTAC 240
30 TTTTTCATCT GCAGCTTGGC TGTGGCTGAT ATGCTGGTGA GCGTTTCAA TGGATCAGAA 300
ACCATTATCA TCACCTATT AACAGTACA GATACGGATG CACAGAGTTT CACAGTGAAT 360
ATTGATAATG TCATTGACTC GGTGATCTGT AGCTCCTTGC TTGCATCCAT TTGCAGCCTG 420

- 62 -

CTTTCAATTG CAGTGGACAG GTACTTTACT ATCTTCTATG CTCTCCAGTA CCATAACATT 480
 ATGACAGTTA AGCGGGTTGG GATCAGCATA AGTTGTATCT GGGCAGCTTG CACGGTTTCA 540
 GGCATTTTGT TCATCATTTA CTCAGATAGT AGTGCTGTCA TCATCTGCCT CATCACCATG 600
 TTCTTCACCA TGCTGGCTCT CATGGCTTCT CTCTATGTCC ACATGTTCTT GATGGCCAGG 660
 5 CTTACACATTA AGAGGATTGC TGTCTCCCCC GGCCTGGTG CCATCCGCCA AGGTGCCAAT 720
 ATGAAGGGAG CGATTACCTT GACCATCCTG ATTGGCGTCT TTGTTGTCTG CTGGGCCCCA 780
 TTCTTCCTCC ACTTAATATT CTACATCTCT TGTCTCAGA ATCCATATTG TGTGTGCTTC 840
 ATGTCTCACT TTAACCTGTA TCTCATACTG ATCATGTGTA ATTCAATCAT CGATCCTCTG 900
 ATTTATGCAC TCCGGAGTCA AGAACTGAGG AAAACCTTCA AAGAGATCAT CTGTTGCTAT 960
 10 CCCCTGGGAG GCCTTTGTGA CTTGTCTAGC AGATATTAA 999

(75) INFORMATION FOR SEQ ID NO:74:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 332 amino acids

(B) TYPE: amino acid

(C) STRANDEDNESS:

(D) TOPOLOGY: not relevant

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:74:

Met Val Asn Ser Thr His Arg Gly Met His Thr Ser Leu His Leu Trp
 1 5 10 15
 Asn Arg Ser Ser Tyr Arg Leu His Ser Asn Ala Ser Glu Ser Leu Gly
 20 25 30
 Lys Gly Tyr Ser Asp Gly Gly Cys Tyr Glu Gln Leu Phe Val Ser Pro
 35 40 45
 25 Glu Val Phe Val Thr Leu Gly Val Ile Ser Leu Leu Glu Asn Ile Leu
 50 55 60
 Val Ile Val Ala Ile Ala Lys Asn Lys Asn Leu His Ser Pro Met Tyr
 65 70 75 80
 30 Phe Phe Ile Cys Ser Leu Ala Val Ala Asp Met Leu Val Ser Val Ser
 85 90 95
 Asn Gly Ser Glu Thr Ile Ile Ile Thr Leu Leu Asn Ser Thr Asp Thr
 100 105 110
 Asp Ala Gln Ser Phe Thr Val Asn Ile Asp Asn Val Ile Asp Ser Val

- 63 -

	115		120		125
	Ile Cys Ser Ser Leu Leu Ala Ser Ile Cys Ser Leu Leu Ser Ile Ala				
	130		135		140
5	Val Asp Arg Tyr Phe Thr Ile Phe Tyr Ala Leu Gln Tyr His Asn Ile				
	145		150		155 160
	Met Thr Val Lys Arg Val Gly Ile Ser Ile Ser Cys Ile Trp Ala Ala				
		165		170	175
	Cys Thr Val Ser Gly Ile Leu Phe Ile Ile Tyr Ser Asp Ser Ser Ala				
		180		185	190
10	Val Ile Ile Cys Leu Ile Thr Met Phe Phe Thr Met Leu Ala Leu Met				
		195		200	205
	Ala Ser Leu Tyr Val His Met Phe Leu Met Ala Arg Leu His Ile Lys				
		210		215	220
15	Arg Ile Ala Val Leu Pro Gly Thr Gly Ala Ile Arg Gln Gly Ala Asn				
		225		230	235 240
	Met Lys Gly Ala Ile Thr Leu Thr Ile Leu Ile Gly Val Phe Val Val				
		245		250	255
	Cys Trp Ala Pro Phe Phe Leu His Leu Ile Phe Tyr Ile Ser Cys Pro				
		260		265	270
20	Gln Asn Pro Tyr Cys Val Cys Phe Met Ser His Phe Asn Leu Tyr Leu				
		275		280	285
	Ile Leu Ile Met Cys Asn Ser Ile Ile Asp Pro Leu Ile Tyr Ala Leu				
		290		295	300
25	Arg Ser Gln Glu Leu Arg Lys Thr Phe Lys Glu Ile Ile Cys Cys Tyr				
		305		310	315 320
	Pro Leu Gly Gly Leu Cys Asp Leu Ser Ser Arg Tyr				
		325		330	

(76) INFORMATION FOR SEQ ID NO:75:

30 (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 32 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

35 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:75:

CCGAAGCTTC GAGCTGAGTA AGGCGGCGGG CT

32

- 64 -

(77) INFORMATION FOR SEQ ID NO:76:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 31 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:76:

GTGGAATTCA TTTGCCCTGC CTCACCCCCC A

31

10 (78) INFORMATION FOR SEQ ID NO:77:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 1344 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:77:

ATGGAGCTGC TAAAGCTGAA CCGGAGCGTG CAGGGAACCG GACCCGGGCC GGGGGCTTCC 60
CTGTGCCGCC CGGGGCGGCC TCTCCTCAAC AGCAGCAGTG TGGGCAACCT CAGCTGCGAG 120
20 CCCCCTCGCA TTCGCGGAGC CGGGACACGA GAATTGGAGC TGGCCATTAG AATCACTCTT 180
TACGCAGTGA TCTTCCTGAT GAGCGTTGGA GGAAATATGC TCATCATCGT GGTCTTGGGA 240
CTGAGCCGCC GCCTGAGGAC TGTCACCAAT GCCTTCCTCC TCTCACTGGC AGTCAGCGAC 300
CTCCTGCTGG CTGTGGCTTG CATGCCCTTC ACCCTCCTGC CCAATCTCAT GGGCACATTC 360
ATCTTTGGCA CCGTCATCTG CAAGGCGGTT TCCTACCTCA TGGGGGTGTC TGTGAGTGTG 420
25 TCCACGCTAA GCCTCGTGCC CATCGCACTG GAGCGATATA GCGCCATCTG CCGACCACTG 480
CAGGCACGAG TGTGGCAGAC GCGCTCCAC GCGGCTCGCG TGATTGTAGC CACGTGGCTG 540
CTGTCCGGAC TACTCATGGT GCCCTACCCC GTGTACACTG TCGTGCAACC AGTGGGGCCT 600
CGTGTGCTGC AGTGCGTGCA TCGCTGGCCC AGTGCGCGGG TCCGCCAGAC CTGGTCCGTA 660
CTGCTGCTTC TGCTCTTGTT CTTCATCCCA GGTGTGGTTA TGGCCGTGGC CTACGGGCTT 720
30 ATCTCTCGCG AGCTCTACTT AGGGCTTCGC TTGACGGCG ACAGTGACAG CGACAGCCAA 780
AGCAGGTGCC GAAACCAAGG CGGGCTGCCA GGGGCTGTTC ACCAGAACGG GCGTTGCCGG 840

- 65 -

CCTGAGACTG GCGCGGTTGG CAAAGACAGC GATGGCTGCT ACGTGCAACT TCCACGTTCC 900
 CGGCCTGCCC TGGAGCTGAC GCGCTGACG GCTCCTGGGC CGGGATCCGG CTCCCGGCCC 960
 ACCCAGGCCA AGCTGCTGGC TAAGAAGCGC GTGGTGCGAA TGTGTCTGGT GATCGTTGTG 1020
 CTTTTTTTTT TGTGTTGGTT GCCAGTTTAT AGTGCCAACA CGTGGCGCGC CTTTGATGGC 1080
 5 CCGGGTGCAC ACCGAGCACT CTCGGGTGCT CCTATCTCCT TCATTCACTT GCTGAGCTAC 1140
 GCCTCGGCCT GTGTCAACCC CCTGGTCTAC TGCTTCATGC ACCGTCGCTT TCGCCAGGCC 1200
 TGCCTGGAAA CTTGCGCTCG CTGCTGCCCC CGGCCTCCAC GAGCTCGCCC CAGGGCTCTT 1260
 CCCGATGAGG ACCCTCCAC TCCCTCCATT GCTTCGCTGT CCAGGCTTAG CTACACCACC 1320
 ATCAGCACAC TGGGCCCTGG CTGA 1344

10 (79) INFORMATION FOR SEQ ID NO:78:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 447 amino acids

(B) TYPE: amino acid

(C) STRANDEDNESS:

15 (D) TOPOLOGY: not relevant

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:78:

Met Glu Leu Leu Lys Leu Asn Arg Ser Val Gln Gly Thr Gly Pro Gly
 1 5 10 15
 20 Pro Gly Ala Ser Leu Cys Arg Pro Gly Ala Pro Leu Leu Asn Ser Ser
 20 25 30
 Ser Val Gly Asn Leu Ser Cys Glu Pro Pro Arg Ile Arg Gly Ala Gly
 35 40 45
 25 Thr Arg Glu Leu Glu Leu Ala Ile Arg Ile Thr Leu Tyr Ala Val Ile
 50 55 60
 Phe Leu Met Ser Val Gly Gly Asn Met Leu Ile Ile Val Val Leu Gly
 65 70 75 80
 Leu Ser Arg Arg Leu Arg Thr Val Thr Asn Ala Phe Leu Leu Ser Leu
 85 90 95
 30 Ala Val Ser Asp Leu Leu Leu Ala Val Ala Cys Met Pro Phe Thr Leu
 100 105 110
 Leu Pro Asn Leu Met Gly Thr Phe Ile Phe Gly Thr Val Ile Cys Lys
 115 120 125

- 66 -

Ala Val Ser Tyr Leu Met Gly Val Ser Val Ser Val Ser Thr Leu Ser
130 135 140

Leu Val Ala Ile Ala Leu Glu Arg Tyr Ser Ala Ile Cys Arg Pro Leu
145 150 155 160

5 Gln Ala Arg Val Trp Gln Thr Arg Ser His Ala Ala Arg Val Ile Val
165 170 175

Ala Thr Trp Leu Leu Ser Gly Leu Leu Met Val Pro Tyr Pro Val Tyr
180 185 190

10 Thr Val Val Gln Pro Val Gly Pro Arg Val Leu Gln Cys Val His Arg
195 200 205

Trp Pro Ser Ala Arg Val Arg Gln Thr Trp Ser Val Leu Leu Leu Leu
210 215 220

Leu Leu Phe Phe Ile Pro Gly Val Val Met Ala Val Ala Tyr Gly Leu
225 230 235 240

15 Ile Ser Arg Glu Leu Tyr Leu Gly Leu Arg Phe Asp Gly Asp Ser Asp
245 250 255

Ser Asp Ser Gln Ser Arg Val Arg Asn Gln Gly Gly Leu Pro Gly Ala
260 265 270

20 Val His Gln Asn Gly Arg Cys Arg Pro Glu Thr Gly Ala Val Gly Lys
275 280 285

Asp Ser Asp Gly Cys Tyr Val Gln Leu Pro Arg Ser Arg Pro Ala Leu
290 295 300

Glu Leu Thr Ala Leu Thr Ala Pro Gly Pro Gly Ser Gly Ser Arg Pro
305 310 315 320

25 Thr Gln Ala Lys Leu Leu Ala Lys Lys Arg Val Val Arg Met Leu Leu
325 330 335

Val Ile Val Val Leu Phe Phe Leu Cys Trp Leu Pro Val Tyr Ser Ala
340 345 350

30 Asn Thr Trp Arg Ala Phe Asp Gly Pro Gly Ala His Arg Ala Leu Ser
355 360 365

Val Ala Pro Ile Ser Phe Ile His Leu Leu Ser Tyr Ala Ser Ala Cys
370 375 380

Val Asn Pro Leu Val Tyr Cys Phe Met His Arg Arg Phe Arg Gln Ala
385 390 395 400

35 Cys Leu Glu Thr Cys Ala Arg Cys Cys Pro Arg Pro Pro Arg Ala Arg
405 410 415

Pro Arg Ala Leu Pro Asp Glu Asp Pro Pro Thr Pro Ser Ile Ala Ser

- 67 -

420

425

430

Leu Ser Arg Leu Ser Tyr Thr Thr Ile Ser Thr Leu Gly Pro Gly
 435 440 445

(80) INFORMATION FOR SEQ ID NO:79:

- 5 (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 30 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

- 10 (ii) MOLECULE TYPE: DNA (genomic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:79:

TGCAAGCTTA AAAAGGAAAA AATGAACAGC

30

(81) INFORMATION FOR SEQ ID NO:80:

- 15 (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 30 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

- (ii) MOLECULE TYPE: DNA (genomic)

- 20 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:80:

TAAGGATCCC TTCCCTTCAA AACATCCTTG

30

(82) INFORMATION FOR SEQ ID NO:81:

- 25 (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 1014 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

- (ii) MOLECULE TYPE: DNA (genomic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:81:

30 ATGAACAGCA CATGTATTGA AGAACAGCAT GACCTGGATC ACTATTTGTT TCCCATTGTT 60
 TACATCTTTG TGATTATAGT CAGCATTCCA GCCAATATTG GATCTCTGTG TGTGTCTTTC 120
 CTGCAACCCA AGAAGGAAAG TGAAGTAGGA ATTTACCTCT TCAGTTTGTC ACTATCAGAT 180
 TTACTCTATG CATTAACCTCT CCCTTTATGG ATTGATTATA CTTGGAATAA AGACAACTGG 240

- 68 -

ACTTTCTCTC CTGCCTTG TG CAAAGGGAGT GCTTTTCTCA TGTACATGAA GTTTTACAGC 300
 AGCACAGCAT TCCTCACCTG CATTGCCGTT GATCGGTATT TGGCTGTTGT CTACCCCTTG 360
 AAGTTTTTTT TCCTAAGGAC AAGAAGAATT GCACTCATGG TCAGCCTGTC CATCTGGATA 420
 TTGGAAACCA TCTTCAATGC TGTCAATGTTG TGGGAAGATG AAACAGTTGT TGAATATTGC 480
 5 GATGCCGAAA AGTCTAATTT TACTTTATGC TATGACAAAT ACCCTTTAGA GAAATGGCAA 540
 ATCAACCTCA ACTTGTTTCTG GACGTGTACA GGCTATGCAA TACCTTTGGT CACCATCCTG 600
 ATCTGTAACC GGAAAGTCTA CCAAGCTGTG CGGCACAATA AAGCCACGGA AAACAAGGAA 660
 AAGAAGAGAA TCATAAACT ACTTGTCAGC ATCACAGTTA CTTTGTCTT ATGCTTTACT 720
 CCCTTTCATG TGATGTTGCT GATTCGCTGC ATTTTAGAGC ATGCTGTGAA CTTCAAGAC 780
 10 CACAGCAATT CTGGGAAGCG AACTTACACA ATGTATAGAA TCACGGTTGC ATTAACAAGT 840
 TTAAATTGTG TTGCTGATCC AATTCTGTAC TGTTTTGTTA CCGAAACAGG AAGATATGAT 900
 ATGTGGAATA TATTAAAATT CTGCACTGGG AGGTGTAATA CATCACAAAG ACAAAGAAAA 960
 CGCATACTTT CTGTGTCTAC AAAAGATACT ATGGAATTAG AGGTCCTTGA GTAG 1014

(83) INFORMATION FOR SEQ ID NO:82:

- 15 (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 337 amino acids
 - (B) TYPE: amino acid
 - (C) STRANDEDNESS:
 - (D) TOPOLOGY: not relevant

- 20 (ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:82:

Met Asn Ser Thr Cys Ile Glu Glu Gln His Asp Leu Asp His Tyr Leu
 1 5 10 15
 Phe Pro Ile Val Tyr Ile Phe Val Ile Ile Val Ser Ile Pro Ala Asn
 25 20 25 30
 Ile Gly Ser Leu Cys Val Ser Phe Leu Gln Pro Lys Lys Glu Ser Glu
 35 40 45
 Leu Gly Ile Tyr Leu Phe Ser Leu Ser Leu Ser Asp Leu Leu Tyr Ala
 50 55 60
 30 Leu Thr Leu Pro Leu Trp Ile Asp Tyr Thr Trp Asn Lys Asp Asn Trp
 65 70 75 80
 Thr Phe Ser Pro Ala Leu Cys Lys Gly Ser Ala Phe Leu Met Tyr Met

- 69 -

	85										90										95											
	Lys	Phe	Tyr	Ser	Ser	Thr	Ala	Phe	Leu	Thr	Cys	Ile	Ala	Val	Asp	Arg																
				100					105						110																	
5	Tyr	Leu	Ala	Val	Val	Tyr	Pro	Leu	Lys	Phe	Phe	Phe	Leu	Arg	Thr	Arg																
			115					120						125																		
	Arg	Ile	Ala	Leu	Met	Val	Ser	Leu	Ser	Ile	Trp	Ile	Leu	Glu	Thr	Ile																
			130				135							140																		
	Phe	Asn	Ala	Val	Met	Leu	Trp	Glu	Asp	Glu	Thr	Val	Val	Glu	Tyr	Cys																
						150								155		160																
10	Asp	Ala	Glu	Lys	Ser	Asn	Phe	Thr	Leu	Cys	Tyr	Asp	Lys	Tyr	Pro	Leu																
					165					170					175																	
	Glu	Lys	Trp	Gln	Ile	Asn	Leu	Asn	Leu	Phe	Arg	Thr	Cys	Thr	Gly	Tyr																
				180					185						190																	
15	Ala	Ile	Pro	Leu	Val	Thr	Ile	Leu	Ile	Cys	Asn	Arg	Lys	Val	Tyr	Gln																
			195					200						205																		
	Ala	Val	Arg	His	Asn	Lys	Ala	Thr	Glu	Asn	Lys	Glu	Lys	Lys	Arg	Ile																
			210				215						220																			
	Ile	Lys	Leu	Leu	Val	Ser	Ile	Thr	Val	Thr	Phe	Val	Leu	Cys	Phe	Thr																
						230						235				240																
20	Pro	Phe	His	Val	Met	Leu	Leu	Ile	Arg	Cys	Ile	Leu	Glu	His	Ala	Val																
					245					250					255																	
	Asn	Phe	Glu	Asp	His	Ser	Asn	Ser	Gly	Lys	Arg	Thr	Tyr	Thr	Met	Tyr																
				260					265					270																		
25	Arg	Ile	Thr	Val	Ala	Leu	Thr	Ser	Leu	Asn	Cys	Val	Ala	Asp	Pro	Ile																
			275					280						285																		
	Leu	Tyr	Cys	P																												

(84) INFORMATION FOR SEQ ID NO:83:

35 (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 40 base pairs
(B) TYPE: nucleic acid

- 70 -

(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:83:

5 CAGGAAGAAG AAACGAGCTG TCATTATGAT GGTGACAGTG
40

(85) INFORMATION FOR SEQ ID NO:84:

(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 40 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:84:

15 CACTGTCACC ATCATAATGA CAGCTCGTTT CTTCTTCCTG
40

(86) INFORMATION FOR SEQ ID NO:85:

(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 30 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:85:

25 GGCCACCGGC AGACCAAACG CGTCCTGCTG
30

(87) INFORMATION FOR SEQ ID NO:86:

(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 31 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:86:

- 71 -

CTCCTTCGGT CCTCCTATCG TTGTCAGAAG T
31

(88) INFORMATION FOR SEQ ID NO:87:

- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 37 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:87:

GGAAAAGAAG AGAATCAAAA AACTACTTGT CAGCATC

37

(89) INFORMATION FOR SEQ ID NO:88:

- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 31 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:88:

CTCCTTCGGT CCTCCTATCG TTGTCAGAAG T

31

(90) INFORMATION FOR SEQ ID NO:89:

- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 1080 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:89:

ATGATTCTCA ACTCTTCTAC TGAAGATGGT ATTAAAAGAA TCCAAGATGA TTGTCCCAA 60

GCTGGAAGGC ATAATTACAT ATTTGTTCATG ATTCCTACTT TATACAGTAT CATCTTTGTG 120

GTGGGAATAT TTGGAAACAG CTTGGTGGTG ATAGTCATTT ACTTTTATAT GAAGCTGAAG 180

ACTGTGGCCA GTGTTTTTCT TTTGAATTTA GCACTGGCTG ACTTATGCTT TTTACTGACT 240

TTGCCACTAT GGGCTGTCTA CACAGCTATG GAATACCGCT GGCCCTTTGG CAATTACCTA 300

- 72 -

TGTAAGATTG CTCAGCCAG CGTCAGTTTC AACCTGTACG CTAGTGTGTT TCTACTCAGC 360
 TGTCTCAGCA TTGATCGATA CCTGGCTATT GTTCACCCAA TGAAGTCCCG CCTTCGACGC 420
 ACAATGCTTG TAGCCAAAGT CACCTGCATC ATCATTGGC TGCTGGCAGG CTTGGCCAGT 480
 TTGCCAGCTA TAATCCATCG AAATGTATTT TTCATTGAGA ACACCAATAT TACAGTTTGT 540
 5 GCTTTCCATT ATGAGTCCCA AAATTCAACC CTCCGATAG GGCTGGGCCT GACCAAAAAT 600
 ATACTGGGTT TCCTGTTTCC TTTCTGATC ATTCTTACAA GTTATACTCT TATTTGGAAG 660
 GCCCTAAAGA AGGCTTATGA AATTCAGAAG AACAAACCAA GAAATGATGA TATTAAAAAG 720
 ATAATTATGG CAATTGTGCT TTTCTTTTTC TTTCTCTGGA TTCCCCACCA AATATTCAC 780
 TTTCTGGATG TATTGATTCA ACTAGGCATC ATACGTGACT GTAGAATTGC AGATATTGTG 840
 10 GACACGGCCA TGCCTATCAC CATTTGTATA GCTTATTTTA ACAATTGCCT GAATCCTCTT 900
 TTTTATGGCT TTCTGGGGAA AAAATTTAAA AGATATTTTC TCCAGCTTCT AAAATATATT 960
 CCCCCAAAAG CCAAATCCCA CTCAAACCTT TCAACAAAAA TGAGCACGCT TTCCTACCGC 1020
 CCCTCAGATA ATGTAAGCTC ATCCACCAAG AAGCCTGCAC CATGTTTTGA GGTGAGTGA 1080

(91) INFORMATION FOR SEQ ID NO:90:

- 15 (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 359 amino acids
 - (B) TYPE: amino acid
 - (C) STRANDEDNESS:
 - (D) TOPOLOGY: not relevant

20 (ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:90:

Met Ile Leu Asn Ser Ser Thr Glu Asp Gly Ile Lys Arg Ile Gln Asp
 1 5 10 15
 Asp Cys Pro Lys Ala Gly Arg His Asn Tyr Ile Phe Val Met Ile Pro
 25 20 25 30
 Thr Leu Tyr Ser Ile Ile Phe Val Val Gly Ile Phe Gly Asn Ser Leu
 35 40 45
 Val Val Ile Val Ile Tyr Phe Tyr Met Lys Leu Lys Thr Val Ala Ser
 50 55 60
 30 Val Phe Leu Leu Asn Leu Ala Leu Ala Asp Leu Cys Phe Leu Leu Thr
 65 70 75 80
 Leu Pro Leu Trp Ala Val Tyr Thr Ala Met Glu Tyr Arg Trp Pro Phe

- 73 -

	85	90	95
	Gly Asn Tyr Leu Cys Lys Ile Ala Ser	Ala Ser Val Ser	Phe Asn Leu
	100	105	110
5	Tyr Ala Ser Val Phe Leu Leu Thr Cys Leu Ser Ile Asp Arg Tyr Leu		
	115	120	125
	Ala Ile Val His Pro Met Lys Ser Arg Leu Arg Arg Thr Met Leu Val		
	130	135	140
	Ala Lys Val Thr Cys Ile Ile Ile Trp Leu Leu Ala Gly Leu Ala Ser		
	145	150	155 160
10	Leu Pro Ala Ile Ile His Arg Asn Val Phe Phe Ile Glu Asn Thr Asn		
	165	170	175
	Ile Thr Val Cys Ala Phe His Tyr Glu Ser Gln Asn Ser Thr Leu Pro		
	180	185	190
15	Ile Gly Leu Gly Leu Thr Lys Asn Ile Leu Gly Phe Leu Phe Pro Phe		
	195	200	205
	Leu Ile Ile Leu Thr Ser Tyr Thr Leu Ile Trp Lys Ala Leu Lys Lys		
	210	215	220
	Ala Tyr Glu Ile Gln Lys Asn Lys Pro Arg Asn Asp Asp Ile Lys Lys		
	225	230	235 240
20	Ile Ile Met Ala Ile Val Leu Phe Phe Phe Phe Ser Trp Ile Pro His		
	245	250	255
	Gln Ile Phe Thr Phe Leu Asp Val Leu Ile Gln Leu Gly Ile Ile Arg		
	260	265	270
25	Asp Cys Arg Ile Ala Asp Ile Val Asp Thr Ala Met Pro Ile Thr Ile		
	275	280	285
	Cys Ile Ala Tyr Phe Asn Asn Cys Leu Asn Pro Leu Phe Tyr Gly Phe		
	290	295	300
	Leu Gly Lys Lys Phe Lys Arg Tyr Phe Leu Gln Leu Leu Lys Tyr Ile		
	305	310	315 320
30	Pro Pro Lys Ala Lys Ser His Ser Asn Leu Ser Thr Lys Met Ser Thr		
	325	330	335
	Leu Ser Tyr Arg Pro Ser Asp Asn Val Ser Ser Ser Thr Lys Lys Pro		
	340	345	350
35	Ala Pro Cys Phe Glu Val Glu		
	355		

(92) INFORMATION FOR SEQ ID NO:91:

- 74 -

- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 35 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:91:

CCAAGAAATG ATGATATTAA AAAGATAATT ATGGC

35

(93) INFORMATION FOR SEQ ID NO:92:

- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 31 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:92:

CTCCTTCGGT CCTCCTATCG TTGTCAGAAG T

31

(94) INFORMATION FOR SEQ ID NO:93:

- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 1080 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:93:

ATGATTCTCA ACTCTTCTAC TGAAGATGGT ATTAAAAGAA TCCAAGATGA TTGTCCCAAA 60

GCTGGAAGGC ATAATTACAT ATTTGTCATG ATTCCTACTT TATACAGTAT CATCTTTGTG 120

GTGGGAATAT TTGGAAACAG CTTGGTGGTG ATAGTCATTT ACTTTTATAT GAAGCTGAAG 180

ACTGTGGCCA GTGTTTTTCT TTTGAATTTA GCACTGGCTG ACTTATGCTT TTTACTGACT 240

TTGCCACTAT GGGCTGTCTA CACAGCTATG GAATACCGCT GGCCCTTTGG CAATTACCTA 300

TGTAAGATTG CTTGAGCCAG CGTCAGTTTC GCCCTGTACG CTAGTGTGTT TCTACTCAGC 360

TGTCTCAGCA TTGATCGATA CCTGGCTATT GTTCACCCAA TGAAGTCCCG CCTTCGACGC 420

- 75 -

ACAATGCTTG TAGCCAAAGT CACCTGCATC ATCATTGGC TGCTGGCAGG CTTGGCCAGT 480
 TTGCCAGCTA TAATCCATCG AAATGTATTT TTCATTGAGA ACACCAATAT TACAGTTTGT 540
 GCTTTCCATT ATGAGTCCCA AAATTCAACC CTTCGATAG GGCTGGGCCT GACCAAAAAT 600
 ATACTGGGTT TCCTGTTTCC TTTCTGATC ATTCTTACAA GTTATACTCT TATTTGGAAG 660
 5 GCCCTAAAGA AGGCTTATGA AATTCAGAAG AACAAACCAA GAAATGATGA TATTTTAAAG 720
 ATAATTATGG CAATTGTGCT TTTCTTTTTC TTTTCCTGGA TTCCCCACCA AATATTCCT 780
 TTTCTGGATG TATTGATTCA ACTAGGCATC ATACGTGACT GTAGAATTGC AGATATTGTG 840
 GACACGGCCA TGCCTATCAC CATTGTGATA GCTTATTTTA ACAATGCTT GAATCCTCTT 900
 TTTTATGGCT TTCTGGGGAA AAAATTTAAA AGATATTTTC TCCAGCTTCT AAAATATATT 960
 10 CCCCCAAAAG CCAATCCCA CTCAAACCTT TCAACAAAA TGAGCAGCT TTCCTACCGC 1020
 CCCTCAGATA ATGTAAGCTC ATCCACCAAG AAGCCTGCAC CATGTTTTGA GGTGAGTGA 1080

(95) INFORMATION FOR SEQ ID NO:94:

(i) SEQUENCE CHARACTERISTICS:

- 15 (A) LENGTH: 359 amino acids
 (B) TYPE: amino acid
 (C) STRANDEDNESS:
 (D) TOPOLOGY: not relevant

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:94:

20 Met Ile Leu Asn Ser Ser Thr Glu Asp Gly Ile Lys Arg Ile Gln Asp
 1 5 10 15
 Asp Cys Pro Lys Ala Gly Arg His Asn Tyr Ile Phe Val Met Ile Pro
 20 25 30
 25 Thr Leu Tyr Ser Ile Ile Phe Val Val Gly Ile Phe Gly Asn Ser Leu
 35 40 45
 Val Val Ile Val Ile Tyr Phe Tyr Met Lys Leu Lys Thr Val Ala Ser
 50 55 60
 Val Phe Leu Leu Asn Leu Ala Leu Ala Asp Leu Cys Phe Leu Leu Thr
 65 70 75 80
 30 Leu Pro Leu Trp Ala Val Tyr Thr Ala Met Glu Tyr Arg Trp Pro Phe
 85 90 95
 Gly Asn Tyr Leu Cys Lys Ile Ala Ser Ala Ser Val Ser Phe Ala Leu

- 76 -

	100	105	110
	Tyr Ala Ser Val Phe Leu Leu Thr Cys Leu Ser Ile Asp Arg Tyr Leu		
	115	120	125
5	Ala Ile Val His Pro Met Lys Ser Arg Leu Arg Arg Thr Met Leu Val		
	130	135	140
	Ala Lys Val Thr Cys Ile Ile Ile Trp Leu Leu Ala Gly Leu Ala Ser		
	145	150	155 160
	Leu Pro Ala Ile Ile His Arg Asn Val Phe Phe Ile Glu Asn Thr Asn		
		165	170 175
10	Ile Thr Val Cys Ala Phe His Tyr Glu Ser Gln Asn Ser Thr Leu Pro		
		180	185 190
	Ile Gly Leu Gly Leu Thr Lys Asn Ile Leu Gly Phe Leu Phe Pro Phe		
		195	200 205
15	Leu Ile Ile Leu Thr Ser Tyr Thr Leu Ile Trp Lys Ala Leu Lys Lys		
		210	215 220
	Ala Tyr Glu Ile Gln Lys Asn Lys Pro Arg Asn Asp Asp Ile Phe Lys		
		225	230 235 240
	Ile Ile Met Ala Ile Val Leu Phe Phe Phe Phe Ser Trp Ile Pro His		
		245	250 255
20	Gln Ile Phe Thr Phe Leu Asp Val Leu Ile Gln Leu Gly Ile Ile Arg		
		260	265 270
	Asp Cys Arg Ile Ala Asp Ile Val Asp Thr Ala Met Pro Ile Thr Ile		
		275	280 285
25	Cys Ile Ala Tyr Phe Asn Asn Cys Leu Asn Pro Leu Phe Tyr Gly Phe		
		290	295 300
	Leu Gly Lys Lys Phe Lys Arg Tyr Phe Leu Gln Leu Leu Lys Tyr Ile		
		305	310 315 320
	Pro Pro Lys Ala Lys Ser His Ser Asn Leu Ser Thr Lys Met Ser Thr		
		325	330 335
30	Leu Ser Tyr Arg Pro Ser Asp Asn Val Ser Ser Ser Thr Lys Lys Pro		
		340	345 350
	Ala Pro Cys Phe Glu Val Glu		
		355	

(97) INFORMATION FOR SEQ ID NO:95:

35

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 26 base pairs

(B) TYPE: nucleic acid

- 77 -

(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(iv) ANTI-SENSE: NO

5 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:95:

CCCAAGCTTC CCCAGGTGTA TTGAT

26

(97) INFORMATION FOR SEQ ID NO:96:

(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 29 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

10

(ii) MOLECULE TYPE: DNA (genomic)

(iv) ANTI-SENSE: YES

15 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:96:

CCTGCAGGCG AACTGACTC TGGCTGAAG

29

(98) INFORMATION FOR SEQ ID NO:97:

(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 42 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

20

(ii) MOLECULE TYPE: DNA (genomic)

(iv) ANTI-SENSE: NO

25 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:97:

CTGTACGCTA GTGTGTTTCT ACTCACGTGT CTCAGCATTG AT

42

(99) INFORMATION FOR SEQ ID NO:98:

(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 26 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

30

(ii) MOLECULE TYPE: DNA (genomic)

- 78 -

(iv) ANTI-SENSE: YES

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:98:

GTTGGATCCA CATAATGCAT TTTCTC

26

(100) INFORMATION FOR SEQ ID NO:99:

- 5 (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 1080 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

10 (ii) MOLECULE TYPE: DNA (genomic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:99:

ATGATTCTCA ACTCTTCTAC TGAAGATGGT ATTAAAAGAA TCCAAGATGA TTGTCCCAAA 60
GCTGGAAGGC ATAATTACAT ATTTGTCATG ATTCCTACTT TATACAGTAT CATCTTTGTG 120
GTGGGAATAT TTGGAACAG CTTGGTGGTG ATAGTCATTT ACTTTTATAT GAAGCTGAAG 180
15 ACTGTGGCCA GTGTTTTTCT TTTGAATTTA GCACTGGCTG ACTTATGCTT TTTACTGACT 240
TTGCCACTAT GGGCTGTCTA CACAGCTATG GAATACCGCT GGCCCTTTGG CAATTACCTA 300
TGTAAGATTG CTTAGCCAG CGTCAGTTTC AACCTGTACG CTAGTGTGTT TCTACTCAGC 360
TGTCTCAGCA TTGATCGATA CCTGGCTATT GTTCACCCAA TGAAGTCCCG CCTTCGACGC 420
ACAATGCTTG TAGCCAAAGT CACCTGCATC ATCATTGGC TGCTGGCAGG CTTGGCCAGT 480
20 TTGCCAGCTA TAATCCATCG AAATGTATTT TTCATTGAGA ACACCAATAT TACAGTTTGT 540
GCTTTCCATT ATGAGTCCCA AAATTCAACC CTTCCGATAG GGCTGGGCCT GACCAAAAAT 600
ATACTGGGTT TCCTGTTTCC TTTTCTGATC ATTCTTACAA GTTATTTTGG AATTCGAAAA 660
CACTTACTGA AGACGAATAG CTATGGGAAG AACAGGATAA CCCGTGACCA AGTTAAGAAG 720
ATAATTATGG CAATTGTGCT TTTCTTTTTC TTTTCTGGA TTCCCCACCA AATATTCACT 780
25 TTTCTGGATG TATTGATTCA ACTAGGCATC ATACGTGACT GTAGAATTGC AGATATTGTG 840
GACACGGCCA TGCCTATCAC CATTTGTATA GCTTATTTTA ACAATTGCCT GAATCCTCTT 900
TTTTATGGCT TTCTGGGGAA AAAATTTAAA AGATATTTTC TCCAGCTTCT AAAATATATT 960
CCCCAAAAG CCAATCCCA CTCAAACCTT TCAACAAAA TGAGCAGCT TTCCTACCGC 1020
CCCTCAGATA ATGTAAGCTC ATCCACCAAG AAGCCTGCAC CATGTTTTGA GGTGAGTGA 1080

- 79 -

(101) INFORMATION FOR SEQ ID NO:100:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 359 amino acids
 (B) TYPE: amino acid
 (C) STRANDEDNESS:
 (D) TOPOLOGY: not relevant

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:100:

10 Met Ile Leu Asn Ser Ser Thr Glu Asp Gly Ile Lys Arg Ile Gln Asp
 1 5 10 15
 Asp Cys Pro Lys Ala Gly Arg His Asn Tyr Ile Phe Val Met Ile Pro
 20 25 30
 Thr Leu Tyr Ser Ile Ile Phe Val Val Gly Ile Phe Gly Asn Ser Leu
 35 40 45
 15 Val Val Ile Val Ile Tyr Phe Tyr Met Lys Leu Lys Thr Val Ala Ser
 50 55 60
 Val Phe Leu Leu Asn Leu Ala Leu Ala Asp Leu Cys Phe Leu Leu Thr
 65 70 75 80
 20 Leu Pro Leu Trp Ala Val Tyr Thr Ala Met Glu Tyr Arg Trp Pro Phe
 85 90 95
 Gly Asn Tyr Leu Cys Lys Ile Ala Ser Ala Ser Val Ser Phe Asn Leu
 100 105 110
 Tyr Ala Ser Val Phe Leu Leu Thr Cys Leu Ser Ile Asp Arg Tyr Leu
 115 120 125
 25 Ala Ile Val His Pro Met Lys Ser Arg Leu Arg Arg Thr Met Leu Val
 130 135 140
 Ala Lys Val Thr Cys Ile Ile Ile Trp Leu Leu Ala Gly Leu Ala Ser
 145 150 155 160
 30 Leu Pro Ala Ile Ile His Arg Asn Val Phe Phe Ile Glu Asn Thr Asn
 165 170 175
 Ile Thr Val Cys Ala Phe His Tyr Glu Ser Gln Asn Ser Thr Leu Pro
 180 185 190
 Ile Gly Leu Gly Leu Thr Lys Asn Ile Leu Gly Phe Leu Phe Pro Phe
 195 200 205
 35 Leu Ile Ile Leu Thr Ser Tyr Phe Gly Ile Arg Lys His Leu Leu Lys
 210 215 220

- 80 -

Thr Asn Ser Tyr Gly Lys Asn Arg Ile Thr Arg Asp Gln Val Lys Lys
 225 230 235 240
 Ile Ile Met Ala Ile Val Leu Phe Phe Phe Phe Ser Trp Ile Pro His
 245 250 255
 5 Gln Ile Phe Thr Phe Leu Asp Val Leu Ile Gln Leu Gly Ile Ile Arg
 260 265 270
 Asp Cys Arg Ile Ala Asp Ile Val Asp Thr Ala Met Pro Ile Thr Ile
 275 280 285
 10 Cys Ile Ala Tyr Phe Asn Asn Cys Leu Asn Pro Leu Phe Tyr Gly Phe
 290 295 300
 Leu Gly Lys Lys Phe Lys Arg Tyr Phe Leu Gln Leu Leu Lys Tyr Ile
 305 310 315 320
 Pro Pro Lys Ala Lys Ser His Ser Asn Leu Ser Thr Lys Met Ser Thr
 325 330 335
 15 Leu Ser Tyr Arg Pro Ser Asp Asn Val Ser Ser Ser Thr Lys Lys Pro
 340 345 350
 Ala Pro Cys Phe Glu Val Glu
 355

(102) INFORMATION FOR SEQ ID NO:101:

- 20 (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 37 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear
 25 (ii) MOLECULE TYPE: DNA (genomic)
 (iv) ANTI-SENSE: YES

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:101:

TCCGAATTCC AAAATAACTT GTAAGAATGA TCAGAAA

37

(103) INFORMATION FOR SEQ ID NO:102:

- 30 (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 33 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear
 35 (ii) MOLECULE TYPE: DNA (genomic)
 (iv) ANTI-SENSE: NO

- 81 -

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:102:

AGATCTTAAG AAGATAATTA TGGCAATTGT GCT

33

(104) INFORMATION FOR SEQ ID NO:103:

(i) SEQUENCE CHARACTERISTICS:

- 5 (A) LENGTH: 62 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

10 (iv) ANTI-SENSE: NO

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:103:

AATTGAAAA CACTTACTGA AGACGAATAG CTATGGGAAG AACAGGATAA CCCGTGACCA

60

AG

62

(105) INFORMATION FOR SEQ ID NO:104:

(i) SEQUENCE CHARACTERISTICS:

- 15 (A) LENGTH: 62 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

20 (ii) MOLECULE TYPE: DNA (genomic)

(iv) ANTI-SENSE: YES

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:104:

TTAACTTGGT CACGGGTAT CTGTTCTTC CCATAGCTAT TCGTCTTCAG TAAGTGT

60

CG

62

25 (106) INFORMATION FOR SEQ ID NO:105:

(i) SEQUENCE CHARACTERISTICS:

- 30 (A) LENGTH: 1083 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:105:

- 82 -

ATGATTCTCA ACTCTTCTAC TGAAGATGGT ATTAAAAGAA TCCAAGATGA TTGTCCCAAA 60
 GCTGGAAGGC ATAATTACAT ATTTGTCATG ATTCCTACTT TATACAGTAT CATCTTTGTG 120
 GTGGGAATAT TTGGAACAG CTTGGTGGTG ATAGTCATTT ACTTTTATAT GAAGCTGAAG 180
 ACTGTGGCCA GTGTTTTTCT TTTGAATTTA GCCTGGCTG ACTTATGCTT TTTACTGACT 240
 5 TTGCCACTAT GGGCTGTCTA CACAGCTATG GAATACCGCT GGCCCTTTGG CAATTACCTA 300
 TGTAAGATTG CTTAGCCAG CGTCAGTTTC AACCTGTACG CTAGTGTGTT TCTACTCACG 360
 TGTCTCAGCA TTGATCGATA CCTGGCTATT GTTCACCCAA TGAAGTCCCG CCTTCGACGC 420
 ACAATGCTTG TAGCCAAAGT CACCTGCATC ATCATTGGC TGCTGGCAGG CTTGGCCAGT 480
 TTGCCAGCTA TAATCCATCG AAATGTATTT TTCATTGAGA ACACCAATAT TACAGTTTGT 540
 10 GCTTTCCATT ATGAGTCCCA AAATTCAACC CTTCCGATAG GGCTGGGCCT GACCAAAAAT 600
 ATACTGGGTT TCCTGTTTCC TTTTCTGATC ATTCTTACAA GTTATACTCT TATTTGGAAG 660
 GCCCTAAAGA AGGCTTATGA AATTCAGAAG AACAAACCAA GAAATGATGA TATTTTAAAG 720
 ATAATTATGG CAGCAATTGT GCTTTTCTTT TTCTTTTCCT GGATTCCCCA CCAAATATTC 780
 ACTTTTCTGG ATGTATTGAT TCAACTAGGC ATCATACGTG ACTGTAGAAT TGCAGATATT 840
 15 GTGGACACGG CCATGCCTAT CACCATTGT ATAGCTTATT TTAACAATTG CCTGAATCCT 900
 CTTTTTTATG GCTTCTGGG GAAAAAATTT AAAAGATATT TTCTCCAGCT TCTAAAATAT 960
 ATTCCCCCAA AAGCCAAATC CCACTCAAAC CTTTCAACAA AAATGAGCAC GCTTTCCTAC 1020
 CGCCCTCAG ATAATGTAAG CTCATCCACC AAGAAGCCTG CACCATGTTT TGAGGTTGAG 1080
 TGA 1083

20 (107) INFORMATION FOR SEQ ID NO:106:

- (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 360 amino acids
 - (B) TYPE: amino acid
 - (C) STRANDEDNESS:
 - 25 (D) TOPOLOGY: not relevant

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:106:

Met Ile Leu Asn Ser Ser Thr Glu Asp Gly Ile Lys Arg Ile Gln Asp
 1 5 10 15

30 Asp Cys Pro Lys Ala Gly Arg His Asn Tyr Ile Phe Val Met Ile Pro

- 83 -

		20		25		30
		Thr Leu Tyr Ser Ile Ile Phe Val Val Gly Ile Phe Gly Asn Ser Leu				
		35		40		45
5		Val Val Ile Val Ile Tyr Phe Tyr Met Lys Leu Lys Thr Val Ala Ser				
		50		55		60
		Val Phe Leu Leu Asn Leu Ala Leu Ala Asp Leu Cys Phe Leu Leu Thr				
		65		70		75 80
		Leu Pro Leu Trp Ala Val Tyr Thr Ala Met Glu Tyr Arg Trp Pro Phe				
			85	90		95
10		Gly Asn Tyr Leu Cys Lys Ile Ala Ser Ala Ser Val Ser Phe Asn Leu				
			100	105		110
		Tyr Ala Ser Val Phe Leu Leu Thr Cys Leu Ser Ile Asp Arg Tyr Leu				
			115	120		125
15		Ala Ile Val His Pro Met Lys Ser Arg Leu Arg Arg Thr Met Leu Val				
			130	135		140
		Ala Lys Val Thr Cys Ile Ile Ile Trp Leu Leu Ala Gly Leu Ala Ser				
			145	150		155 160
		Leu Pro Ala Ile Ile His Arg Asn Val Phe Phe Ile Glu Asn Thr Asn				
			165	170		175
20		Ile Thr Val Cys Ala Phe His Tyr Glu Ser Gln Asn Ser Thr Leu Pro				
			180	185		190
		Ile Gly Leu Gly Leu Thr Lys Asn Ile Leu Gly Phe Leu Phe Pro Phe				
			195	200		205
25		Leu Ile Ile Leu Thr Ser Tyr Thr Leu Ile Trp Lys Ala Leu Lys Lys				
			210	215		220
		Ala Tyr Glu Ile Gln Lys Asn Lys Pro Arg Asn Asp Asp Ile Phe Lys				
			225	230		235 240
		Ile Ile Met Ala Ala Ile Val Leu Phe Phe Phe Phe Ser Trp Ile Pro				
			245	250		255
30		His Gln Ile Phe Thr Phe Leu Asp Val Leu Ile Gln Leu Gly Ile Ile				
			260	265		270
		Arg Asp* Cys Arg Ile Ala Asp Ile Val Asp Thr Ala Met Pro Ile Thr				
			275	280		285
35		Ile Cys Ile Ala Tyr Phe Asn Asn Cys Leu Asn Pro Leu Phe Tyr Gly				
			290	295		300
		Phe Leu Gly Lys Lys Phe Lys Arg Tyr Phe Leu Gln Leu Leu Lys Tyr				
			305	310		315 320

- 84 -

Ile Pro Pro Lys Ala Lys Ser His Ser Asn Leu Ser Thr Lys Met Ser
325 330 335

Thr Leu Ser Tyr Arg Pro Ser Asp Asn Val Ser Ser Ser Thr Lys Lys
340 345 350

5 Pro Ala Pro Cys Phe Glu Val Glu
355 360

(108) INFORMATION FOR SEQ ID NO:107:

- 10 (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 26 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(iv) ANTI-SENSE: NO

15 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:107:

CCCAAGCTTC CCCAGGTGTA TTTGAT

26

(109) INFORMATION FOR SEQ ID NO:108:

- 20 (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 38 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(iv) ANTI-SENSE: YES

25 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:108:

AAGCACAATT GCTGCATAAT TATCTTAAAA ATATCATC

38

(110) INFORMATION FOR SEQ ID NO:109:

- 30 (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 39 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(iv) ANTI-SENSE: NO

- 85 -

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:109:

AAGATAATTA TGGCAGCAAT TGTGCTTTTC TTTTCTTT

39

(111) INFORMATION FOR SEQ ID NO:110:

(i) SEQUENCE CHARACTERISTICS:

- 5 (A) LENGTH: 26 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic).

10 (iv) ANTI-SENSE: YES

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:110:

GTTGGATCCA CATAATGCAT TTTCTC

26

(112) INFORMATION FOR SEQ ID NO:111:

(i) SEQUENCE CHARACTERISTICS:

- 15 (A) LENGTH: 1344 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic).

20 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:111:

ATGGAGCTGC TAAAGCTGAA CCGGAGCGTG CAGGGAACCG GACCCGGGCC GGGGGCTTCC 60
CTGTGCCGCC CGGGGGCGCC TCTCCTCAAC AGCAGCAGTG TGGGCAACCT CAGCTGCGAG 120
CCCCCTCGCA TTCGCGGAGC CGGGACACGA GAATTGGAGC TGGCCATTAG AATCACTCTT 180
TACGCAGTGA TCTTCCTGAT GAGCGTTGGA GGAAATATGC TCATCATCGT GGTCTCTGGA 240
25 CTGAGCCGCC GCCTGAGGAC TGTACCAAT GCCTTCCTCC TCTCACTGGC AGTCAGCGAC 300
CTCCTGCTGG CTGTGGCTTG CATGCCCTTC ACCCTCCTGC CCAATCTCAT GGGCACATTC 360
ATCTTTGGCA CCGTCATCTG CAAGGCGGTT TCCTACCTCA TGGGGGTGTC TGTGAGTGTG 420
TCCACGCTAA GCCTCGTGGC CATCGCACTG GAGCGATATA GCGCCATCTG CCGACCACTG 480
CAGGCACGAG TGTGGCAGAC GCGCTCCAC GCGGCTCGCG TGATTGTAGC CACGTGGCTG 540
30 CTGTCCGGAC TACTCATGGT GCCCTACCCC GTGTACACTG TCGTGCAACC AGTGGGGCCT 600
CGTGTGCTGC AGTGCGTGCA TCGCTGGCCC AGTGC GCGGG TCCGCCAGAC CTGGTCCGTA 660

- 86 -

CTGCTGCTTC TGCTCTTGTT CTTTCATCCCA GGTGTGGTTA TGGCCGTGGC CTACGGGCTT 720
 ATCTCTCGCG AGCTCTACTT AGGGCTTCGC TTTGACGGCG ACAGTGACAG CGACAGCCAA 780
 AGCAGGGTCC GAAACCAAGG CGGGCTGCCA GGGGCTGTTC ACCAGAACGG GCGTTGCCGG 840
 CCTGAGACTG GCGCGGTTGG CAAAGACAGC GATGGCTGCT ACGTGCAACT TCCACGTTCC 900
 5 CGGCCTGCCC TGGAGCTGAC GGCCTGACG GCTCCTGGGC CGGGATCCGG CTCCCGGCCC 960
 ACCCAGGCCA AGCTGCTGGC TAAGAAGCGC GTGAAACGAA TGTGCTGGT GATCGTTGTG 1020
 CTTTTTTTTC TGTGTTGGTT GCCAGTTTAT AGTGCCAACA CGTGGCGCGC CTTTGATGGC 1080
 CCGGGTGAC ACCGAGCACT CTCGGGTGCT CCTATCTCCT TCATTCACTT GCTGAGCTAC 1140
 GCCTCGGCCT GTGTCAACCC CCTGGTCTAC TGCTTCATGC ACCGTCGCTT TCGCCAGGCC 1200
 10 TGCCTGGAAA CTTGCGCTCG CTGCTGCCCC CGGCCTCCAC GAGCTCGCCC CAGGGCTCTT 1260
 CCCGATGAGG ACCCTCCAC TCCCTCCATT GCTTCGCTGT CCAGGCTTAG CTACACCACC 1320
 ATCAGCACAC TGGGCCCTGG CTGA 1344

(113) INFORMATION FOR SEQ ID NO:112:

- (i) SEQUENCE CHARACTERISTICS:
 15 (A) LENGTH: 447 amino acids
 (B) TYPE: amino acid
 (C) STRANDEDNESS:
 (D) TOPOLOGY: not relevant

(ii) MOLECULE TYPE: protein

20 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:112:

Met Glu Leu Leu Lys Leu Asn Arg Ser Val Gln Gly Thr Gly Pro Gly
 1 5 10 15
 Pro Gly Ala Ser Leu Cys Arg Pro Gly Ala Pro Leu Leu Asn Ser Ser
 20 25 30
 25 Ser Val Gly Asn Leu Ser Cys Glu Pro Pro Arg Ile Arg Gly Ala Gly
 35 40 45
 Thr Arg Glu Leu Glu Leu Ala Ile Arg Ile Thr Leu Tyr Ala Val Ile
 50 55 60
 Phe Leu Met Ser Val Gly Gly Asn Met Leu Ile Ile Val Val Leu Gly
 30 65 70 75 80
 Leu Ser Arg Arg Leu Arg Thr Val Thr Asn Ala Phe Leu Leu Ser Leu
 85 90 95

- 87 -

Ala Val Ser Asp Leu Leu Leu Ala Val Ala Cys Met Pro Phe Thr Leu
100 105 110

Leu Pro Asn Leu Met Gly Thr Phe Ile Phe Gly Thr Val Ile Cys Lys
115 120 125

5 Ala Val Ser Tyr Leu Met Gly Val Ser Val Ser Val Ser Thr Leu Ser
130 135 140

Leu Val Ala Ile Ala Leu Glu Arg Tyr Ser Ala Ile Cys Arg Pro Leu
145 150 155 160

10 Gln Ala Arg Val Trp Gln Thr Arg Ser His Ala Ala Arg Val Ile Val
165 170 175

Ala Thr Trp Leu Leu Ser Gly Leu Leu Met Val Pro Tyr Pro Val Tyr
180 185 190

Thr Val Val Gln Pro Val Gly Pro Arg Val Leu Gln Cys Val His Arg
195 200 205

15 Trp Pro Ser Ala Arg Val Arg Gln Thr Trp Ser Val Leu Leu Leu Leu
210 215 220

Leu Leu Phe Phe Ile Pro Gly Val Val Met Ala Val Ala Tyr Gly Leu
225 230 235 240

20 Ile Ser Arg Glu Leu Tyr Leu Gly Leu Arg Phe Asp Gly Asp Ser Asp
245 250 255

Ser Asp Ser Gln Ser Arg Val Arg Asn Gln Gly Gly Leu Pro Gly Ala
260 265 270

Val His Gln Asn Gly Arg Cys Arg Pro Glu Thr Gly Ala Val Gly Lys
275 280 285

25 Asp Ser Asp Gly Cys Tyr Val Gln Leu Pro Arg Ser Arg Pro Ala Leu
290 295 300

Glu Leu Thr Ala Leu Thr Ala Pro Gly Pro Gly Ser Gly Ser Arg Pro
305 310 315 320

30 Thr Gln Ala Lys Leu Leu Ala Lys Lys Arg Val Lys Arg Met Leu Leu
325 330 335

Val Ile Val Val Leu Phe Phe Leu Cys Trp Leu Pro Val Tyr Ser Ala
340 345 350

Asn Thr Trp Arg Ala Phe Asp Gly Pro Gly Ala His Arg Ala Leu Ser
355 360 365

35 Val Ala Pro Ile Ser Phe Ile His Leu Leu Ser Tyr Ala Ser Ala Cys
370 375 380

Val Asn Pro Leu Val Tyr Cys Phe Met His Arg Arg Phe Arg Gln Ala

385					390					395					400
Cys	Leu	Glu	Thr	Cys	Ala	Arg	Cys	Cys	Pro	Arg	Pro	Pro	Arg	Ala	Arg
				405					410					415	
Pro	Arg	Ala	Leu	Pro	Asp	Glu	Asp	Pro	Pro	Thr	Pro	Ser	Ile	Ala	Ser
			420					425					430		
Leu	Ser	Arg	Leu	Ser	Tyr	Thr	Thr	Ile	Ser	Thr	Leu	Gly	Pro	Gly	
			435				440					445			

(i) SEQUENCE CHARACTERISTICS:

10. (A) LENGTH: 34 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

15 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:113:

34

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 33 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: not relevant

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:114:

35

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 33 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(iv) ANTI-SENSE: NO

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:115:

- 89 -

ATGGAGAAAA GAATCAAAAG AATGTTCTAT ATA

33

(117) INFORMATION FOR SEQ ID NO:116:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 33 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(iv) ANTI-SENSE: YES

10 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:116:

TATATAGAAC ATTCTTTTGA TTCTTTTCTC CAT

33

(118) INFORMATION FOR SEQ ID NO:117:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 30 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(iv) ANTI-SENSE: NO

20 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:117:

CGCTCTCTGG CCTTGAAGCG CACGCTCAGC

30

(119) INFORMATION FOR SEQ ID NO:118:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 30 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(iv) ANTI-SENSE: YES

30 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:118:

GCTGAGCGTG CGCTTCAAGG CCAGAGAGCG

30

(120) INFORMATION FOR SEQ ID NO:119:

- 90 -

(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 30 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
5 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(iv) ANTI-SENSE: NO

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:119:

CCCAGGAAAA AGGTGAAAGT CAAAGTTTTC

30

10 (121) INFORMATION FOR SEQ ID NO:120:

(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 30 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
15 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(iv) ANTI-SENSE: YES

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:120:

GAAAACTTTG ACTTTCACCT TTTTCCTGGG

30

20 (122) INFORMATION FOR SEQ ID NO:121:

(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 27 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
25 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(iv) ANTI-SENSE: NO

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:121:

GGGGCGCGGG TGAAACGGCT GGTGAGC

27

30 (123) INFORMATION FOR SEQ ID NO:122:

(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 27 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single

- 91 -

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(iv) ANTI-SENSE: YES

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:122:

5 GCTCACCAGC CGTTTCACCC GCGCCCC

27

(124) INFORMATION FOR SEQ ID NO:123:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 30 base pairs

(B) TYPE: nucleic acid

10 (C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(iv) ANTI-SENSE: NO

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:123:

15 CCCCTTGAAA AGCCTAAGAA CTTGGTCATC

30

(125) INFORMATION FOR SEQ ID NO:124:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 30 base pairs

(B) TYPE: nucleic acid

20 (C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(iv) ANTI-SENSE: YES

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:124:

25 GATGACCAAG TTCTTAGGCT TTTCAAGGGG

30

(126) INFORMATION FOR SEQ ID NO:125:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 32 base pairs

(B) TYPE: nucleic acid

30 (C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

- 92 -

(iv) ANTI-SENSE: NO

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:125:

GATCTCTAGA ATGAACAGCA CATGTATTGA AG

32

(127) INFORMATION FOR SEQ ID NO:126:

- 5 (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 35 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

10 (ii) MOLECULE TYPE: DNA (genomic)

(iv) ANTI-SENSE: YES

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:126:

CTAGGGTACC CGCTCAAGGA CCTCTAATTC CATAG

35

(128) INFORMATION FOR SEQ ID NO:127:

- 15 (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 1296 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

20 (ii) MOLECULE TYPE: DNA (genomic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:127:

ATGCAGGCGC TTAACATTAC CCCGGAGCAG TTCTCTCGGC TGCTGCGGGA CCACAACCTG 60
 ACGCGGGAGC AGTTCATCGC TCTGTACCGG CTGCGACCGC TCGTCTACAC CCCAGAGCTG 120
 CCGGGACGCG CCAAGCTGGC CCTCGTGCTC ACCGGCGTGC TCATCTTCGC CCTGGCGCTC 180
 25 TTTGGCAATG CTCTGGTGTT CTACGTGGTG ACCCGCAGCA AGGCCATGCG CACCGTCACC 240
 AACATCTTTA TCTGCTCCTT GCGCTCAGT GACCTGCTCA TCACCTTCTT CTGCATTCCC 300
 GTCACCATGC TCCAGAACAT TTCCGACAAC TGGCTGGGGG GTGCTTTTCAT TTGCAAGATG 360
 GTGCCATTGG TCCAGTCTAC CGCTGTTGTG ACAGAAATGC TCACTATGAC CTGCATTGCT 420
 GTGGAAAGGC ACCAGGGACT TGTGCATCCT TTTAAATGA AGTGGCAATA CACCAACCGA 480

- 93 -

AGGGCTTTCA CAATGCTAGG TGTGGTCTGG CTGGTGGCAG TCATCGTAGG ATCACCCATG 540
 TGGCACGTGC AACAACTTGA GATCAAATAT GACTTCCTAT ATGAAAAGGA ACACATCTGC 600
 TGCTTAGAAG AGTGGACCAG CCCTGTGCAC CAGAAGATCT ACACCACCTT CATCCTTGTC 660
 ATCCTCTTCC TCCTGCCTCT TATGGTGATG CTTATTCTGT ACAGTAAAAT TGGTTATGAA 720
 5 CTTTGGATAA AGAAAAGAGT TGGGGATGGT TCAGTGCTTC GAACTATTCA TGGAAAAGAA 780
 ATGTCCAAAA TAGCCAGGAA GAAGAAACGA GCTAAGATTA TGATGGTGAC AGTGGTGGCT 840
 CTCTTTGCTG TGTGCTGGGC ACCATTCCAT GTTGTCCATA TGATGATTGA ATACAGTAAT 900
 TTTGAAAAGG AATATGATGA TGTCACAATC AAGATGATTT TTGCTATCGT GCAAATTATT 960
 GGATTTTCCA ACTCCATCTG TAATCCCATT GTCTATGCAT TTATGAATGA AAACCTTCAA 1020
 10 AAAAAATGTTT TGTCTGCAGT TTGTTATTGC ATAGTAAATA AAACCTTCTC TCCAGCACAA 1080
 AGGCATGGAA ATTCAAGGAAT TACAATGATG CGGAAGAAAG CAAAGTTTTT CCTCAGAGAG 1140
 AATCCAGTGG AGGAAACCAA AGGAGAAGCA TTCAGTGATG GCAACATTGA AGTCAAATTG 1200
 TGTGAACAGA CAGAGGAGAA GAAAAAGCTC AAACGACATC TTGCTCTCTT TAGGTCTGAA 1260
 CTGGCTGAGA ATTCTCCTTT AGACAGTGGG CATTAA 1296

15 (129) INFORMATION FOR SEQ ID NO:128:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 431 amino acids

(B) TYPE: amino acid

(C) STRANDEDNESS:

20 (D) TOPOLOGY: not relevant

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:128:

Met Gln Ala Leu Asn Ile Thr Pro Glu Gln Phe Ser Arg Leu Leu Arg
 1 5 10 15
 25 Asp His Asn Leu Thr Arg Glu Gln Phe Ile Ala Leu Tyr Arg Leu Arg
 20 25 30
 Pro Leu Val Tyr Thr Pro Glu Leu Pro Gly Arg Ala Lys Leu Ala Leu
 35 40 45
 30 Val Leu Thr Gly Val Leu Ile Phe Ala Leu Ala Leu Phe Gly Asn Ala
 50 55 60
 Leu Val Phe Tyr Val Val Thr Arg Ser Lys Ala Met Arg Thr Val Thr
 65 70 75 80

- 94 -

	Asn	Ile	Phe	Ile	Cys	Ser	Leu	Ala	Leu	Ser	Asp	Leu	Leu	Ile	Thr	Phe	85	90	95
	Phe	Cys	Ile	Pro	Val	Thr	Met	Leu	Gln	Asn	Ile	Ser	Asp	Asn	Trp	Leu	100	105	110
5	Gly	Gly	Ala	Phe	Ile	Cys	Lys	Met	Val	Pro	Phe	Val	Gln	Ser	Thr	Ala	115	120	125
	Val	Val	Thr	Glu	Met	Leu	Thr	Met	Thr	Cys	Ile	Ala	Val	Glu	Arg	His	130	135	140
10	Gln	Gly	Leu	Val	His	Pro	Phe	Lys	Met	Lys	Trp	Gln	Tyr	Thr	Asn	Arg	145	150	155
	Arg	Ala	Phe	Thr	Met	Leu	Gly	Val	Val	Trp	Leu	Val	Ala	Val	Ile	Val	165	170	175
	Gly	Ser	Pro	Met	Trp	His	Val	Gln	Gln	Leu	Glu	Ile	Lys	Tyr	Asp	Phe	180	185	190
15	Leu	Tyr	Glu	Lys	Glu	His	Ile	Cys	Cys	Leu	Glu	Glu	Trp	Thr	Ser	Pro	195	200	205
	Val	His	Gln	Lys	Ile	Tyr	Thr	Thr	Phe	Ile	Leu	Val	Ile	Leu	Phe	Leu	210	215	220
20	Leu	Pro	Leu	Met	Val	Met	Leu	Ile	Leu	Tyr	Ser	Lys	Ile	Gly	Tyr	Glu	225	230	235
	Leu	Trp	Ile	Lys	Lys	Arg	Val	Gly	Asp	Gly	Ser	Val	Leu	Arg	Thr	Ile	245	250	255
	His	Gly	Lys	Glu	Met	Ser	Lys	Ile	Ala	Arg	Lys	Lys	Lys	Arg	Ala	Lys	260	265	270
25	Ile	Met	Met	Val	Thr	Val	Val	Ala	Leu	Phe	Ala	Val	Cys	Trp	Ala	Pro	275	280	285
	Phe	His	Val	Val	His	Met	Met	Ile	Glu	Tyr	Ser	Asn	Phe	Glu	Lys	Glu	290	295	300
30	Tyr	Asp	Asp	Val	Thr	Ile	Lys	Met	Ile	Phe	Ala	Ile	Val	Gln	Ile	Ile	305	310	315
	Gly	Phe	Ser	Asn	Ser	Ile	Cys	Asn	Pro	Ile	Val	Tyr	Ala	Phe	Met	Asn	325	330	335
	Glu	Asn	Phe	Lys	Lys	Asn	Val	Leu	Ser	Ala	Val	Cys	Tyr	Cys	Ile	Val	340	345	350
35	Asn	Lys	Thr	Phe	Ser	Pro	Ala	Gln	Arg	His	Gly	Asn	Ser	Gly	Ile	Thr	355	360	365
	Met	Met	Arg	Lys	Lys	Ala	Lys	Phe	Ser	Leu	Arg	Glu	Asn	Pro	Val	Glu			

- 95 -

370 375 380

Glu Thr Lys Gly Glu Ala Phe Ser Asp Gly Asn Ile Glu Val Lys Leu
385 390 395 400

Cys Glu Gln Thr Glu Glu Lys Lys Lys Leu Lys Arg His Leu Ala Leu
5 405 410 415

Phe Arg Ser Glu Leu Ala Glu Asn Ser Pro Leu Asp Ser Gly His
420 425 430

(130) INFORMATION FOR SEQ ID NO:129:

10 (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 2040 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

15 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:129:

ATGGGCAGCC CCTGGAACGG CAGCGACGGC CCCGAGGGGG CGCGGGAGCC GCCGTGGCCC
60

GCGCTGCCGC CTTGCGACGA GCGCCGCTGC TCGCCCTTTC CCCTGGGGGC GCTGGTGCCG
120

20 GTGACCGCTG TGTGCCTGTG CCTGTTCGTC GTCGGGGTGA GCGGCAACGT GGTGACCGTG
180

ATGCTGATCG GCGCTACCG GGACATGCGG ACCACCACCA ACTTGATCCT GGGCAGCATG
240

GCCGTGTCCG ACCTACTCAT CCTGCTCGGG CTGCCGTTCT ACCTGTACCG CCTCTGGCGC
25 300

TCGCGGCCCT GGGTGTTCGG GCCGCTGCTC TGCCGCCTGT CCCTCTACGT GGGCGAGGGC
360

TGCACCTACG CCACGCTGCT GCACATGACC GCGCTCAGCG TCGAGCGCTA CCTGGCCATC
30 420

TGCCGCCCCG TCCGCGCCCC CGTCTTGGTC ACCCGGCGCC GCGTCCGCGC GCTCATCGCT
480

GTGCTCTGGG CCGTGGCGCT GCTCTCTGCC GGTCCCTTCT TGTTCCTGGT GGGCGTCGAG
35 540

CAGGACCCCG GCATCTCCGT AGTCCCGGGC CTCAATGGCA CCGCGCGGAT CGCCTCCTCG
600

40 CCTCTCGCCT CGTCGCGGCC TCTCTGGCTC TCGCGGGCGC CACCGCCGTC CCCGCCGTCG

- 96 -

660
GGGCCCCGAGA CCGCGGAGGC CGCGGCGCTG TTCAGCCGCG AATGCCGGCC GAGCCCCGCG
720
5 CAGCTGGGCG CGCTGCGTGT CATGCTGTGG GTCACCACCG CCTACTTCTT CTTGCCCTTT
780
CTGTGCCTCA GCATCCTCTA CGGGCTCATC GGGCGGGAGC TGTGGAGCAG CCGGCGGCCG
10 840
CTGCGAGGCC CGGCCGCTC GGGGCGGGAG AGAGGCCACC GGCAGACCAA ACGCGTCCTG
900
15 CGTAAGTGGA GCCGCCGTGG TTCCAAAGAC GCCTGCCTGC AGTCCGCCCC GCCGGGGACC
960
GCGCAAACGC TGGGTCCCCT TCCCCTGCTC GCCCAGCTCT GGGCGCCGCT TCCAGCTCCC
1020
20 TTTCTATTTT CGATTCCAGC CTCCACCCGC CGGTACTTCC CATCCCCCGA GAAAACCATG
1080
TCCTGTCCCC CAGGAGCTCT GGGGGACCCC AGGGCGCTTT GAGGGTGGA TCCCCGGATC
25 1140
CGATTAGTA ACCAGCAGTG CTTTTCCAGA GCCTCTGAGA CCAGAAAGGA GAGTTGGTAA
1200
30 TTCCTAATCC AACCACCTGT TAGATGCCAC AAATGAGGAG TCCTCACAGT GCTCTTGAGA
1260
AGACGAGGGA GATTTTATTA AGCTAAAATT TTTTATTTAA TGTTAAGTGA TGCTGAAGGC
1320
35 TAAAGTAAAC CTTGCTCGTA TCAAAAAGTA AAGATTGTGC AGACCTGTTG TAGAATCTTT
1380
TTCAACAGAG AACAGAAAAC TTGTCTCCGA AGTGGGTTTG TGGAAGGAAG CCTGCCAAGG
40 1440
CGGCTTGTTT AGAGAAATTG CTCCTTCTGG TTTATGTCCA GCCTTGATAA CACATATGGG
1500
45 AGCCTACTAT GCAGTTTAA AGCAAGTATC CATGCAGCCT GCAGCCTGGT CATTTTTTCT
1560
GGGGTGAGGA TCTGCCTAGG TAGAAGTTTT CTCTAATTTA TTTTGCTGTT ACTTGTTATT
1620
50 GCAGATGGTT CTTGTGCGG GTGGGGGGTT TATTTGCTTC CCAATGCTTT TGTTAATCCC
1680
GGTGTGTGT CTTATGTTGC AGTGGTGGTG GTTCTGGCAT TTATAATTG CTGGTTGCCC
55 1740

- 97 -

TTCCACGTTG GCAGAATCAT TTACATAAAC ACGGAAGATT CGCGGATGAT GTACTTCTCT
 1800
 5 CAGTACTTTA ACATCGTCGC TCTGCAACTT TTCTATCTGA GCGCATCTAT CAACCCAATC
 1860
 CTCTACAACC TCATTTCAAA GAAGTACAGA GCGGCGGCCT TTAACTGCT GCTCGCAAGG
 1920
 10 AAGTCCAGGC CGAGAGGCTT CCACAGAAGC AGGGACACTG CGGGGGAAGT TGCAGGGGAC
 1980
 ACTGGAGGAG ACACGGTGGG CTACACCGAG ACAAGCGCTA ACGTGAAGAC GATGGGATAA
 2040
 15
 (131) INFORMATION FOR SEQ ID NO:130:
 (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 412 amino acids
 (B) TYPE: amino acid
 20 (C) STRANDEDNESS:
 (D) TOPOLOGY: not relevant
 (ii) MOLECULE TYPE: protein
 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:130:
 Met Gly Ser Pro Trp Asn Gly Ser Asp Gly Pro Glu Gly Ala Arg Glu
 25 1 5 10 15
 Pro Pro Trp Pro Ala Leu Pro Pro Cys Asp Glu Arg Arg Cys Ser Pro
 20 25 30
 Phe Pro Leu Gly Ala Leu Val Pro Val Thr Ala Val Cys Leu Cys Leu
 35 40 45
 30 Phe Val Val Gly Val Ser Gly Asn Val Val Thr Val Met Leu Ile Gly
 50 55 60
 Arg Tyr Arg Asp Met Arg Thr Thr Thr Asn Leu Tyr Leu Gly Ser Met
 65 70 75 80
 35 Ala Val Ser Asp Leu Leu Ile Leu Leu Gly Leu Pro Phe Asp Leu Tyr
 85 90 95
 Arg Leu Trp Arg Ser Arg Pro Trp Val Phe Gly Pro Leu Leu Cys Arg
 100 105 110
 Leu Ser Leu Tyr Val Gly Glu Gly Cys Thr Tyr Ala Thr Leu Leu His
 115 120 125
 40 Met Thr Ala Leu Ser Val Glu Arg Tyr Leu Ala Ile Cys Arg Pro Leu
 130 135 140

- 98 -

	Arg	Ala	Arg	Val	Leu	Val	Thr	Arg	Arg	Arg	Val	Arg	Ala	Leu	Ile	Ala	
	145						150					155				160	
	Val	Leu	Trp	Ala	Val	Ala	Leu	Leu	Ser	Ala	Gly	Pro	Phe	Leu	Phe	Leu	
					165					170					175		
5	Val	Gly	Val	Glu	Gln	Asp	Pro	Gly	Ile	Ser	Val	Val	Pro	Gly	Leu	Asn	
				180					185					190			
	Gly	Thr	Ala	Arg	Ile	Ala	Ser	Ser	Pro	Leu	Ala	Ser	Ser	Pro	Pro	Leu	
			195					200					205				
10	Trp	Leu	Ser	Arg	Ala	Pro	Pro	Ser	Pro	Pro	Ser	Gly	Pro	Glu	Thr		
	210					215					220						
	Ala	Glu	Ala	Ala	Ala	Leu	Phe	Ser	Arg	Glu	Cys	Arg	Pro	Ser	Pro	Ala	
	225					230					235					240	
	Gln	Leu	Gly	Ala	Leu	Arg	Val	Met	Leu	Trp	Val	Thr	Thr	Ala	Tyr	Phe	
					245					250					255		
15	Phe	Leu	Pro	Phe	Leu	Cys	Leu	Ser	Ile	Leu	Tyr	Gly	Leu	Ile	Gly	Arg	
				260					265					270			
	Glu	Leu	Trp	Ser	Ser	Arg	Arg	Pro	Leu	Arg	Gly	Pro	Ala	Ala	Ser	Gly	
			275					280					285				
20	Arg	Glu	Arg	Gly	His	Arg	Gln	Thr	Lys	Arg	Val	Leu	Leu	Val	Val	Val	
	290						295					300					
	Leu	Ala	Phe	Ile	Ile	Cys	Trp	Leu	Pro	Phe	His	Val	Gly	Arg	Ile	Ile	
	305					310					315					320	
	Tyr	Ile	Asn	Thr	Glu	Asp	Ser	Arg	Met	Met	Tyr	Phe	Ser	Gln	Tyr	Phe	
					325					330					335		
25	Asn	Ile	Val	Ala	Leu	Gln	Leu	Phe	Tyr	Leu	Ser	Ala	Ser	Ile	Asn	Pro	
				340					345					350			
	Ile	Leu	Tyr	Asn	Leu	Ile	Ser	Lys	Lys	Tyr	Arg	Ala	Ala	Ala	Phe	Lys	
			355					360					365				
30	Leu	Leu	Leu	Ala	Arg	Lys	Ser	Arg	Pro	Arg	Gly	Phe	His	Arg	Ser	Arg	
	370						375					380					
	Asp	Thr	Ala	Gly	Glu	Val	Ala	Gly	Asp	Thr	Gly	Gly	Asp	Thr	Val	Gly	
	385					390					395					400	
	Tyr	Thr	Glu	Thr	Ser	Ala	Asn	Val	Lys	Thr	Met	Gly					
					405					410							

35 (132) INFORMATION FOR SEQ ID NO:131:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 1344 base pairs

- 99 -

(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

5 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:131:

ATGGAGCTGC TAAAGCTGAA CCGGAGCGTG CAGGGAACCG GACCCGGGCC GGGGGCTTCC
60

CTGTGCCGCC CGGGGGCGCC TCTCCTCAAC AGCAGCAGTG TGGGCAACCT CAGCTGCGAG
120

10 CCCCCTCGCA TTCGCGGAGC CGGGACACGA GAATTGGAGC TGGCCATTAG AATCACTCTT
180

TACGCAGTGA TCTTCCTGAT GAGCGTTGGA GGAAATATGC TCATCATCGT GGTCTGGGA
240

15 CTGAGCCGCC GCCTGAGGAC TGTCACCAAT GCCTTCCTCC TCTCACTGGC AGTCAGCGAC
300

CTCCTGCTGG CTGTGGCTTG CATGCCCTTC ACCCTCCTGC CCAATCTCAT GGGCACATTC
360

ATCTTTGGCA CCGTCATCTG CAAGGCGGTT TCCTACCTCA TGGGGGTGTC TGTGAGTGTG
420

20 TCCACGCTAA GCCTCGTGGC CATCGCACTG GAGCGATATA GCGCCATCTG CCGACCACTG
480

CAGGCACGAG TGTGGCAGAC GCGCTCCAC GCGGCTCGCG TGATTGTAGC CACGTGGCTG
540

25 CTGTCEGGAC TACTCATGGT GCCCTACCCC GTGTACACTG TCGTGCAACC AGTGGGGCCT
600

CGTGTGCTGC AGTGCGTGCA TCGCTGGCCC AGTGCGCGGG TCCGCCAGAC CTGGTCCGTA
660

CTGCTGCTTC TGCTCTTGTT CTTCAATCCA GGTGTGGTTA TGGCCGTGGC CTACGGGCTT
720

30 ATCTCTCGCG AGCTCTACTT AGGGCTTCGC TTTGACGGCG ACAGTGACAG CGACAGCCAA
780

AGCAGGGTCC GAAACCAAGG CGGGCTGCCA GGGGCTGTTC ACCAGAACGG GCGTTGCCGG
840

35 CCTGAGACTG GCGCGGTTGG CAAAGACAGC GATGGCTGCT ACGTGCAACT TCCACGTTC
900

CGGCCTGCCC TGGAGCTGAC GCGCTGACG GCTCCTGGGC CGGGATCCGG CTCCCGCCCC

- 100 -

960

ACCCAGGCCA AGCTGCTGGC TAAGAAGCGC GTGAAACGAA TGTGCTGGT GATCGTTGTG
1020

5 CTTTTTTTTC TGTGTTGGTT GCCAGTTTAT AGTGCCAACA CGTGGCGCGC CTTTGATGGC
1080

CCGGGTGCAC ACCGAGCACT CTCGGGTGCT CCTATCTCCT TCATTCACCT GCTGAGCTAC
1140

GCCTCGGCCT GTGTCAACCC CCTGGTCTAC TGCTTCATGC ACCGTCGCTT TCGCCAGGCC
1200

10 TGCCTGGAAA CTTGCGCTCG CTGCTGCCCC CGGCCTCCAC GAGCTCGCCC CAGGGCTCTT
1260

CCCGATGAGG ACCCTCCCAC TCCCTCCATT GCTTCGCTGT CCAGGCTTAG CTACACCACC
1320

15 ATCAGCACAC TGGGCCCTGG CTGA
1344

(133) INFORMATION FOR SEQ ID NO:132:

(i) SEQUENCE CHARACTERISTICS:

- 20 (A) LENGTH: 447 amino acids
(B) TYPE: amino acid
(C) STRANDEDNESS:
(D) TOPOLOGY: not relevant

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:132:

25 Met Glu Leu Leu Lys Leu Asn Arg Ser Val Gln Gly Thr Gly Pro Gly
1 5 10 15
Pro Gly Ala Ser Leu Cys Arg Pro Gly Ala Pro Leu Leu Asn Ser Ser
20 25 30
Ser Val Gly Asn Leu Ser Cys Glu Pro Pro Arg Ile Arg Gly Ala Gly
35 40 45
30 Thr Arg Glu Leu Glu Leu Ala Ile Arg Ile Thr Leu Tyr Ala Val Ile
50 55 60
Phe Leu Met Ser Val Gly Gly Asn Met Leu Ile Ile Val Val Leu Gly
65 70 75 80
35 Leu Ser Arg Arg Leu Arg Thr Val Thr Asn Ala Phe Leu Leu Ser Leu
85 90 95
Ala Val Ser Asp Leu Leu Leu Ala Val Ala Cys Met Pro Phe Thr Leu

- 101 -

	100	105	110
	Leu Pro Asn Leu Met Gly Thr Phe Ile Phe Gly Thr Val Ile Cys Lys 115 120 125		
5	Ala Val Ser Tyr Leu Met Gly Val Ser Val Ser Val Ser Thr Leu Ser 130 135 140		
	Leu Val Ala Ile Ala Leu Glu Arg Tyr Ser Ala Ile Cys Arg Pro Leu 145 150 155 160		
	Gln Ala Arg Val Trp Gln Thr Arg Ser His Ala Ala Arg Val Ile Val 165 170 175		
10	Ala Thr Trp Leu Leu Ser Gly Leu Leu Met Val Pro Tyr Pro Val Tyr 180 185 190		
	Thr Val Val Gln Pro Val Gly Pro Arg Val Leu Gln Cys Val His Arg 195 200 205		
15	Trp Pro Ser Ala Arg Val Arg Gln Thr Trp Ser Val Leu Leu Leu Leu 210 215 220		
	Leu Leu Phe Phe Ile Pro Gly Val Val Met Ala Val Ala Tyr Gly Leu 225 230 235 240		
	Ile Ser Arg Glu Leu Tyr Leu Gly Leu Arg Phe Asp Gly Asp Ser Asp 245 250 255		
20	Ser Asp Ser Gln Ser Arg Val Arg Asn Gln Gly Gly Leu Pro Gly Ala 260 265 270		
	Val His Gln Asn Gly Arg Cys Arg Pro Glu Thr Gly Ala Val Gly Lys 275 280 285		
25	Asp Ser Asp Gly Cys Tyr Val Gln Leu Pro Arg Ser Arg Pro Ala Leu 290 295 300		
	Glu Leu Thr Ala Leu Thr Ala Pro Gly Pro Gly Ser Gly Ser Arg Pro 305 310 315 320		
	Thr Gln Ala Lys Leu Leu Ala Lys Lys Arg Val Lys Arg Met Leu Leu 325 330 335		
30	Val Ile Val Val Leu Phe Phe Leu Cys Trp Leu Pro Val Tyr Ser Ala 340 345 350		
	Asn Thr Trp Arg Ala Phe Asp Gly Pro Gly Ala His Arg Ala Leu Ser 355 360 365		
35	Val Ala Pro Ile Ser Phe Ile His Leu Leu Ser Tyr Ala Ser Ala Cys 370 375 380		
	Val Asn Pro Leu Val Tyr Cys Phe Met His Arg Arg Phe Arg Gln Ala 385 390 395 400		

- 102 -

Cys Leu Glu Thr Cys Ala Arg Cys Cys Pro Arg Pro Arg Ala Arg
 405 410 415

Pro Arg Ala Leu Pro Asp Glu Asp Pro Pro Thr Pro Ser Ile Ala Ser
 420 425 430

5 Leu Ser Arg Leu Ser Tyr Thr Thr Ile Ser Thr Leu Gly Pro Gly
 435 440 445

(134) INFORMATION FOR SEQ ID NO:133:

(i) SEQUENCE CHARACTERISTICS:

- 10 (A) LENGTH: 1014 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:133:

15 ATGAACAGCA CATGTATTGA AGAACAGCAT GACCTGGATC ACTATTTGTT TCCCATTTGTT 60
 TACATCTTTG TGATTATAGT CAGCATTCCA GCCAATATTG GATCTCTGTG TGTGTCCTTC 120
 CTGCAAGCAA AGAAGGAAAG TGAAGTAGGA ATTTACCTCT TCAGTTTGTC ACTATCAGAT 180
 TTACTCTATG CATTAAGTCT CCCTTTATGG ATTGATTATA CTTGGAATAA AGACAACTGG 240
 ACTTTCTCTC CTGCCTTGTG CAAAGGGAGT GCTTTTCTCA TGTACATGAA TTTTACAGC 300
 20 AGCACAGCAT TCCTCACCTG CATTGCCGTT GATCGGTATT TGGCTGTTGT CTACCCTTTG 360
 AAGTTTTTTT TCCTAAGGAC AAGAAGATTT GCACTCATGG TCAGCCTGTC CATCTGGATA 420
 TTGGAAACCA TCTTCAATGC TGTCATGTTG TGGGAAGATG AAACAGTTGT TGAATATTGC 480
 GATGCCGAAA AGTCTAATTT TACTTTATGC TATGACAAAT ACCCTTTAGA GAAATGGCAA 540
 ATCAACCTCA ACTTGTTTCTG GACGTGTACA GGCTATGCAA TACCTTTGGT CACCATCCTG 600
 25 ATCTGTAAAC GGAAAGTCTA CCAAGCTGTG CGGCACAATA AAGCCACGGA AAACAAGGAA 660
 AAGAAGAGAA TCAAAAACT ACTTGTCAGC ATCACAGTTA CTTTGTCTT ATGCTTTACT 720
 CCCTTTCATG TGATGTTGCT GATTGCTGTC ATTTTAGAGC ATGCTGTGAA CTTCGAAGAC 780
 CACAGCAATT CTGGGAAGCG AACTTACACA ATGTATAGAA TCACGGTTGC ATTAACAAGT 840
 TTAAATTGTG TTGCTGATCC AATTCTGTAC TGTTTTGTGA CCGAAACAGG AAGATATGAT 900
 30 ATGTGGAATA TATTAAATT CTGCACTGGG AGGTGTAATA CATCACAAAG ACAAAGAAAA 960
 CGCATACTTT CTGTGTCTAC AAAAGATACT ATGGAATTAG AGGTCCTTGA GTAG 1014

- 103 -

(135) INFORMATION FOR SEQ ID NO:134:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 337 amino acids
 (B) TYPE: amino acid
 (C) STRANDEDNESS:
 (D) TOPOLOGY: not relevant

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:134:

10	Met	Asn	Ser	Thr	Cys	Ile	Glu	Glu	Gln	His	Asp	Leu	Asp	His	Tyr	Leu	1	5	10	15
	Phe	Pro	Ile	Val	Tyr	Ile	Phe	Val	Ile	Ile	Val	Ser	Ile	Pro	Ala	Asn	20	25	30	
	Ile	Gly	Ser	Leu	Cys	Val	Ser	Phe	Leu	Gln	Ala	Lys	Lys	Glu	Ser	Glu	35	40	45	
15	Leu	Gly	Ile	Tyr	Leu	Phe	Ser	Leu	Ser	Leu	Ser	Asp	Leu	Leu	Tyr	Ala	50	55	60	
	Leu	Thr	Leu	Pro	Leu	Trp	Ile	Asp	Tyr	Thr	Trp	Asn	Lys	Asp	Asn	Trp	65	70	75	80
20	Thr	Phe	Ser	Pro	Ala	Leu	Cys	Lys	Gly	Ser	Ala	Phe	Leu	Met	Tyr	Met	85	90	95	
	Asn	Phe	Tyr	Ser	Ser	Thr	Ala	Phe	Leu	Thr	Cys	Ile	Ala	Val	Asp	Arg	100	105	110	
	Tyr	Leu	Ala	Val	Val	Tyr	Pro	Leu	Lys	Phe	Phe	Phe	Leu	Arg	Thr	Arg	115	120	125	
25	Arg	Phe	Ala	Leu	Met	Val	Ser	Leu	Ser	Ile	Trp	Ile	Leu	Glu	Thr	Ile	130	135	140	
	Phe	Asn	Ala	Val	Met	Leu	Trp	Glu	Asp	Glu	Thr	Val	Val	Glu	Tyr	Cys	145	150	155	160
30	Asp	Ala	Glu	Lys	Ser	Asn	Phe	Thr	Leu	Cys	Tyr	Asp	Lys	Tyr	Pro	Leu	165	170	175	
	Glu	Lys	Trp	Gln	Ile	Asn	Leu	Asn	Leu	Phe	Arg	Thr	Cys	Thr	Gly	Tyr	180	185	190	
	Ala	Ile	Pro	Leu	Val	Thr	Ile	Leu	Ile	Cys	Asn	Arg	Lys	Val	Tyr	Gln	195	200	205	
35	Ala	Val	Arg	His	Asn	Lys	Ala	Thr	Glu	Asn	Lys	Glu	Lys	Lys	Arg	Ile	210	215	220	

- 104 -

Lys Lys Leu Leu Val Ser Ile Thr Val Thr Phe Val Leu Cys Phe Thr
 225 230 235 240
 Pro Phe His Val Met Leu Leu Ile Arg Cys Ile Leu Glu His Ala Val
 245 250 255
 5 Asn Phe Glu Asp His Ser Asn Ser Gly Lys Arg Thr Tyr Thr Met Tyr
 260 265 270
 Arg Ile Thr Val Ala Leu Thr Ser Leu Asn Cys Val Ala Asp Pro Ile
 275 280 285
 10 Leu Tyr Cys Phe Val Thr Glu Thr Gly Arg Tyr Asp Met Trp Asn Ile
 290 295 300
 Leu Lys Phe Cys Thr Gly Arg Cys Asn Thr Ser Gln Arg Gln Arg Lys
 305 310 315 320
 Arg Ile Leu Ser Val Ser Thr Lys Asp Thr Met Glu Leu Glu Val Leu
 325 330 335
 15 Glu

(136) INFORMATION FOR SEQ ID NO:135:

(i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 999 base pairs
 20 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:135:

25 ATGGTGAACT CCACCCACCG TGGGATGCAC ACTTCTCTGC ACCTCTGGAA CCGCAGCAGT
 60
 TACAGACTGC ACAGCAATGC CAGTGAGTCC CTTGGAAAAG GCTACTCTGA TGGAGGGTGC
 120
 TACGAGCAAC TTTTGTCTC TCCTGAGGTG TTTGTGACTC TGGGTGTCAT CAGCTTGTTG
 30 180
 GAGAATATCT TAGTGATTGT GGCAATAGCC AAGAACAAGA ATCTGCATT CACCATGTAC
 240
 TTTTTCATCT GCAGCTTGGC TGTGGCTGAT ATGCTGGTGA GCGTTTCAA TGGATCAGAA
 300
 35 ACCATTATCA TCACCCTATT AAACAGTACA GATACGGATG CACAGAGTTT CACAGTGAAT
 360

- 105 -

(137) INFORMATION FOR SEO ID NO:136:

(ii) MOLECULE TYPE: protein

35 Lys Gly Tyr Ser Asp Gly Gly Cys Tyr Glu Gln Leu Phe Val Ser Pro.
 35 40 45

- 106 -

	Glu	Val	Phe	Val	Thr	Leu	Gly	Val	Ile	Ser	Leu	Leu	Glu	Asn	Ile	Leu	
	50						55					60					
	Val	Ile	Val	Ala	Ile	Ala	Lys	Asn	Lys	Asn	Leu	His	Ser	Pro	Met	Tyr	
	65					70					75					80	
5	Phe	Phe	Ile	Cys	Ser	Leu	Ala	Val	Ala	Asp	Met	Leu	Val	Ser	Val	Ser	
				85						90					95		
	Asn	Gly	Ser	Glu	Thr	Ile	Ile	Ile	Thr	Leu	Leu	Asn	Ser	Thr	Asp	Thr	
				100					105					110			
10	Asp	Ala	Gln	Ser	Phe	Thr	Val	Asn	Ile	Asp	Asn	Val	Ile	Asp	Ser	Val	
			115					120					125				
	Ile	Cys	Ser	Ser	Leu	Leu	Ala	Ser	Ile	Cys	Ser	Leu	Leu	Ser	Ile	Ala	
		130					135					140					
	Val	Asp	Arg	Tyr	Phe	Thr	Ile	Phe	Tyr	Ala	Leu	Gln	Tyr	His	Asn	Ile	
	145					150					155					160	
15	Met	Thr	Val	Lys	Arg	Val	Gly	Ile	Ser	Ile	Ser	Cys	Ile	Trp	Ala	Ala	
				165						170					175		
	Cys	Thr	Val	Ser	Gly	Ile	Leu	Phe	Ile	Ile	Tyr	Ser	Asp	Ser	Ser	Ala	
				180					185					190			
20	Val	Ile	Ile	Cys	Leu	Ile	Thr	Met	Phe	Phe	Thr	Met	Leu	Ala	Leu	Met	
			195					200					205				
	Ala	Ser	Leu	Tyr	Val	His	Met	Phe	Leu	Met	Ala	Arg	Leu	His	Ile	Lys	
		210					215					220					
	Arg	Ile	Ala	Val	Leu	Pro	Gly	Thr	Gly	Ala	Ile	Arg	Gln	Gly	Ala	Asn	
	225				230						235					240	
25	Met	Lys	Gly	Lys	Ile	Thr	Leu	Thr	Ile	Leu	Ile	Gly	Val	Phe	Val	Val	
				245						250					255		
	Cys	Trp	Ala	Pro	Phe	Phe	Leu	His	Leu	Ile	Phe	Tyr	Ile	Ser	Cys	Pro	
			260					265						270			
30	Gln	Asn	Pro	Tyr	Cys	Val	Cys	Phe	Met	Ser	His	Phe	Asn	Leu	Tyr	Leu	
			275					280					285				
	Ile	Leu	Ile	Met	Cys	Asn	Ser	Ile	Ile	Asp	Pro	Leu	Ile	Tyr	Ala	Leu	
		290					295					300					
	Arg	Ser	Gln	Glu	Leu	Arg	Lys	Thr	Phe	Lys	Glu	Ile	Ile	Cys	Cys	Tyr	
	305				310						315					320	
35	Pro	Leu	Gly	Gly	Leu	Cys	Asp	Leu	Ser	Ser	Arg	Tyr					
				325						330							

(138) INFORMATION FOR SEQ ID NO:137:

- 107 -

- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 33 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

5

(ii) MOLECULE TYPE: DNA (genomic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:137:

GCCAATATGA AGGGAATAAT TACCTTGACC ATC
33

10 (137) INFORMATION FOR SEQ ID NO:138:

- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 31 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

15

(ii) MOLECULE TYPE: DNA (genomic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:138:

CTCCTTCGGT CCTCCTATCG TTGTCAGAAG T
31

20 (140) INFORMATION FOR SEQ ID NO:139:

- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 1842 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

25

(ii) MOLECULE TYPE: DNA (genomic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:139:

ATGGGGCCCA CCCTAGCGGT TCCCACCCCC TATGGCTGTA TTGGCTGTAA GCTACCCAG 60
CCAGAATACC CACCGGCTCT AATCATCTTT ATGTTCTGCG CGATGGTTAT CACCATCGTT 120
30 GTAGACCTAA TCGGCAACTC CATGGTCATT TTGGCTGTGA CGAAGAACAA GAAGCTCCGG 180
AATTCTGGCA ACATCTTCGT GGTCACTCTC TCTGTGGCCG ATATGCTGGT GGCCATCTAC 240
CCATACCCTT TGATGCTGCA TGCCATGTCC ATTGGGGGCT GGGATCTGAG CCAGTTACAG 300
TGCCAGATGG TCGGGTTCAT CACAGGGCTG AGTGTGGTCG GCTCCATCTT CAACATCGTG 360

- 108 -

GCAATCGCTA TCAACCGTTA CTGCTACATC TGCCACAGCC TCCAGTACGA ACGGATCTTC 420
 AGTGTGCGCA ATACCTGCAT CTACCTGGTC ATCACCTGGA TCATGACCGT CCTGGCTGTC 480
 CTGCCCCAACA TGTACATTGG CACCATCGAG TACGATCCTC GCACCTACAC CTGCATCTTC 540
 AACTATCTGA ACAACCCTGT CTTCACTGTT ACCATCGTCT GCATCCACTT CGTCCTCCCT 600
 5 CTCCTCATCG TGGGTTTCTG CTACGTGAGG ATCTGGACCA AAGTGCTGGC GGCCCGTGAC 660
 CCTGCAGGGC AGAATCCTGA CAACCAACTT GCTGAGGTTC GCAATTTTCT AACCATGTTT 720
 GTGATCTTCC TCCTCTTTGC AGTGTGCTGG TGCCCTATCA ACGTGCTCAC TGTCTTGGTG 780
 GCTGTCAGTC CGAAGGAGAT GGCAGGCAAG ATCCCCAACT GGCTTTATCT TGCAGCCTAC 840
 TTCATAGCCT ACTTCAACAG CTGCCTCAAC GCTGTGATCT ACGGGCTCCT CAATGAGAAT 900
 10 TTCCGAAGAG AATACTGGAC CATCTTCCAT GCTATGCGGC ACCCTATCAT ATTCTTCCCT 960
 GGCCTCATCA GTGATATTCG TGAGATGCAG GAGGCCCGTA CCCTGGCCCG CGCCCGTGCC 1020
 CATGCTCGCG ACCAAGCTCG TGAACAAGAC CGTGCCCATG CCTGTCCTGC TGTGGAGGAA 1080
 ACCCCGATGA ATGTCCGGAA TGTTCATTA CCTGGTGATG CTGCAGCTGG CCACCCGAC 1140
 CGTGCCTCTG GCCACCCTAA GCCCCATTCC AGATCCTCCT CTGCCTATCG CAAATCTGCC 1200
 15 TCTACCCACC ACAAGTCTGT CTTTAGCCAC TCCAAGGCTG CCTCTGGTCA CCTCAAGCCT 1260
 GTCTCTGGCC ACTCCAAGCC TGCCCTCTGGT CACCCCAAGT CTGCCACTGT CTACCCTAAG 1320
 CCTGCCTCTG TCCATTTCOA GGGTGACTCT GTCCATTTCOA AGGGTGACTC TGTCCATTTC 1380
 AAGCCTGACT CTGTTTCATTT CAAGCCTGCT TCCAGCAACC CCAAGCCCAT CACTGGCCAC 1440
 CATGTCTCTG CTGGCAGCCA CTCCAAGTCT GCCTTCAGTG CTGCCACCAG CCACCCTAAA 1500
 20 CCCATCAAGC CAGCTACCAG CCATGCTGAG CCCACCACTG CTGACTATCC CAAGCCTGCC 1560
 ACTACCAGCC ACCCTAAGCC CGCTGCTGCT GACAACCCTG AGCTCTCTGC CTCCCATTCG 1620
 CCCGAGATCC CTGCCATTGC CCACCCTGTG TCTGACGACA GTGACCTCCC TGAGTCGGCC 1680
 TCTAGCCCTG CCGCTGGGCC CACCAAGCCT GCTGCCAGCC AGCTGGAGTC TGACACCATC 1740
 GCTGACCTTC CTGACCCTAC TGTAGTCACT ACCAGTACCA ATGATTACCA TGATGTCGTG 1800
 25 GTTGTGATG TTGAAGATGA TCCTGATGAA ATGGCTGTGT GA 1842

(141) INFORMATION FOR SEQ ID NO:140:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 613 amino acids

(B) TYPE: amino acid

- 109 -

(C) STRANDEDNESS:

(D) TOPOLOGY: not relevant

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:140:

5 Met Gly Pro Thr Leu Ala Val Pro Thr Pro Tyr Gly Cys Ile Gly Cys
 1 5 10 15
 Lys Leu Pro Gln Pro Glu Tyr Pro Pro Ala Leu Ile Ile Phe Met Phe
 20 25 30
 Cys Ala Met Val Ile Thr Ile Val Val Asp Leu Ile Gly Asn Ser Met
 10 35 40 45
 Val Ile Leu Ala Val Thr Lys Asn Lys Lys Leu Arg Asn Ser Gly Asn
 50 55 60
 Ile Phe Val Val Ser Leu Ser Val Ala Asp Met Leu Val Ala Ile Tyr
 65 70 75 80
 15 Pro Tyr Pro Leu Met Leu His Ala Met Ser Ile Gly Gly Trp Asp Leu
 85 90 95
 Ser Gln Leu Gln Cys Gln Met Val Gly Phe Ile Thr Gly Leu Ser Val
 100 105 110
 Val Gly Ser Ile Phe Asn Ile Val Ala Ile Ala Ile Asn Arg Tyr Cys
 115 120 125
 Tyr Ile Cys His Ser Leu Gln Tyr Glu Arg Ile Phe Ser Val Arg Asn
 130 135 140
 Thr Cys Ile Tyr Leu Val Ile Thr Trp Ile Met Thr Val Leu Ala Val
 145 150 155 160
 25 Leu Pro Asn Met Tyr Ile Gly Thr Ile Glu Tyr Asp Pro Arg Thr Tyr
 165 170 175
 Thr Cys Ile Phe Asn Tyr Leu Asn Asn Pro Val Phe Thr Val Thr Ile
 180 185 190
 Val Cys Ile His Phe Val Leu Pro Leu Leu Ile Val Gly Phe Cys Tyr
 195 200 205
 Val Arg Ile Trp Thr Lys Val Leu Ala Ala Arg Asp Pro Ala Gly Gln
 210 215 220
 Asn Pro Asp Asn Gln Leu Ala Glu Val Arg Asn Phe Leu Thr Met Phe
 225 230 235 240
 35 Val Ile Phe Leu Leu Phe Ala Val Cys Trp Cys Pro Ile Asn Val Leu
 245 250 255

- 110 -

	Thr	Val	Leu	Val	Ala	Val	Ser	Pro	Lys	Glu	Met	Ala	Gly	Lys	Ile	Pro	
				260					265					270			
	Asn	Trp	Leu	Tyr	Leu	Ala	Ala	Tyr	Phe	Ile	Ala	Tyr	Phe	Asn	Ser	Cys	
			275					280					285				
5	Leu	Asn	Ala	Val	Ile	Tyr	Gly	Leu	Leu	Asn	Glu	Asn	Phe	Arg	Arg	Glu	
		290					295					300					
	Tyr	Trp	Thr	Ile	Phe	His	Ala	Met	Arg	His	Pro	Ile	Ile	Phe	Phe	Pro	
	305					310					315					320	
	Gly	Leu	Ile	Ser	Asp	Ile	Arg	Glu	Met	Gln	Glu	Ala	Arg	Thr	Leu	Ala	
10					325					330					335		
	Arg	Ala	Arg	Ala	His	Ala	Arg	Asp	Gln	Ala	Arg	Glu	Gln	Asp	Arg	Ala	
				340					345					350			
	His	Ala	Cys	Pro	Ala	Val	Glu	Glu	Thr	Pro	Met	Asn	Val	Arg	Asn	Val	
			355					360					365				
15	Pro	Leu	Pro	Gly	Asp	Ala	Ala	Ala	Gly	His	Pro	Asp	Arg	Ala	Ser	Gly	
		370					375					380					
	His	Pro	Lys	Pro	His	Ser	Arg	Ser	Ser	Ser	Ala	Tyr	Arg	Lys	Ser	Ala	
	385					390					395					400	
	Ser	Thr	His	His	Lys	Ser	Val	Phe	Ser	His	Ser	Lys	Ala	Ala	Ser	Gly	
20					405					410					415		
	His	Leu	Lys	Pro	Val	Ser	Gly	His	Ser	Lys	Pro	Ala	Ser	Gly	His	Pro	
				420					425					430			
	Lys	Ser	Ala	Thr	Val	Tyr	Pro	Lys	Pro	Ala	Ser	Val	His	Phe	Lys	Gly	
			435					440					445				
25	Asp	Ser	Val	His	Phe	Lys	Gly	Asp	Ser	Val	His	Phe	Lys	Pro	Asp	Ser	
		450					455					460					
	Val	His	Phe	Lys	Pro	Ala	Ser	Ser	Asn	Pro	Lys	Pro	Ile	Thr	Gly	His	
	465					470					475					480	
	His	Val	Ser	Ala	Gly	Ser	His	Ser	Lys	Ser	Ala	Phe	Ser	Ala	Ala	Thr	
30					485					490					495		
	Ser	His	Pro	Lys	Pro	Ile	Lys	Pro	Ala	Thr	Ser	His	Ala	Glu	Pro	Thr	
				500					505					510			
	Thr	Ala	Asp	Tyr	Pro	Lys	Pro	Ala	Thr	Thr	Ser	His	Pro	Lys	Pro	Ala	
			515					520					525				
35	Ala	Ala	Asp	Asn	Pro	Glu	Leu	Ser	Ala	Ser	His	Cys	Pro	Glu	Ile	Pro	
		530					535					540					
	Ala	Ile	Ala	His	Pro	Val	Ser	Asp	Asp	Ser	Asp	Leu	Pro	Glu	Ser	Ala	

- 111 -

[illegible]

10 (142) INFORMATION FOR SEQ ID NO:141:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 1842 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:141:

	ATGGGGCCCCA	CCCTAGCGGT	TCCCACCCCC	TATGGCTGTA	TTGGCTGTAA	GCTACCCCAG	60
	CCAGAATACC	CACCGGCTCT	AATCATCTTT	ATGTTCTGCG	CGATGGTTAT	CACCATCGTT	120
20	GTAGACCTAA	TCGGCAACTC	CATGGTCATT	TTGGCTGTGA	CGAAGAACAA	GAAGCTCCGG	180
	AATTCTGGCA	ACATCTTCGT	GGTCAGTCTC	TCTGTGGCCG	ATATGCTGGT	GGCCATCTAC	240
	CCATACCCTT	TGATGCTGCA	TGCCATGTCC	ATTGGGGGCT	GGGATCTGAG	CCAGTTACAG	300
	TGCCAGATGG	TCGGGTTTCA	CACAGGGCTG	AGTGTGGTCG	GCTCCATCTT	CAACATCGTG	360
	GCAATCGCTA	TCAACCGTTA	CTGCTACATC	TGCCACAGCC	TCCAGTACGA	ACGGATCTTC	420
25	AGTGTGCGCA	ATACCTGCAT	CTACCTGGTC	ATCACCTGGA	TCATGACCGT	CCTGGCTGTC	480
	CTGCCCAACA	TGTACATTGG	CACCATCGAG	TACGATCCTC	GCACCTACAC	CTGCATCTTC	540
	AACTATCTGA	ACAACCCTGT	CTTCACTGTT	ACCATCGTCT	GCATCCACTT	CGTCCTCCCT	600
	CTCCTCATCG	TGGGTTTCTG	CTACGTGAGG	ATCTGGACCA	AAGTGCTGGC	GGCCCGTGAC	660
	CCTGCAGGGC	AGAATCCTGA	CAACCAACTT	GCTGAGGTTC	GCAATAAACT	AACCATGTTT	720
30	GTGATCTTCC	TCCTCTTTGC	AGTGTGCTGG	TGCCCTATCA	ACGTGCTCAC	TGTCTTGGTG	780
	GCTGTCAATC	CGAAGGAGAT	GGCAGGCAAG	ATCCCCAACT	GGCTTTTATCT	TGCAGCCTAC	840

- 112 -

TTCATAGCCT ACTTCAACAG CTGCCTCAAC GCTGTGATCT ACGGGCTCCT CAATGAGAAT 900
 TTCCGAAGAG AATACTGGAC CATCTTCCAT GCTATGCGGC ACCCTATCAT ATTCTTCTCT 960
 GGCCTCATCA GTGATATTCG TGAGATGCAG GAGGCCCCGTA CCCTGGCCCCG CGCCCGTGCC 1020
 CATGCTCGCG ACCAAGCTCG TGAACAAGAC CGTGCCCATG CCTGTCCTGC TGTGGAGGAA 1080
 5 ACCCCGATGA ATGTCCGGAA TGTTCATTA CCTGGTGATG CTGCAGCTGG CCACCCCGAC 1140
 CGTGCTCTG GCCACCCTAA GCCCCATTCC AGATCCTCCT CTGCCTATCG CAAATCTGCC 1200
 TCTACCCACC ACAAGTCTGT CTTTAGCCAC TCCAAGGCTG CCTCTGGTCA CCTCAAGCCT 1260
 GTCTCTGGCC ACTCCAAGCC TGCCTCTGGT CACCCCAAGT CTGCCACTGT CTACCCTAAG 1320
 CCTGCCTCTG TCCATTTCAA GGCTGACTCT GTCCATTTCA AGGGTGACTC TGTCCATTTC 1380
 10 AAGCTGACT CTGTTTCAAT CAAGCCTGCT TCCAGCAACC CCAAGCCCAT CACTGGCCAC 1440
 CATGTCTCTG CTGGCAGCCA CTCCAAGTCT GCCTTCAATG CTGCCACCAG CCACCCTAAA 1500
 CCCATCAAGC CAGCTACCAG CCATGCTGAG CCCACCACTG CTGACTATCC CAAGCCTGCC 1560
 ACTACCAGCC ACCCTAAGCC CGCTGCTGCT GACAACCCTG AGCTCTCTGC CTCCCATTGC 1620
 CCCGAGATCC CTGCCATTGC CCACCCTGTG TCTGACGACA GTGACCTCCC TGAGTCGGCC 1680
 15 TCTAGCCCTG CCGCTGGGCC CACCAAGCCT GCTGCCAGCC AGCTGGAGTC TGACACCATC 1740
 GCTGACCTTC CTGACCCTAC TGTAGTCACT ACCAGTACCA ATGATTACCA TGATGTCGTG 1800
 GTTGTGTGATG TTGAAGATGA TCCTGATGAA ATGGCTGTGT GA 1842

(143) INFORMATION FOR SEQ ID NO:142:

- 20 (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 613 amino acids
 (B) TYPE: amino acid
 (C) STRANDEDNESS:
 (D) TOPOLOGY: not relevant

(ii) MOLECULE TYPE: protein

- 25 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:142:

Met Gly Pro Thr Leu Ala Val Pro Thr Pro Tyr Gly Cys Ile Gly Cys
 1 5 10 15

Lys Leu Pro Gln Pro Glu Tyr Pro Pro Ala Leu Ile Ile Phe Met Phe
 20 25 30

30 Cys Ala Met Val Ile Thr Ile Val Val Asp Leu Ile Gly Asn Ser Met
 35 40 45

- 113 -

Val Ile Leu Ala Val Thr Lys Asn Lys Lys Leu Arg Asn Ser Gly Asn
 50 55 60
 Ile Phe Val Val Ser Leu Ser Val Ala Asp Met Leu Val Ala Ile Tyr
 65 70 75 80
 5 Pro Tyr Pro Leu Met Leu His Ala Met Ser Ile Gly Gly Trp Asp Leu
 85 90 95
 Ser Gln Leu Gln Cys Gln Met Val Gly Phe Ile Thr Gly Leu Ser Val
 100 105 110
 10 Val Gly Ser Ile Phe Asn Ile Val Ala Ile Ala Ile Asn Arg Tyr Cys
 115 120 125
 Tyr Ile Cys His Ser Leu Gln Tyr Glu Arg Ile Phe Ser Val Arg Asn
 130 135 140
 Thr Cys Ile Tyr Leu Val Ile Thr Trp Ile Met Thr Val Leu Ala Val
 145 150 155 160
 15 Leu Pro Asn Met Tyr Ile Gly Thr Ile Glu Tyr Asp Pro Arg Thr Tyr
 165 170 175
 Thr Cys Ile Phe Asn Tyr Leu Asn Asn Pro Val Phe Thr Val Thr Ile
 180 185 190
 20 Val Cys Ile His Phe Val Leu Pro Leu Leu Ile Val Gly Phe Cys Tyr
 195 200 205
 Val Arg Ile Trp Thr Lys Val Leu Ala Ala Arg Asp Pro Ala Gly Gln
 210 215 220
 Asn Pro Asp Asn Gln Leu Ala Glu Val Arg Asn Lys Leu Thr Met Phe
 225 230 235 240
 25 Val Ile Phe Leu Leu Phe Ala Val Cys Trp Cys Pro Ile Asn Val Leu
 245 250 255
 Thr Val Leu Val Ala Val Ser Pro Lys Glu Met Ala Gly Lys Ile Pro
 260 265 270
 30 Asn Trp Leu Tyr Leu Ala Ala Tyr Phe Ile Ala Tyr Phe Asn Ser Cys
 275 280 285
 Leu Asn Ala Val Ile Tyr Gly Leu Leu Asn Glu Asn Phe Arg Arg Glu
 290 295 300
 Tyr Trp Thr Ile Phe His Ala Met Arg His Pro Ile Ile Phe Phe Ser
 305 310 315 320
 35 Gly Leu Ile Ser Asp Ile Arg Glu Met Gln Glu Ala Arg Thr Leu Ala
 325 330 335
 Arg Ala Arg Ala His Ala Arg Asp Gln Ala Arg Glu Gln Asp Arg Ala

- 114 -

	340		345		350
	His Ala Cys Pro Ala Val Glu Glu Thr Pro Met Asn Val Arg Asn Val				
	355		360		365
5	Pro Leu Pro Gly Asp Ala Ala Ala Gly His Pro Asp Arg Ala Ser Gly				
	370		375		380
	His Pro Lys Pro His Ser Arg Ser Ser Ser Ala Tyr Arg Lys Ser Ala				
	385		390		395
					400
	Ser Thr His His Lys Ser Val Phe Ser His Ser Lys Ala Ala Ser Gly				
			405		410
					415
10	His Leu Lys Pro Val Ser Gly His Ser Lys Pro Ala Ser Gly His Pro				
			420		425
					430
	Lys Ser Ala Thr Val Tyr Pro Lys Pro Ala Ser Val His Phe Lys Ala				
			435		440
					445
15	Asp Ser Val His Phe Lys Gly Asp Ser Val His Phe Lys Pro Asp Ser				
			450		455
					460
	Val His Phe Lys Pro Ala Ser Ser Asn Pro Lys Pro Ile Thr Gly His				
			465		470
					475
					480
	His Val Ser Ala Gly Ser His Ser Lys Ser Ala Phe Asn Ala Ala Thr				
			485		490
					495
20	Ser His Pro Lys Pro Ile Lys Pro Ala Thr Ser His Ala Glu Pro Thr				
			500		505
					510
	Thr Ala Asp Tyr Pro Lys Pro Ala Thr Thr Ser His Pro Lys Pro Ala				
			515		520
					525
25	Ala Ala Asp Asn Pro Glu Leu Ser Ala Ser His Cys Pro Glu Ile Pro				
			530		535
					540
	Ala Ile Ala His Pro Val Ser Asp Asp Ser Asp Leu Pro Glu Ser Ala				
			545		550
					555
					560
	Ser Ser Pro Ala Ala Gly Pro Thr Lys Pro Ala Ala Ser Gln Leu Glu				
			565		570
					575
30	Ser Asp Thr Ile Ala Asp Leu Pro Asp Pro Thr Val Val Thr Thr Ser				
			580		585
					590
	Thr Asn Asp Tyr His Asp Val Val Val Val Asp Val Glu Asp Asp Pro				
			595		600
					605
35	Asp Glu Met Ala Val				
			610		

(144) INFORMATION FOR SEQ ID NO:143:

- 115 -

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 33 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

5

(ii) MOLECULE TYPE: DNA (genomic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:143:

GCTGAGGTTC GCAATAAACT AACCATGTTT GTG

33

(145) INFORMATION FOR SEQ ID NO:144:

- 10 (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 30 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

15 (ii) MOLECULE TYPE: DNA (genomic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:144:

CTCCTTCGGT CTCCTATCG TTGTCAGAAG T

31

(146) INFORMATION FOR SEQ ID NO:145:

- 20 (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 27 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

25 (iv) ANTI-SENSE: NO

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:145:

TTAGATATCG GGGCCACCC TAGCGGT

33

(147) INFORMATION FOR SEQ ID NO:146:

- 30 (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 29 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

- 116 -

(iv) ANTI-SENSE: YES

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:146:

GGTACCCCCA CAGCCATTTC ATCAGGATC

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
20 April 2000 (20.04.2000)

PCT

(10) International Publication Number
WO 00/22131 A3

(51) International Patent Classification⁷: C12N 15/16,
C07K 14/72

(71) Applicant (for all designated States except US): ARENA
PHARMACEUTICALS, INC. [US/US]; 6166 Nancy
Ridge Drive, San Diego, CA 92121 (US).

(21) International Application Number: PCT/US99/24065

(22) International Filing Date: 13 October 1999 (13.10.1999)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:

09/170,496	13 October 1998 (13.10.1998)	US
60/108,029	12 November 1998 (12.11.1998)	US
60/109,213	20 November 1998 (20.11.1998)	US
60/110,060	27 November 1998 (27.11.1998)	US
60/120,416	16 February 1999 (16.02.1999)	US
60/121,852	26 February 1999 (26.02.1999)	US
60/123,944	12 March 1999 (12.03.1999)	US
60/123,945	12 March 1999 (12.03.1999)	US
60/123,948	12 March 1999 (12.03.1999)	US
60/123,946	12 March 1999 (12.03.1999)	US
60/123,949	12 March 1999 (12.03.1999)	US
60/123,951	12 March 1999 (12.03.1999)	US
60/136,436	28 May 1999 (28.05.1999)	US
60/136,437	28 May 1999 (28.05.1999)	US
60/136,439	28 May 1999 (28.05.1999)	US
60/136,567	28 May 1999 (28.05.1999)	US
60/137,127	28 May 1999 (28.05.1999)	US
60/137,131	28 May 1999 (28.05.1999)	US
60/141,448	30 June 1999 (30.06.1999)	US
60/151,114	27 August 1999 (27.08.1999)	US
60/152,524	3 September 1999 (03.09.1999)	US
60/156,653	29 September 1999 (29.09.1999)	US
60/156,633	29 September 1999 (29.09.1999)	US
60/156,555	29 September 1999 (29.09.1999)	US
60/156,634	29 September 1999 (29.09.1999)	US
60/157,280	1 October 1999 (01.10.1999)	US
60/157,294	1 October 1999 (01.10.1999)	US
60/157,281	1 October 1999 (01.10.1999)	US
60/157,293	1 October 1999 (01.10.1999)	US
60/157,282	1 October 1999 (01.10.1999)	US
09/417,044	12 October 1999 (12.10.1999)	US
09/416,760	12 October 1999 (12.10.1999)	US

(72) Inventors; and

(75) Inventors/Applicants (for US only): BEHAN, Dominic,
P. [GB/US]; 11472 Roxboro Court, San Diego, CA 92131
(US). LEHMANN-BRUINSMA, Karin [DE/US]; 12565
Pathos Lane, San Diego, CA 92129 (US). CHALMERS,
Derek, T. [GB/US]; 347 Longden Lane, Solana Beach,
CA 92150 (US). CHEN, Ruoping [CN/US]; 5296 Timber
Branch Way, San Diego, CA 92130 (US). DANG, Huong,
T. [US/US]; 5352 Oak Park Drive, San Diego, CA 92105
(US). GORE, Martin [GB/US]; 6868 Estrella Avenue,
San Diego, CA 92120 (US). LIAW, Chen, W. [US/US];
7668 Salix Place, San Diego, CA 92129 (US). LIN, I-Lin
[—/US]; 8291-7 Gold Coast Drive, San Diego, CA 92126
(US). LOWITZ, Kevin [US/US]; Apartment C, 8031
Caminito de Pizza, San Diego, CA 92108 (US). WHITE,
Carol [US/US]; 4260 Cleveland Avenue, San Diego, CA
92103 (US).

(74) Agents: MILLER, Suzanne, E. et al.; Woodcock Wash-
burn Kurtz Mackiewicz & Norris LLP, 46th floor, One Lib-
erty Place, Philadelphia, PA 19103 (US).

(81) Designated States (national): AE, AL, AM, AT, AU, AZ,
BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DE, DK,
DM, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL,
IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU,
LV, MA, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT,
RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA,
UG, US, UZ, VN, YU, ZA, ZW.

(84) Designated States (regional): ARIPO patent (GH, GM,
KE, LS, MW, SD, SL, SZ, TZ, UG, ZW), Eurasian patent
(AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent
(AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU,
MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM,
GA, GN, GW, ML, MR, NE, SN, TD, TG).

Published:

— With international search report.

(88) Date of publication of the international search report:
22 February 2001

(63) Related by continuation (CON) or continuation-in-part
(CIP) to earlier application:

US 09/170,496 (CIP)
Filed on 13 October 1998 (13.10.1998)

For two-letter codes and other abbreviations, refer to the "Guid-
ance Notes on Codes and Abbreviations" appearing at the begin-
ning of each regular issue of the PCT Gazette.

(54) Title: NON-ENDOGENOUS, CONSTITUTIVELY ACTIVATED HUMAN G PROTEIN-COUPLED RECEPTORS

(57) Abstract: The invention disclosed in this patent document relates to transmembrane receptors, more particularly to a human G protein-coupled receptor for which the endogenous ligand is unknown ("orphan GPCR receptors"), and most particularly to mutated (non-endogenous) versions of the human GPCRs for evidence of constitutive activity.

WO 00/22131 A3

INTERNATIONAL SEARCH REPORT

International Application No
PCT/US 99/24065

A. CLASSIFICATION OF SUBJECT MATTER
IPC 7 C12N15/16 C07K14/72

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)
IPC 7 C12N C07K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	WO 97 21731 A (NEW ENGLAND MEDICAL CENTER INC) 19 June 1997 (1997-06-19) page 18, line 16-- line 26 figures 2,3	1-4
A	SCHEER A. ET AL.: "CONSTITUTIVELY ACTIVE G PROTEIN-COUPLED RECEPTORS: POTENTIAL MECHANISMS OF RECEPTOR ACTIVATION" JOURNAL OF RECEPTOR AND SIGNAL TRANSDUCTION RESEARCH, vol. 17, no. 1/03, 1997, pages 57-73, XP000867531 ISSN: 1079-9893 the whole document	1-4
	-/-	

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents :

- *A* document defining the general state of the art which is not considered to be of particular relevance
- *E* earlier document but published on or after the international filing date
- *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- *O* document referring to an oral disclosure, use, exhibition or other means
- *P* document published prior to the international filing date but later than the priority date claimed

T later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

X document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

Y document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

Z document member of the same patent family

Date of the actual completion of the international search

2 March 2000

Date of mailing of the international search report

14 06. 2000

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax (+31-70) 340-3016

Authorized officer

Mandl, B

INTERNATIONAL SEARCH REPORT

International Application No
PCT/US 99/24065

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	WO 98 38217 A (HERRICK DAVIS KATHARINE ;TEITLER MILT (US); EGAN CHRISTINA C. (US)) 3 September 1998 (1998-09-03) figure 4	1-4
A	KJELSBORG M. A. ET AL.: "CONSTITUTIVE ACTIVATION OF THE ALPHA1B-ADRENERGIC RECEPTOR BY ALL AMINO ACID SUBSTITUTIONS AT A SINGLE SITE" JOURNAL OF BIOLOGICAL CHEMISTRY, vol. 267, no. 3, 25 January 1992 (1992-01-25), pages 1430-1433, XP002911764 ISSN: 0021-9258 the whole document	1-4
P,A	PAUWELS P. J. ET AL.: "REVIEW:AMINO ACID DOMAINS INVOLVED IN CONSTITUTIVE ACTIVATION OF G-PROTEIN-COUPLED RECEPTORS" MOLECULAR NEUROBIOLOGY, vol. 17, no. 1/03, 1998, pages 109-135, XP000866477 ISSN: 0893-7648 the whole document	1-4
P,A	WO 99 24569 A (ONO PHARMACEUTICAL CO ;HAGA HISANORI (JP); NAKADE SHINJI (JP); FUK) 20 May 1999 (1999-05-20) SEQ.IDs. 1-3	1-4

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US 99/24065

Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☐ Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:
2. ☐ Claims Nos.:
because they relate to parts of the International Application that do not comply with the prescribed requirements to such an extent that no meaningful International Search can be carried out, specifically:
3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this International Search Report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☒ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

1-4

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☐ No protest accompanied the payment of additional search fees.

FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

1. Claims: 1-4

A cDNA encoding a non-endogenous, constitutively activated version of a human G-protein-coupled receptor comprising hARE-3(F313K); the receptor encoded by said cDNA; a plasmid comprising said cDNA; and a host cell comprising said plasmid.

2. Claims: 5-8

A cDNA encoding a non-endogenous, constitutively activated version of a human G-protein-coupled receptor comprising hARE-4(V233K); the receptor encoded by said cDNA; a plasmid comprising said cDNA; and a host cell comprising said plasmid.

3. Claims: 9-12

A cDNA encoding a non-endogenous, constitutively activated version of a human G-protein-coupled receptor comprising hARE-5(A240K); the receptor encoded by said cDNA; a plasmid comprising said cDNA; and a host cell comprising said plasmid.

4. Claims: 13-16

A cDNA encoding a non-endogenous, constitutively activated version of a human G-protein-coupled receptor comprising hGPCR14(L257K); the receptor encoded by said cDNA; a plasmid comprising said cDNA; and a host cell comprising said plasmid.

5. Claims: 17-20

A cDNA encoding a non-endogenous, constitutively activated version of a human G-protein-coupled receptor comprising hGPCR27(C283K); the receptor encoded by said cDNA; a plasmid comprising said cDNA; and a host cell comprising said plasmid.

6. Claims: 21-24

A cDNA encoding a non-endogenous, constitutively activated version of a human G-protein-coupled receptor comprising hARE-1(E232K); the receptor encoded by said cDNA; a plasmid comprising said cDNA; and a host cell comprising said plasmid.

7. Claims: 25-28

FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

A cDNA encoding a non-endogenous, constitutively activated version of a human G-protein-coupled receptor comprising hARE-2(G285K); the receptor encoded by said cDNA; a plasmid comprising said cDNA; and a host cell comprising said plasmid.

8. Claims: 29-32

A cDNA encoding a non-endogenous, constitutively activated version of a human G-protein-coupled receptor comprising hPPR1(L239K); the receptor encoded by said cDNA; a plasmid comprising said cDNA; and a host cell comprising said plasmid.

9. Claims: 33-36

A cDNA encoding a non-endogenous, constitutively activated version of a human G-protein-coupled receptor comprising hG2A(K232A); the receptor encoded by said cDNA; a plasmid comprising said cDNA; and a host cell comprising said plasmid.

10. Claims: 37-40

A cDNA encoding a non-endogenous, constitutively activated version of a human G-protein-coupled receptor comprising hRUP3(L224K); the receptor encoded by said cDNA; a plasmid comprising said cDNA; and a host cell comprising said plasmid.

11. Claims: 41-44

A cDNA encoding a non-endogenous, constitutively activated version of a human G-protein-coupled receptor comprising hRUP5(A236K); the receptor encoded by said cDNA; a plasmid comprising said cDNA; and a host cell comprising said plasmid.

12. Claims: 45-48

A cDNA encoding a non-endogenous, constitutively activated version of a human G-protein-coupled receptor comprising hRUP6(N267K); the receptor encoded by said cDNA; a plasmid comprising said cDNA; and a host cell comprising said plasmid.

13. Claims: 49-52

FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

A cDNA encoding a non-endogenous, constitutively activated version of a human G-protein-coupled receptor comprising hRUP7(A302K); the receptor encoded by said cDNA; a plasmid comprising said cDNA; and a host cell comprising said plasmid.

14. Claims: 53-56

A cDNA encoding a non-endogenous, constitutively activated version of a human G-protein-coupled receptor comprising hCHN4(V236K); the receptor encoded by said cDNA; a plasmid comprising said cDNA; and a host cell comprising said plasmid.

15. Claims: 57-60

A cDNA encoding a non-endogenous, constitutively activated version of a human G-protein-coupled receptor comprising hMC4(A244K); the receptor encoded by said cDNA; a plasmid comprising said cDNA; and a host cell comprising said plasmid.

16. Claims: 61-64

A cDNA encoding a non-endogenous, constitutively activated version of a human G-protein-coupled receptor comprising hCHN3(S284K); the receptor encoded by said cDNA; a plasmid comprising said cDNA; and a host cell comprising said plasmid.

17. Claims: 65-68.

A cDNA encoding a non-endogenous, constitutively activated version of a human G-protein-coupled receptor comprising hCHN6(L352K); the receptor encoded by said cDNA; a plasmid comprising said cDNA; and a host cell comprising said plasmid.

18. Claims: 69-72

A cDNA encoding a non-endogenous, constitutively activated version of a human G-protein-coupled receptor comprising hCHN8(N235K); the receptor encoded by said cDNA; a plasmid comprising said cDNA; and a host cell comprising said plasmid.

19. Claims: 73-76

A cDNA encoding a non-endogenous, constitutively activated

FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

version of a human G-protein-coupled receptor comprising hH9(F236K); the receptor encoded by said cDNA; a plasmid comprising said cDNA; and a host cell comprising said plasmid.

20. Claims: 77-80

A cDNA encoding a non-endogenous, constitutively activated version of a human G-protein-coupled AT1 receptor selected from the group consisting of hAT1(F239K), hAT1(N111A), hAT1(AT2K2551C3) and hAT1 (A243+); the receptor encoded by said cDNA; a plasmid comprising said cDNA; and a host cell comprising said plasmid.

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/US 99/24065

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 9721731 A	19-06-1997	US 5750353 A	12-05-1998
		AU 715611 B	03-02-2000
		AU 1334397 A	03-07-1997
		CA 2239293 A	19-06-1997
		EP 0869975 A	14-10-1998
WO 9838217 A	03-09-1998	AU 6343998 A	18-09-1998
WO 9924569 A	20-05-1999	NONE	